

THE INSTITUTE OF  
TRANSPLANTATION




The Newcastle upon Tyne Hospitals  
NHS Foundation Trust

# Donor infection and the impact on transplantation



Dr Emma Montgomery

Consultant Nephrologist, Freeman Hospital,  
Newcastle Upon Tyne



Donor  
derived  
infection

Organ transplant is a  
low risk but important  
route of transmission  
of infection from  
donor to recipient.



# Agenda

1. Blood born Viruses
  - Hepatitis B
  - Hepatitis C
  - HIV
2. EBV & CMV virus
3. All the others
4. COVID 19



## **MICROBIOLOGICAL SAFETY GUIDELINES**


**PREVIOUSLY KNOWN AS**

**GUIDANCE ON THE MICROBIOLOGICAL SAFETY  
OF HUMAN ORGANS, TISSUES  
AND CELLS USED IN TRANSPLANTATION**

The most  
useful  
resource

What  
are we  
dealing  
with ?

*The degree of risk for  
transmission of infection  
carried with grafts,  
notably of viruses, is  
largely unknown and,  
for a specific organ,  
difficult to assess.*



# What are the consequences..


Days from Transplant to Recognition of Infection					
	0-30 days	31-90 days	91-180 days	>181 days	TOTAL
Viral	11	7	4	8	30
Bacterial	23	1	0	0	24
Fungal	20	5	2	1	27
Mycobacterial	2	2	2	0	6
Parasitic	7	4	6	1	18
TOTAL	63	19	14	10	106
	59%	18%	13%	9%	

# Transmission of Blood born virus

- **Human immunodeficiency virus (HIV)**
- **Hepatitis C virus (HCV)**
- **Hepatitis B virus (HBV)**
- **CMV and EBV**


# Other infections

- Human T-lymphotropic virus-1 (HTLV-1),
- Influenza
- *Treponema pallidum*
- *Mycobacterium tuberculosis*,
- Multidrug-resistant bacteria,
- *Strongyloides stercoralis*,
- *Toxoplasma gondii*
- Hepatitis E
- Transmissible spongiform encephalopathy disease.





## Blood born virus ( Hep B, C & HIV)



Exact rates of transmission are not well understood.

There are certain things from the history however that will give you indications for risk.

# High Risk activity

Pathogens	Behavioral characteristics	Nonbehavioral characteristics
HIV	<ul style="list-style-type: none"> <li>• MSM</li> <li>• IVDU</li> <li>• Noninjection illicit drug use</li> <li>• Multiple sex partners</li> <li>• Sex with partner known to be HIV-infected</li> <li>• Age <math>\leq 18</math> at first sexual intercourse</li> </ul>	<ul style="list-style-type: none"> <li>• STI</li> <li>• Marital status</li> </ul>
HCV	<ul style="list-style-type: none"> <li>• IVDU</li> <li>• Noninjection illicit drug use</li> <li>• Multiple sex partners</li> <li>• Sex worker</li> <li>• Inmates</li> <li>• Age <math>\leq 18</math> at first sexual intercourse</li> <li>• Sex with partner known to be HCV-infected</li> <li>• Sex with an injection drug user</li> <li>• Tattooing performed by a nonprofessional</li> </ul>	<ul style="list-style-type: none"> <li>• Hemodialysis</li> <li>• Receipt of blood transfusion</li> <li>• Signs and symptoms (eg, jaundice, elevated ALT)</li> <li>• STI</li> <li>• Marital status</li> </ul>
HBV	<ul style="list-style-type: none"> <li>• MSM</li> <li>• IVDU</li> <li>• Multiple sex partners</li> </ul>	<ul style="list-style-type: none"> <li>• Hemodialysis</li> <li>• STI</li> <li>• Marital status</li> </ul>


HIV, human immunodeficiency virus; HCV, hepatitis C virus; HBV, hepatitis B virus; MSM, men who have sex with men; IVDU, injection drug use; STI, sexually transmitted infection.

## Required Donor Information


- Treatment received in the illness before donation (including duration and dose of antimicrobial and other drug therapy);
- Vaccination history and immunisation status
- History of receipt of blood, blood components, blood products, tissue or organ graft.
- Previous or current immunosuppression (by disease or drugs) as this may affect the interpretation of test results or the donor's suitability.
- Travel History

# Required Donor Information

- History of contact with animals and other vectors. Transplantation may transmit zoonotic infections.
- History that may have put the donor at increased risk of transmissible spongiform encephalopathies (TSEs).
- History of malignancy, recent infectious disease or exposure to an infectious disease.
- Behavioural history that could have put the donor at risk of transmissible pathogens This will include questions about risk behaviours such as recreational drug use, men who have sex with men (MSM), sex with commercial sex workers, sex with a partner know to have a sexually transmissible disease, acupuncture, tattooing and body piercing.
- Results of any recent microbiological tests should be reviewed.




Twenty-year trends in the percentage of donors with drug overdose (intended or unintended) as a cause of death in Australia and New Zealand (ANZ) compared with the United Kingdom (UK) and United States (data sources: Australia and New Zealand Organ Donation Registry [ANZOD], Organ Procurement and Transplantation Network [OPTN], National Health Service Blood and Transplant [NHSBT]).




Estimated prevalence of human immunodeficiency virus (HIV) and hepatitis C virus (HCV) among people who inject drugs in selected high-income countries. HCV prevalence estimates represent mid-range estimates (source of HCV data: United Nations Office on Drugs and Crime <http://unodc.org>; source of HIV data: UNAIDS [aidsinfo.unaids.org](http://aidsinfo.unaids.org)). \*HCV estimate for Germany represents high range estimate for the year 2011. IDU, intravenous drug users.

# Interpreting donor test results

- Following exposure to, and infection by, a microbiological agent there is a period of time during which no microbe can be readily recovered from the host; this is classically called the eclipse period.
- Donations taken during this period are unlikely to be infectious but in practice this would not be safe and should be avoided.



- This period of infectivity which cannot be detected is colloquially called a “window” and represents the duration of undetectable infectivity.
- This “window” is shortest for genomic (nucleic acid technology testing, NAT) and antigen tests, and longest for antibody tests.





# Window

**Table 1:** Estimates of window period length for different testing methods\*


Pathogen	Standard serology	Enhanced serology (fourth generation or combined antibody-antigen tests)	Nucleic acid testing
HIV	17–22 days (5–8)	~7–16 days (9,10)	5–6 days (5,6)
HCV	~70 days (5,8,11)	~40–50 days (12–14)	3–5 days (5,11)
HBV	35–44 days (15,16)	Not applicable	20–22 days (8,15)

## Interpreting Hepatitis B Blood Test Results

Interpretation & Action Needed	HBsAg Hepatitis B Surface Antigen	HBsAb (anti-HBs) Hepatitis B Surface Antibody	HBcAb (anti-HBc) Hepatitis B Core Antibody
<b>Not Immune - Not Protected</b>  Has not been infected, but still at risk for possible hep B infection.  <b>Vaccine is needed.</b>	—	—	—
<b>*Immune Controlled - Protected</b>  Surface antibodies present due to natural infection. Has recovered from a prior hep B infection. Cannot infect others.  <b>No vaccine is needed.</b>	—	+	+
<b>Immune - Protected</b>  Has been vaccinated. Does not have the virus and has never been infected.  <b>No vaccine is needed.</b>	—	+	—
<b>Infected</b>  Positive HBsAg indicates hep B virus is present. Virus can spread to others. Find a doctor who is knowledgeable about hep B for further evaluation.  <b>More Testing Needed.</b>	+	—	+
<b>*Could be Infected</b>  Result unclear - possible past or current hep B infection. Find a doctor who is knowledgeable about hep B for further evaluation.  <b>More Testing Needed.</b>	—	—	+

\*Inform all doctors about a prior or current hepatitis B infection and include this information as part of your health history.  
Talk to doctors before taking immune system suppressing medications to understand the risk for possible hep B reactivation.

# Hepatitis C tests



# HIV


- Screening for HIV infection must include a combined HIV antigen/antibody assay.
- Samples giving repeat reactivity in antibody or combined antigen/antibody assays must undergo additional testing to confirm HIV infection including nucleic acid tests (NAT) for HIV RNA.
- Confirmed detection of specific anti-HIV 1/2 antibodies and/or HIV RNA and/or HIV antigen indicates current infection.
- The use of organ and cells from HIV-infected individuals may be considered in the setting of HIV-infected recipients.

# CMV and EBV


---

- The majority of adult populations worldwide are latently infected with CMV and/or EBV,
- CMV: 20% to 100%
- EBV: 50% to 90%

of populations older than 18 years, respectively


- 
- CMV and EBV cause lifelong infection, and organs from seropositive donors may transmit infection, potentially causing severe disease in a seronegative recipient.
  - Latent CMV and EBV may also reactivate in immunosuppressed seropositive patients post transplantation. No contraindications exist for organ donation in the case of donors with latent CMV infection, although recipient morbidity increases in the case of D+/R- combinations.

# CMV



# EBV - *Epstein-Barr virus*

- Epstein-Barr virus (EBV) infection is associated with the development of post-transplant lymphoproliferative disorders (PTLDs).
- EBV transmission to a seronegative recipient is the greatest risk factor for PTLD
- EBV positive donors does not prevent transplant but monitoring is recommended





# Human T cell Lymphotropic Virus (HTLV)

---

- The Human T-cell lymphocytic virus-1 (HTLV-1) is an oncogenic retrovirus that preferentially infects CD4+ T-cells.
- Transmission may occur as a result of breast feeding, IV drug use, sexual intercourse or blood transfusion.
- Although infection is usually asymptomatic in most individuals, approximately 2% to 5% of infected individuals will subsequently develop acute T-cell leukaemia/lymphoma (ATL) around 20 to 30 years after infection.
- A smaller proportion (0.25–4%) will develop HTLV-1–associated myelopathy/tropical spastic paraparesis (HAM/TSP) soon after the initial infection.



# Human T cell Lymphotropic Virus (HTLV)

- **The majority of HTLV-1–infected individuals will not develop clinical manifestations of ATL or HAM/TSP in their lifetime.**
  - However, infection with HTLV-1 suppresses immune surveillance and increases susceptibility to other infections including parasitic infection with *Strongyloides stercoralis* and scabies, bacterial infections including *Mycobacterium tuberculosis*, *Mycobacterium leprae*, and infectious dermatitis, and viral infections including HIV, HCV, and HBV.
-

# HTLV

- HTLV is not mandatory for all donors of tissues and cells but is for donors living in, or originating from high-prevalence areas, or with sexual partners originating from those areas or where the donor's parents originate from those areas. There are also requirements for the repeat testing after at least 180 days for those donors at risk of HTLV infection
- Confirmation of specific anti-HTLV antibodies indicates current infection
- The decision to proceed with solid organ transplantation following an initial reactive HTLV antibody test is dependent on an assessment of the net benefit of receiving that transplant when compared to the risk of not receiving that specific transplant.





# Toxoplasma gondii


- Transmission of an infection with *T. gondii* occurs most commonly when a seronegative recipient receives an organ from a seropositive donor. Whereas the occurrence of toxoplasmosis following non-cardiac organ transplantation is low, the reported prevalence in serologically mismatched (D+/R-) heart and heart-lung recipients in the absence of antimicrobial prophylaxis can be as high as 75% due to the transmission of *T. gondii* cysts present within cardiac tissue.
- Less commonly, seropositive recipients may manifest reactivation of latent infection.

If positive

- Donation permitted
  - Informs need for prophylaxis in heart recipients
-

# TB - Tuberculosis


- Donation of organs and tissues is contraindicated from donors with active disease or within the first six months of anti-tuberculosis treatment.
- Previous disease or from a risk area - requires donor prophylaxis.



# Hepatitis E Virus


---

- Hepatitis E virus (HEV) is overall the world's most common cause of acute viral hepatitis.
- If positive
- Donation permitted. Informs post transplant management



## *Strongyloides stercoralis*

- Asymptomatic carriage with strongyloides stercoralis has been reported most often in donors who were both born in and lived for some while in endemic areas which include most of the Tropics and Sub-tropics.
- An Eosinophilia may or may not be present. Transmission to immuno-compromised recipients is often associated with significant morbidity and a high mortality rate.
- Pre-donation identification from stool sampling and serology, most practicable for a live donor allows for effective recipient prophylaxis.







## Syphilis


Treponema  
pallidum (Syphilis)

- Syphilis is never a contraindication for using organs
- Penicillin should be administered to recipients of serologically reactive donors.

Drug resistant  
bacteria e.g.  
methicillin resistant  
Staphylococcus  
aureus (MRSA),  
vancomycin resistant  
Enterococcus (VRE),  
carbapenemase-  
producing  
Enterobacteriaceae  
(CPE)

Drug resistant bacteria can be transmitted from donor to recipient. Transmitted infections are difficult to treat and are associated with poorer outcome in the recipient.

- The presence of drug resistant bacteria in the donor is a relative contraindication to solid organ transplantation.
- Specialist microbiological advice must be sought.
- Careful consideration of benefits from transplant is required.



## Yearly Epidemic Influenza

- UK guidelines state that lungs and bowel should not be used from donors with confirmed influenza infection. Other organs may be offered, and the final decision lies with the transplanting surgeon, weighing the balance of risks for the recipient and noting that pathogenicity of some strains of virus may be enhanced by immunosuppression.

# Meningoencephalitis of unknown cause

- Donors with undiagnosed meningoencephalitis are an uncommon but potentially lethal source of donor-derived infection.
- Transmission of rabies, LCMV, WNV, *Mycobacterium tuberculosis*, *Cryptococcus*, *Coccidioides immitis*, *Aspergillus*, and *Balamuthia* have occurred when donors with meningitis or encephalitis of unknown cause have been used as organ donors.
- For this reason, any meningitis or encephalitis without a proven cause should be an absolute contraindication to transplantation, according to the international guidelines

# Bacterial meningitis

- If bacterial meningitis has been confirmed, and there is no visible damage or local infection in the organ or tissues required at retrieval, the donation of the organs, tissues and cells are acceptable provided appropriate treatment has been administered to the donor.



# Transmissible Spongiform Encephalopathies (TSEs) (Prion )

TSEs (otherwise known as prion diseases) are a group of fatal transmissible neurodegenerative disorders that in humans occur in sporadic, genetic and acquired forms.

The commonest human TSE, Creutzfeldt-Jakob disease, occurs in all three forms:

1. Genetic (gCJD),
2. Sporadic (sCJD)
3. Acquired (Variant CJD,vCJD, and iatrogenic CJD,iCJD).

# Exclusions from organ and/or tissue donation based on possible TSE exposure

	<b>Solid Organ Donors</b>
Definite, probable or possible case of human TSE, including CJD and vCJD	Absolute contra-indication
Individual with a neurological disease of unknown aetiology	Absolute contra-indication
Individual whose blood relatives have had familial CJD <sup>1</sup>	Absolute contra-indication
Individual "presumed infected" with vCJD <sup>2</sup>	Absolute contra-indication
Individual "at increased risk of CJD/vCJD" (for public health purposes) <sup>3</sup>	<b>Individual assessment required <sup>4</sup></b>
History of definite <sup>5</sup> or probable <sup>6</sup> blood transfusion since 1980	<b>Individual assessment required <sup>4</sup></b>
History of receipt of <i>dura mater</i> graft	<b>Individual assessment required <sup>4</sup></b>
History of definite receipt of tissue since 1980	<b>Individual assessment required <sup>4</sup></b>
History of receipt of pituitary derived growth hormone and/or gonadotrophin	<b>Individual assessment required <sup>4</sup></b>
History of receipt of organ	<b>Individual assessment required <sup>4</sup></b>

## Bacterial infection tests

Tests for bacterial infection	
Rapid plasma reagin (RPR) or other serological test for syphilis	Not contraindicated but treat the recipient
Tuberculin skin test	(test recommended only for recipient)
Blood cultures	Not contraindicated but treat the recipient. Individual decision in the case of MDR bacteria.



## Viral infection Tests


Tests for viral infection	
HIV 1/2 antibody	Contraindicated but considered for HIV-positive recipient
Cytomegalovirus IgG antibody	Not contraindicated but essential to define prophylactic strategy after procedure depending on recipient serology
EBV IgG antibody	Not contraindicated but essential to monitor EBV-negative recipients, especially children
HBsAg	Contraindicated but considered for HBsAg+ recipients or HBV protective immunity
HBcAc/'HBc alone'	Not contraindicated but consider antiviral prophylaxis for liver and HBV non-immune recipients
HCV antibody	Contraindicated but considered for HCV+ recipients

## Tests for parasitic infection

Tests for parasitic infection	
Toxoplasma IgG antibody	Not contraindicated but consider prophylaxis for heart transplant

# COVID 19

Flowchart 1 – Excluding Confirmed and Possible COVID-19





# COVID 19

- The organs that have a high expression of hACE2 receptors are the lungs, heart and kidneys
- Organs with high hACE2 expressing cells should be considered high risk for SARS-CoV-2 infection.
- Donor derived COVID 19 in solid organ transplant recipients has never been described.
- Current COVID 19 infection is currently an absolute contraindication but 28 days after full recovery can be considered.

Questions?

