

Pathology

Professor Derek Manas

Professor of HPB and Transplant Surgery

Newcastle University

Associate Medical Director OTDT: Clinical Governance (Retrieval and Transplantation)

Introduction

- Histopathological review is necessary when:
 - a) A suspicion of malignancy identified in a donor
 - i. 'lump' in a potentially transplantable organ
 - ii. Suspected 'cancer' in a hollow viscus
 - b) Further assessment of organ quality is required
 - i. Liver (fat)
 - ii. Kidney (glomerulosclerosis)
- Urgent histopathological analysis can enable the utilisation of donor organs that would otherwise have been discarded

Case 1

- 42 year old male
- 'liked a drink'
- RTA
 - Head injury
 - DBD donor
 - LFTs – Normal
 - U+E – Normal
- 6 Organs accepted:
 - 2 kidneys
 - Heart and lungs
 - Pancreas
 - Liver


Creatinine	73
Amylase	40
Glucose	8
Bilirubin	18
ALT	17
GGT	78
PT	12
U Output	3400mls

Case 1


- Retrieval

- All organs retrieved
- Heart and Lungs – dispatched
- Abdominal organs were retrieved 'En-bloc'
- On the back table – lesion was noted posteriorly in the Right lobe of the liver

- WHAT NEXT?




Case 1

- Options:
 - Ignore – surgeon says its benign
 - Discard all organs
 - Biopsy 



- The pathologist on call that night could not decide if it was malignant or not

Case 1



Hepatocellular Carcinoma



Focal Nodular Hyperplasia

Case 1

- Options:
 - Ignore – surgeon says its benign
 - Discard all organs
 - Biopsy results awaited. . . .



- What about the heart and lungs – already dispatched?

Case 1: FNH (benign)


- FNH is the second most common hepatic lesion and is found at autopsy
 - prevalence of 0.3 – 3 %
- Clinically relevant cases of FNH are rare
 - reported prevalence in US studies of 0.03 %
- Caused by an injury to the portal tract resulting in the formation and enlargement of arterial to venous shunts
 - This causes hyper-perfusion in local arteries resulting in oxidative stress that triggers a response from hepatic stellate cells to produce the **central scar typically seen in cases of FNH**

Types of FNH lesions

- **Traditional**
 - those containing abnormal nodular architecture, malformed appearing vessels and cholangiolar proliferation)
 - most likely to be associated with symptoms.
- **Telangiectatic**
- **Mixed**
- **Atypical forms**
 - less likely to be associated with symptoms.



RC pathology guidelines

 The Royal College of Pathologists
Pathology: the science behind the cure

Clarification of the use of the College publication 'Guidelines on staffing and workload for histopathology and cytopathology departments' in limiting the workload of pathologists

Professor Peter Furness, Director of Professional Standards

In recent months the College and RCPaTH Consulting have undertaken invited reviews of a number of cellular pathology departments and have become aware that the College workload guidance may, on occasions, be misinterpreted and used in an unintended and rigid manner in discussions on working practices between medical staff and managers. The national consultant contract is time-based (programmed activities) and the intention of the College guidance is to indicate an appropriate workload for a period of time. The College acknowledges that there will normally be variations in the amount of work that can be safely delivered by pathologists depending on their experience and the types of work being performed. It is therefore appropriate to be flexible in interpreting the guidance in the interests of good patient care, and not limiting work by a rigid interpretation of a fixed number of points per day.

The College is concerned to be told that its guidance could be used in this unintended and rigid manner. As a result, this clarification is being published and the matter will be emphasised in the next full update of the guidance.

The guidance document states that it is intended to:

- a) support departments of pathology in balancing staffing with the anticipated workload, so as to ensure that a sustainable, high-quality service is provided for the benefit of patients
- b) facilitate equitable distribution of work among pathologists within a department
- c) provide information for job planning.



It goes on to add:

"It is expected that service users and commissioners will find this guidance helpful in predicting the resource implications of changes in demand."


It also states:

"These guidelines are not intended to provide a basis for a 'fee per case' system of payment."

Nothing in these guidelines is intended to alter the nationally agreed terms and conditions of service of consultants or associate specialists, in which time is the basis of remuneration.

 050118 1  INVESTOR IN PEOPLE

Report of the Review of NHS Pathology Services in England
Chaired by Lord Carter of Coles



An Independent Review for the Department of Health


Specialist On-call Rota's


Case 2

- 42 year old male
- 'liked a drink'
- BMI - 36
- RTA
 - Head injury
 - DBD donor
 - LFTs – Abnormal
 - U+E – Normal
- 6 Organs accepted:
 - 2 kidneys
 - Heart and lungs
 - Pancreas
 - Liver


Creatinine	73
Amylase	40
Glucose	8
Bilirubin	28
ALT	17
GGT	378
PT	17
U Output	3400mls

Case 2

- At retrieval:
 - Liver noted to be 'moderately fatty'
- What next:
 - Ignore
 - Biopsy 
 - Contact recipient Centre



Case 2: 16% to 20% Macrovesicular steatosis




There are two forms of Liver 'Graft–Steatosis'

- **Macrovesicular steatosis**

- fat vacuoles occupy most of the hepatocytes cytoplasm and displaces the nucleus peripherally
- associated with ***excessive alcohol, obesity, diabetes and hyperlipidaemia***

- **Microvesicular steatosis**

- fat vacuoles are smaller and have a centrilobular distribution
- associated with ***mitochondrial injury such as acute viral or drug induced injury, sepsis and some metabolic disorders***



Large droplet Macro-VS, small droplet Macro-VS, Micro-VS

Hepatic Steatosis – Outcome after LT

Definition

- < 30% **Mild**
- 30% - 60% **Moderate**
- > 60% **Severe**

Impact of a 'fatty' graft

- Primary non-function
 - No steatosis: < 5%
 - Mild steatosis: 5 %
 - Moderate steatosis: 10-15%
 - Severe steatosis: > 50%

D'Alessandro A. Transplantation 1991

Strasberg SM, Hepatology, 1994
Ploeg R, Transplantation, 1993

MORE SPECIFIC: LARGE DROPLET MacVS, SMALL DROPLET MacVS, MicVS, TOTAL STEATOSIS

Case 2: best utilisation?

- **Recipient**

- 40 year old ALD
 - Abstinent for 2 years
- 5 cm HCC
 - Previously 6.2 cm
 - Down-sized with TACE and RFA
 - Meeting the 'new criteria'
- Been waiting 100 days
 - First offer of a liver


Case 3

- Female donor:
 - 63 years old
 - SAH
 - BMI - 32
 - History of
 - Hypertension and MODM
 - Smoker for 20 years
 - 'liked a drink'
 - Previous breast cancer – stage I – 10yrs before
 - Given the 'all clear'
 - DBD
 - Offer of liver and kidneys

Creatinine	92
Amylase	40
Glucose	14
Bilirubin	18
ALT	47
GGT	118
ICU stay	3days
U Output	1400mls
Po2	10

Case 3

- At retrieval:
 - L kidney was noted to have multiple cysts
- What next:
 - Ignore
 - Biopsy
 - 'De-roof'
 - Contact the recipient Centre



Case 3

- What if the cyst was more complex
- What next:
 - Ignore
 - Biopsy
 - Contact the recipient Centre



Renal cysts

Category	CT features	Significance
Class I	Water density homogenous Noncalcified, smooth margin No enhancing component	Benign
Class II	Thin septae (<1 mm) Thin calcification (<1 mm) Hemorrhagic cyst	Benign
Class IIF		Likely benign Follow-up imaging indicated
Class III	Thick septa Thick calcification Thick wall Multilocular +/- enhancement	≈ 50% malignant
Class IV	Criteria of category III Enhancing solid mass of wall or septa	Definitely malignant



Which one should be biopsied?



Case 4

- 72 year old man
 - Hypertension
 - Diabetes
- Sudden collapse
- Plan to withdraw treatment
- Offered kidneys as a DCD
- NORS surgeon noted significant scarring
 - Both kidneys

Creatinine	102
Amylase	24
Glucose	14
Bilirubin	18
ALT	27
GGT	78
ICU stay	3days
U Output	400mls
Po2	12

Case 4

- At the implanting Centre there were concerns

- Options:

- Ignore and implant
- NMP
- Discard
- Biopsy




REM UZZI SCORE: 8




Case 4

Tubulointerstitial fibrosis



Glomerulosclerosis



Scarred kidney. . .




- With **increasing age** there is an increase in chronic vascular changes, tubulointerstitial scarring and glomerulosclerosis in kidneys
 - The histological extent does not correlate with the serum creatinine
- The only reliable way to determine the extent of scarring is by an **adequately sized biopsy**, that takes in the full thickness of the cortex:
 - to prevent over representation of subcapsular accentuation of glomerulosclerosis
 - includes arcuate arteries more likely to show hypertensive type intimal changes impacting the luminal area
- The use of a biopsy has been shown to increase the utilization of elderly donors with good outcomes

Remuzzi Score (Cambridge modification)


Glomerular sclerosis

- >25 gloms (Karpinski >20) – should be 50-75
- G0 no sclerosed gloms
- G1 1% –<20%
- G2 20%–50%
- G3 >50%
- If 100 gloms
- G1 1GS to 19GS




Tubular atrophy

- We are defining tubular atrophy as < 50% diameter of normal tubule (Banff)
- Percent of cortex involved
- TA0 no atrophic tubules
- TA1 >0 –<20%
Banff cut off 25%
- TA2 20%-50%
- TA3 >50%



Interstitial fibrosis

- % cortex scarred
- IF0 no fibrosis
- IF1 >0 –<20%
Banff IF0 (d0) up to 5%
- IF2 20%-50%
Banff cut off 25%
- IF3 >50%




Vessels

- Cambridge modification
- Do not score arterioles
- Original Remuzzi – worst of artery and arteriole
- If arterioles bad – they will mention in comments and discuss with the surgeon – but not is score.
- WORST ARTERY IN BIOPSY SCORED
- A0 normal artery
- A1 wall thickness < lumen diameter
- A2 wall thickness = or slightly > lumen diameter
- A3 wall thickness >> lumen diameter
Cambridge modification

Remuzzi – wall thickness far exceeds with severe luminal narrowing

REM UZZI VASULAR 1: wall thickness < lumen diameter
wall thickness 108.6 & lumen diameter 243.7



REM UZZI SCORE (Cambridge modification)

(G + TA + IF + A)
0 1 2 3 4 5 6 7 8 9

ADEQUACY

Is the biopsy adequate (≥ 25 glomeruli AND ≥ 3 arteries)

Yes No

OTHER COMMENTS:

REPORTING PATHOLOGIST: Desley Neil

DATE: 22/9/17

TIME: 2.30 PM

Remuzzi score

- G + TA + IF + A (0-12)

CURRENT PRACTICE IN CAMBRIDGE WITH REM UZZI SCORE (CAMBRIDGE MOD)

≤4 Single transplant





5-6 Dual transplant*

≥7 Discard

* "good 5s" with minimal (<5%) glomerular sclerosis, tubular atrophy and interstitial fibrosis

the need

- For those unexpected lesions identified during organ retrieval or at the time of examination of the organ/s at the implanting centre.
 - Histopathological analysis becomes necessary before safe transplantation can proceed.
 - Once its biopsied all is put on hold until there is an outcome
 - At present there is NO FORMAL 'JOINED-UP' process across TRANSPLANTING CENTRES IN THE UK for obtaining 'out-of-hours' histopathological review of retrieved organs
 - Some centers have an on-call
 - Some centers – 'good-will'
 - Some centers have no service




Why is pathological analysis important?

- Risk of malignant lesions is increased with:
 - Increasing age
 - Obesity
 - Excess alcohol
 - Smoking
 - History of previous malignancy (metastases to liver and lungs)
- Further assessment of organ quality is required
 - enable the better utilization of donor organs


Age and BMI

Figure 3.1 Age of deceased donors in the UK, 1 April 2010 - 31 March 2020



37% > 60

Figure 3.2 BMI of deceased donors in the UK, 1 April 2010 - 31 March 2020



30% > 30


Case 5

- Male donor:
 - 18yrs old
 - Attempted suicide by hanging
 - Was found by friends and cut down
 - 'hanging-time' unknown
 - ??? WIT
 - Resuscitated by paramedics
 - No past medical history
 - All his organs offered as a DBD

Creatinine	375 → 101
Amylase	300 → 88
Glucose	4
Bilirubin	18
ALT	702 → 204
Po2	18
ICU stay	5days
U Output	800mls

Case 5

- At retrieval a large para-aortic lymph node mass (2.5 cms) was found
- SNOD attempted to get a histology assessment @ 2am
- No pathologist was available
- NORS surgeon assessed it as benign



Case 5

- What next:
 - Cardiac patient was asleep
 - R kidney was allocated to a recipient for a beneficial match (waited 3 years)
 - Liver was allocated to a Sero-negative hepatitis 27 year old female
 - Pancreas was allocated to an islet patient with severe hypoglycemic unawareness
- 6 organs were discarded and 6 patients missed out
 - 2 died

The National Histopathology Audit (2013/14)

Aim of the audit

1. Define the incidence of 'urgent' histopathological analysis requests
 1. *Urgent biopsies were defined as those biopsies where the report was awaited in order to proceed either for retrieval or for transplantation*
2. Define numbers of retrieved organs utilised following histology
3. Identify impact of an out-of-hours histopathology service on:
 - Donor/organ utilisation
 - Recipient safety

What was done. . .

- **Prospective audit** over a 6 month period
- All NORS team retrievals and All Transplant Centers were involved

- 654 Retrievals
- 2322 Organs Retrieved
 - Utilized: **2064 (88%)**
 - Taken, Accepted and Not Utilized: **258 (12%)**
- *100% Data Returned*


Number of biopsies requested

- **142 urgent biopsies in 654 retrievals (21.7%)**
 - 42 (29.6%) - Suspected Malignancy (Type 1)
 - 100 (70.4%) – Quality (Type 2)


- **51% out-of-hours (1900 to 0700 and weekends)**

- 95% biopsies sent to pathology services at NORS centers
- The organs biopsied at Transplanting Centres were mostly **Livers and Kidneys**
- There was only **one biopsy taken at CT Transplant Centre**
- **NO** biopsies from **Pancreases**

Results – biopsy incidence by age



Results – biopsy incidence by donor type



56% Biopsies were performed on DBD organs
44% Biopsies were performed on DCD organs

Results – Type 1 (potential malignancy) Biopsies

- 42/654 Biopsies for suspected malignancy (6.4%)
 - 3/654 Malignancies identified (0.45%)
 - 3/42 Biopsies confirmed malignancy (7%)
- There were **119 organs** safely transplanted thanks to negative Biopsy report


Results – Type 2 Biopsy

- 100 Type 2 Biopsies
 - 22 liver
 - 78 kidney
- Unsuitable organ quality in 5% (4 kidneys;1 liver)
- 21 Livers and 74 Kidneys utilised following Type 2 Biopsy

Impact of Biopsies (All Donors)

	Number of Organs Retrieved	Number of Organs Used WITHOUT ANY Biopsy (%)	Number of Organs used after Type 1 Bx (%)	Number of Organs used after Type 2 Bx (%)	Total N. Organs Used	Increase Utilisation %
Kidneys	1220	980 (80.3)	69 (5.6)	73 (6)	1122 (92)	11.7
Whole Livers	481	384 (80)	25 (5.2)	19 (4)	428 (89)	9
Pancreases	166	102 (61.4)	5 (3)	N/A	107 (64.4)	3
Hearts	97	87 (89.7)	5 (5.1)	N/A	92 (95)	5.3
Lungs	229	199 (87)	13 (5.6)	N/A	212 (92.5)	5.5
Total	2193	1752 (79.9)	117 (5.5)	92 (4.2)	1961 (89.4)	9.6

Retrospective audit for suspected malignancy: 1/9/19 – 1/3/20



Retrospective audit 19/20: Histopathology undertaken

Organ type	Number
Liver	16
Kidney	15
Pancreas	12
Lung	10
Lymph nodes	8
Ovary	7
Uterus	5
Other	20


Retrospective audit 19/20: Where were they performed

Abdominal NORS centres undertaking histopathology	Number
Leeds	19
Cambridge	15
Kings College	14
Newcastle	11
Royal Free	8
Birmingham	7
Edinburgh	4
Cardiff	2
Non NORS centres	13

Increased vulnerability of 'out of hours' Histopathology from October 2020


Abdominal NORS	24/7 Histopathology Service- Current	Histopathology undertaken September 1 st – February 28th	24/7 Histopathology Service - October	Additional Info
Birmingham	Yes	7	Yes	No formal rota, 2 pathologists will undertake out of hours histopathology, not contracted or paid to do so
Cambridge	Yes	15	Yes	Will only process their own histopathology
Cardiff	Yes	2	Yes	
Edinburgh	No	4	No	Stopped 2 years ago
King's College	Yes	14	Yes	Only Liver and related specimens. No BMS/lab staff on call just Pathologists
Manchester	No	0	No	
Leeds	Yes	19	No	Likely to cease in October 2020 although not confirmed
Newcastle	Yes	11	No	Will cease in October 2020
Royal Free	Yes	8	Yes	
Oxford	No	0	No	

The current process: 'vulnerable'



Glass Slide - Histopathology Processing/ Histopathology Assessment

Advent of digital technology




Digital Pathology


3DHistech scanner



Pannoramic DESK



Slide Scanner - Histopathology Processing/ Histopathology Assessment



PITHIA TRIAL: Slides Scanners currently in 6 centres



• 6 Scanner Centres

- Cambridge
- Royal Free
- Birmingham
- Leeds
- Newcastle
- Edinburgh

Option 1. National Histopathology Assessment Centre/ NORS Histopathology Processing Centre (With Slide Scanner)

Single National Histopathology Assessment Centre

BMS on call at 6 Scanner Centres

6 subspecialty pathologists on call

- Urothelial Kidney
- Liver and Hepatobiliary
- Gynaecology
- Lung
- Gastrointestinal
- Haematological

Single National Histopathology Assessment Centre

BMS on call at 6 Scanner Centres

4 subspecialty pathologists on call

- Urothelial Kidney
- Liver and Hepatobiliary
- Gynaecology
- Lung

Option 2. NORS Histopathology Assessment Centre/ NORS Histopathology Processing Centre (With Slide Scanner)

6 Histopathology Assessment Centres

BMS on call at 6 Scanner Centres

6 subspecialty pathologists on call

- Urothelial Kidney
- Liver and Hepatobiliary
- Gynaecology
- Lung
- Gastrointestinal
- Haematological

6 Histopathology Assessment Centres

BMS on call at 6 Scanner Centres

4 subspecialty pathologists on call

- Urothelial Kidney
- Liver and Hepatobiliary
- Gynaecology
- Lung

Option 3. National Histopathology Assessment Centre/ NORS Histopathology Processing Centre (Slide Scanner) – Informal Rota (Histopathology rota leads)

Single National Histopathology
Assessment Centre

BMS on call at 6 Scanner Centres

6 subspecialty pathologists – Informal
Rota

- Urothelial Kidney
- Liver and Hepatobiliary
- Gynaecology
- Lung
- Gastrointestinal
- Haematological

Single National Histopathology
Assessment Centre

BMS on call at 6 Scanner Centres

4 subspecialty pathologists – Informal
Rota

- Urothelial Kidney
- Liver and Hepatobiliary
- Gynaecology
- Lung

Next steps

- Workforce (Staffing/ standby/ call out)
- Transport requirements
- Distance of travel for samples
- Impact on SNODs/ HUB/ BMS/ Pathologists
- Cost of equipment (initial, recurring)

History of cancer in a potential organ donor

Original article

Estimated risk of cancer transmission from organ donor to graft recipient in a national transplantation registry

R. Desai¹, D. Collett¹, C. J. E. Watson², P. Johnson³, T. Evans⁴ and J. Neuberger¹

¹NHS Blood and Transplant, Bristol, ²University Department of Surgery and Cambridge National Institute for Health Research Biomedical Campus, Addenbrooke's Hospital, Cambridge, and ³School of Cancer Sciences, University of Birmingham, and ⁴Public Health England, Birmingham, UK
Correspondence to: Dr R. Desai, NHS Blood and Transplant, Fox Den Road, Stoke Gifford, Bristol BS34 8RR, UK (e-mail: rajeev.desai@nhs.net)

Background: Transplanted organs carry the risk of inadvertent donor cancer transmission. Some cancers in organ donors have been classified as being associated with a high or unacceptable risk, but the evidence for such recommendations is scanty.

Methods: The risk of cancer transmission from donors characterized as high or unacceptable risk was studied by analysing transplant and cancer registry data. Donors and recipients from England (1990–2008) were identified from the UK Transplant Registry. Cancer details were obtained from cancer registries and classified using guidelines from the Council of Europe and Organ Procurement and Transplantation Network/United Network for Organ Sharing.

Results: Of 17 639 donors, 202 (1.1 per cent) had a history of cancer, including 61 donors with cancers classed as having an unacceptable/high risk of transmission. No cancer transmission was noted in 133 recipients of organs from these 61 donors. At 10 years after transplantation, the additional survival benefit gained by transplanting organs from donors with unacceptable/high-risk cancer was 944 (95 per cent confidence interval (c.i.) 851 to 1037) life-years, with a mean survival of 7.1 (95 per cent c.i. 6.4 to 7.8) years per recipient.

Conclusion: Strict implementation of present guidelines is likely to result in overestimation of cancer transmission risk in some donors. Organs from some donors with cancers defined as unacceptable/high risk can be used safely.

Paper accepted 16 January 2014

Published online 28 April 2014 in Wiley Online Library (www.bjs.co.uk). DOI: 10.1002/bjs.9460

Transplantation Reviews 28 (2014) 169–175

Contents lists available at ScienceDirect

Transplantation Reviews

journal homepage: www.elsevier.com/locate/trre

Organ transplantation from donors (cadaveric or living) with a history of malignancy: Review of the literature

Sheng Zhang¹, Jin Yuan¹, Wei Li¹, Qifa Ye^{*,1}

¹Third Xiangya Hospital of Central South University, Changsha, China

ABSTRACT

The evolution of organ transplantation has resulted in extended lifespan as well as better life quality of patients with end-stage diseases, which in turn causes an increased demand for organs. The persistent organ shortage requires a careful reconsideration of potential donors (living or cadaveric) that have current or historical malignancies. Donors with low-grade skin tumors, carcinomas in situ of the uterine cervix, and primary central nervous system (CNS) tumors can be considered as potential donors for recipients dying on wait list long for organ transplantation. Recently, transplant centers have turned to other types of malignancies including low grade renal cell carcinoma, prostate, uterine, endometrial and breast cancer, and favorable outcomes have been shown in such innovations. When considering donors with a history of malignancy, general biologic behavior of the tumor type, histology and stage at the time of diagnosis, and the length of disease-free interval should be considered (Transplantation 2002;74(12):1657–1663). With the review of literatures, we illustrate the organ utilization from donors with malignancies all around the world since earlier times and give some suggestions for decision making under the circumstance of whether to choose those marginal donors or not on the basis of reviewed literatures.
© 2014 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

1. Introduction

The persistent shortage of organ supplies is a major obstacle to carry out organ transplantation for the large number of people waiting on the list. Both the size of the candidate waiting list and the number of deaths on the waiting list are progressively increasing [1]. The disparity between organ demand and organ supply has never been moderated. In order to decrease the mortality on the waiting list, transplant centers make every effort to increase the number of donors. Thus, utilization of extended criteria donors (ECD) has been suggested [2]. In the early 2000s, the concept of extended criteria kidney donor was defined to older individuals with hypertension, diabetes, or renal dysfunction, who were expected to produce allografts at greater risk of graft loss than standard donors, albeit sufficiently adequate for transplantation [3]. While the definition of extended criteria liver donor was characterized by individuals with advanced age, steatotic livers, donation after cardiac death (DCD), livers with seropositivity for hepatitis B virus (HBV) and hepatitis C virus (HCV). Besides, occult malignancies become a part of extended criteria donor factors [4]. Using organs from donors with malignancy is not uncommon, and it has played an important role in expanding the donor pool. Though

this may carry risk of malignancy transmission, the risk of tumor transmission or donor related death is extremely small when compared with the benefits of organ transplantation.

2. Review of the literature

Buell et al. [5] reviewed all cases reported to the Israel Penn International Transplant Tumor Registry (PIPTR) that demonstrated a potential for donor-transmitted malignancy from the year 1905 to 2003. 296 cases of high-risk transplants performed using donors with known or incidentally discovered malignancies were reviewed. From the overall series, 124 cases (42%) were identified with confirmed donor transmission. Among them, CNS tumor, malignant melanoma, choriocarcinoma, renal cell carcinoma (RCC), lung cancer, colon cancer, and breast cancer were discovered with donor malignancy transmission. The transmission rate ranges from 23% to 93%. This study showed a relatively higher rate of tumor transmission, given that the donors might have high grade malignancies or misdiagnosed CNS tumors, which carry a much higher transmission risk.

Later, an analysis of Organ Procurement and Transplantation Network/United Network for Organ Sharing (OPTN/UNOS) data on 39,455 deceased donors from 2000 to 2005 showed 1069 donors had a past history of malignancy, resulting in 2508 transplants. The most common type of previous cancer in the donor was nonmelanoma skin cancer (n = 776) followed by central nervous system malignancies (n = 642) and carcinoma of the uterine cervix (n = 336). Four recipients died from donor transmitted malignancy. However, these

* Corresponding author at: Third Xiangya Hospital, No 152 Tongzipo Road, Changsha City, Hunan Province, P.R. China, 410013. Tel.: +86 15116313340; fax: +86 7318948395.
E-mail address: yeqifaj@163.com (Q. Ye).

¹ No conflict of interest.

History of cancer in a potential organ donor

- 61 donors donated 140 organs
- 133 recipients, comprising a total of
 - 86 Kidneys
 - 22 Livers
 - 10 Hearts
 - 8 Lungs
 - 7 multiple organs
 - (4 kidney–pancreas, 2 heart–lung and 1 kidney–heart).
- Comparison of the survival of recipients of single organs from donors with an unacceptable/high risk and standard/non-standard risk of cancer transmission revealed no significant difference in unadjusted survival or risk-adjusted hazard of death

- At 10 years after transplantation, the additional survival benefit of transplanting the organs from donors with an unacceptable/high risk of cancer transmission was 944 (95 per cent C.I. 851 to 1037) life-years, with a mean survival of 7.1 (95 per cent C.I. 6.4 to 7.8) years per recipient.
- 8 of these recipients developed post-transplant cancers, but none had the same cancer type as their donor, indicating these were likely to be de novo cancers

Recipient group	Transplants from donors with an unacceptable/high risk of cancer transmission			Transplants from donors with a standard/non-standard risk of cancer transmission			Risk-adjusted hazard of death for recipients from donors with unacceptable/high risk‡		
	n	Mean age (years)	Recipient survival (years)	n	Mean age (years)	Recipient survival (years)	P†	Hazard ratio	P
Kidney	86	47.4 (43.7, 51.0)	8.79 (3.80, –)*	23 994	42.6 (42.4, 42.8)	10.96 (10.69, 11.27)	0.522	0.87 (0.55, 1.39)	0.566
Liver	22	41.2 (32.6, 49.9)	5.37 (0.11, –)*	6560	39.4 (39.0, 39.8)	4.86 (4.43, 5.42)	0.807	1.07 (0.43, 2.64)	0.884
Heart	10	34.3 (22.8, 45.8)	3.75 (0.01, –)*	2720	32.2 (31.7, 32.7)	3.56 (2.72, 4.17)	0.686	0.73 (0.17, 3.18)	0.670
Lung	8	39.0 (28.1, 49.9)	0.43 (0.04, 5.94)	1245	36.6 (35.8, 37.3)	0.94 (0.70, 1.29)	0.400	2.85 (0.94, 8.62)	0.063
Pancreas	0	–	–	149	32.7 (30.7, 34.6)	6.20 (5.84, 10.32)	–	–	–

Bad for transplantation. . .

Soldier died after being given smoker's lungs in transplant

Cancer developed and drugs prescribed by
Papworth hospital served to speed up disease, coroner hears

[The Guardian](#), Monday 12 October 2009

Transplant patients given kidneys from donor with cancer

Investigation under way into how two transplant patients
were given kidneys from a donor with a rare form of lymphoma

•[The Guardian](#), Tuesday 22 March 2011

A kidney operation changed Robert's life. He got cancer

Father of four was assured stringent tests
had been carried out on the organ he was receiving – but there was one vital flaw

•[The Guardian](#), Tuesday 22 March 2011

Cancer Spreads from Organ Donor to 4 People in 'Extraordinary' Case

By [Rachael Rettner](#) September 15, 2018



(Image: © Shutterstock)

It's well known that organ transplants can pass infectious diseases from donors to recipients in rare cases. But even more rarely, transplants can transmit cancer, as a new case shows.

Donor Malignancy Transmission Risk Assessment

Table 2: Suggested risk categorizations for specific tumor types¹

Risk category	Tumors
No significant risk	Benign tumors in which malignancy is excluded (see Table 3 and Supporting Table S4)
Minimal risk (<0.1% transmission)	Basal cell carcinoma, skin Squamous cell carcinoma, skin without metastases Carcinoma <i>in situ</i> , skin (nonmelanoma) <i>In situ</i> cervical carcinoma <i>In situ</i> vocal cord carcinoma Superficial (noninvasive) papillary carcinoma of bladder (T0N0M0 by TNM stage) (nonrenal transplant only) ⁵
Low risk (0.1–1% transmission)	Solitary papillary thyroid carcinoma, ≤0.5 cm Minimally invasive follicular carcinoma, thyroid, ≤ 1.0 cm (Resected) solitary renal cell carcinoma, ≤1.0 cm, well differentiated (Fuhrman 1–2) ⁴ (Resected) solitary renal cell carcinoma, > 1.0 cm ≤2.5 cm, well differentiated (Fuhrman 1–2) ⁴ Low grade CNS tumor (WHO grade I or II) Primary CNS mature teratoma Solitary papillary thyroid carcinoma, 0.5–2.0 cm Minimally invasive follicular carcinoma, thyroid, 1.0–2.0 cm History of treated non-CNS malignancy (≥5 years prior) with >99% probability of cure
Intermediate risk (1–10% transmission)	Breast carcinoma (stage 0 i.e. carcinoma <i>in situ</i>) Colon carcinoma (stage 0 i.e. carcinoma <i>in situ</i>) (Resected) solitary renal cell carcinoma T1b (4–7 cm) well differentiated (Fuhrman 1–2) stage I ^{4,6} History of treated non-CNS malignancy (≥5 years prior) with probability of cure between 90–99%
High risk (>10% transmission)	Malignant melanoma Breast carcinoma >stage 0 (active) ² Colon carcinoma >stage 0 (active) ² Choriocarcinoma CNS tumor (any) with ventriculooperitoneal or ventriculoatrial shunt, surgery (other than uncomplicated biopsy), irradiation or extra-CNS metastasis CNS Tumor WHO grade III or IV (see Supporting Table S3) ⁷ Leukemia or lymphoma History of melanoma, leukemia or lymphoma, small cell lung/neuroendocrine carcinoma Any other history of treated non-CNS malignancy either (a) insufficient follow-up to predict behavior, (b) considered incurable or (c) with probability of cure <90% Metastatic carcinoma Sarcoma Lung cancer (stages I–IV) ⁶ Renal cell carcinoma >7 cm or stage II–IV ⁶ Small cell/neuroendocrine carcinoma, any site of origin Active cancer not listed elsewhere ³

Nalesnik MA et al, 2011

Discussion

- Histopathology is vital to improve donor characterization
 - Donors are now much older and the risk of malignancy is significantly higher
 - Risk averse practices can be reduced when histopathology analysis is available
 - Organ utilization can be improved
-
- There is a recognition that there is a need for organ specific pathological analysis
 - Need for sustainability
 - Need for a robust service