



Blood and Transplant

ANNUAL REPORT ON PANCREAS AND ISLET TRANSPLANTATION

**REPORT FOR 2021/2022
(1 APRIL 2012 – 31 MARCH 2022)**

PUBLISHED DECEMBER 2022

PRODUCED IN COLLABORATION WITH NHS ENGLAND



Contents

1	Executive Summary	4
2	Introduction	6
	PANCREAS	12
3	Pancreas transplant list	12
3.1	Patients on the pancreas transplant list as at 31 March, 2013 – 2022	13
3.2	Post-registration outcomes, 1 April 2018 – 31 March 2019.....	15
3.3	Demographic characteristics, 1 April 2021 – 31 March 2022	16
3.4	Patient waiting times for those currently on the list, 31 March 2022	19
3.5	Median active waiting time to transplant, 1 April 2016 - 31 March 2020	19
4	Response to pancreas offers.....	21
4.1	Offer decline rates, 1 April 2019 – 31 March 2022.....	22
5	Pancreas transplants.....	25
5.1	Pancreas transplants, 1 April 2012 – 31 March 2022	26
5.2	Demographic characteristics, 1 April 2021 - 31 March 2022.....	29
6	Pancreas outcomes.....	36
6.1	Deceased donor graft and patient survival for first SPK transplant	37
6.2	Deceased donor graft and patient survival for first PO transplants	40
7	Survival from listing	42
7.1	Patient survival from listing for SPK transplant	43
8	Form return rates.....	45
8.1	Pancreas form return rates, 1 January – 31 December 2021	46
	ISLET.....	48
9	Islet transplant list.....	48
9.1	Patients on the islet transplant list as at 31 March, 2013 – 2022	49
9.2	Post-registration outcomes, 1 April 2018 – 31 March 2019.....	51
9.3	Demographic characteristics, 1 April 2021 – 31 March 2022	52
9.4	Patient waiting times for those currently on the list, 31 March 2022	53
9.5	Median active waiting time to transplant, 1 April 2016 - 31 March 2020	53
10	Response to islet offers	55
10.1	Offer decline rates, 1 April 2019 – 31 March 2022.....	56
11	Islet transplants	57
11.1	Islet transplants, 1 April 2012 – 31 March 2022	58
11.2	Demographic characteristics, 1 April 2021 - 31 March 2022.....	61
12	Islet outcomes	63
12.1	Outcome measures for routine islet transplants.....	64

13 Form return rates.....	70
13.1 Islet form return rates, 1 January – 31 December 2021	71
APPENDIX.....	72
A1 Glossary of terms	73
A2 Methods.....	78

Executive Summary

This report presents key figures about pancreas and islet transplantation in the UK. The period reported covers ten years of pancreas and islet transplant data, from 1 April 2012. The report presents information on the number of transplants and survival analysis after first simultaneous pancreas and kidney and pancreas only transplantation on a national and centre-specific basis. Also reported on a national basis is survival analysis after islet transplantation and additional outcome measures.

Key findings

- On the 31 March 2022, there were 278 patients on the UK active pancreas and islet [transplant list](#), which represents a 3% increase in number of patients a year earlier. The number of patients on the active pancreas list increased by 65% to 252 in 2022 and the active islet [transplant list](#) increased by 37% to 26 patients in the same time period.
- There were 1736 pancreas transplants performed in the UK in the ten year period and 269 islet transplants performed in the same time period. The number of transplants from [donations after brain death](#) has increased by 45% in the last year to 112. The number of transplants from [donations after circulatory death](#) has increased by 71% in the last year to 41.
- The national rates of [patient](#) survival one- and five-years after first simultaneous pancreas and kidney transplant from deceased donors are 98% and 91%, respectively. These rates vary between centres, ranging from 95% to 100% at one-year and 84% to 96% at five-years. All centre rates are [risk-adjusted](#).
- The national rates of [graft](#) survival one- and five-years after first simultaneous pancreas and kidney transplant from deceased donors are 94% and 82%, respectively. These rates vary between centres, ranging from 89% to 100% at one-year and 75% to 93% at five-years. All centre rates are [risk-adjusted](#).
- The national rates of [patient](#) survival one- and five-years after first pancreas only transplant from deceased donors are 100% and 88%, respectively. The national rates of [graft](#) survival at one- and five-years are 87% and 66%. Centre specific estimates of these rates must be interpreted with caution due to the small number of transplants upon which they are based.
- The national rate of ten-year [patient](#) survival from listing for deceased donor simultaneous pancreas and kidney transplant is 77%. The rates were similar at centres, ranging from 77% to 78%. All centre rates are [risk-adjusted](#).
- The national rates of one- and five-years [graft](#) survival for patients receiving a first routine islet transplant are 76% and 57%. For patients with a functioning graft at one-year post-transplant, the national rate of five year [graft](#) survival was 73% for patients receiving an additional priority islet graft and 57% for patients who did not.
- Reductions in annual rate of severe [hypoglycaemic](#) events, median [HbA1c](#) and median insulin requirements have been reported at one-year post routine islet transplant.

Use of the contents of this report should be acknowledged as follows:

Annual Report on Pancreas and Islet Transplantation 2021/22, NHS Blood and Transplant.

Introduction

This report presents information on pancreas and islet transplant activity between 1 April 2012 and 31 March 2022, for all eight centres performing pancreas transplantation and seven centres performing islet transplantation in the UK. Cambridge, Cardiff, Guy's and WLRTC only perform pancreas transplants while Bristol, King's College and the Royal Free only perform islet transplants. Throughout this report West London Renal and Transplant Centre is labeled as WLRTC, simultaneous pancreas and kidney transplants and simultaneous islet and kidney transplants are reported as SPK and SIK transplants respectively.

Data were obtained from the UK Transplant Registry, at NHS Blood & Transplant, that holds information relating to donors, recipients and outcomes for all pancreas and islet transplants performed in the UK. [Graft](#) and [patient](#) pancreas survival estimates at one-year post-transplant for the period 1 April 2017 to 31 March 2021 and five-year post-transplant for the period 1 April 2013 to 31 March 2017, will be presented at a later date due to insufficient follow-up at time of analysis.

Islet transplant survival is measured by four key variables: graft survival, and a reduction in [HbA1c](#), insulin requirements and the annual rate of severe [hypoglycaemic](#) events. Islet outcomes at one-year post-transplant for the period 1 April 2017 to 31 March 2021, and [graft](#) survival at five-year post-transplant for the period 1 April 2012 to 31 March 2021, for the national cohort only, will be presented at a later date due to insufficient follow-up at time of analysis. Islet outcomes are [unadjusted](#) for risk and islet outcome data from the UK Transplant Registry is supplemented by data collected from the UK Islet Transplant Consortium.

Pancreas [patient](#) survival from listing is reported at one, five and ten years post registration for a deceased donor simultaneous pancreas and kidney transplants between 1 January 2009 and 31 December 2021.

The centre specific results for survival estimates are adjusted for differences in [risk factors](#) between the centres. The risk models and methods used are described in the Appendix.

Patients requiring [multi-organ transplants](#) (except simultaneous pancreas and kidney or islets and kidney transplants (SPK and SIK)) are excluded from all analyses apart from the introduction. All results are described separately for pancreas and islet transplant recipients other than those presented in this introduction section. Intestinal transplants that involve a pancreas are excluded from all sections of the report.

The COVID-19 pandemic has led to unprecedented challenges for UK transplantation. Concerns about the ability to care for transplant recipients, lack of access to resource because it is being used for patients in the pandemic, and the risk versus benefit for immunosuppressed transplant recipients, have resulted in a major reduction in the number of organ transplants undertaken.

Waiting list figures at the 31 March 2020 and 2021 do not accurately reflect the need for pancreas and islet transplantation due to the COVID-19 pandemic. In 2020, different practices were established across the UK with regards to waiting list management.

Figure 2.1 shows the number of patients on the pancreas and islet [transplant list](#) at 31 March each year between 2013 and 2022. The number of patients actively waiting for a pancreas or islet transplant has increased slightly by 3% from 271 in 2013 to 278 in 2022.




Figure 2.2 shows the number of patients on the pancreas and islet [transplant list](#) at 31 March 2022 for each transplant centre. Manchester has the largest [transplant list](#) with 83 patients registered for a pancreas or islet transplant. Of these patients, 70 are registered for a SPK, three for a pancreas only and four for an islet only transplant. Edinburgh, Manchester and Oxford have patients waiting for an SIK transplant, eight in total. There were no patients on the active islet list at Bristol, King's College or The Royal Free at 31 March 2022.




Figure 2.3 shows the total number of pancreas and islet transplants performed in the last ten financial years. Transplant numbers decreased gradually from 246 in 2013/14 to 203 in 2019/20 and then halved to 101 in 2020/21 due to the COVID-19 pandemic. In 2021/22 transplants numbers increased to 153 transplants. In particular, the number of pancreas only transplants decreased from 37 transplants in 2012/13 to 11 in 2021/22.




Figure 2.4 shows the total number of pancreas and islet transplants performed in 2021/22 at each transplant centre. Oxford performed the most pancreas and islet transplants last year, a total of 51 transplants, whilst Edinburgh performed the most islet and SIK transplants (13). A total of seven SIK transplants were performed at Manchester, Edinburgh and Oxford. King's College, The Royal Free and Bristol performed no transplants during this time period and Bristol no longer offer an islet service.





Figure 2.5 details the 153 pancreas and islet transplants performed in the UK between 1 April 2021 and 31 March 2022. Data for transplants performed in 2020/21 are also presented. The overall number of whole pancreas transplants performed in 2021/22 has increased by 45 compared with 2020/21 to 131. The number of islet transplants has increased by seven compared with 2020/21 to 22.

**Figure 2.5 Pancreas and islet transplants performed in the UK,
1 April 2021 – 31 March 2022 (1 April 2020 – 31 March 2021)**



Geographical variation in registration and transplant rates

Figure 2.6 shows rates of registration to the pancreas and islet transplant list per million population (pmp) between 1 April 2021 and 31 March 2022 compared with pancreas and islet transplant rates pmp for the same time period, by recipient country/NHS region of residence. **Table 2.2** shows the breakdown of these numbers by recipient country/NHS region of residence. No adjustments have been made for potential demographic differences in populations. If a patient has had more than one registration/transplant in the period, each registration/transplant is considered. Note that this analysis only considered NHS Group 1 patients.

Since there will inevitable be some random variation in rates between areas, the [systematic component of variation](#) (SCV) was used to identify if the variation is more (or less) than a random effect for the different NHS regions in England only. Only first registrations and transplants in this period were considered. The larger the SCV the greater the evidence of a high level of systematic variation between areas. Registration and transplant rates yielded an SCV of 0.0172 (p-value = 0.098) and 0 (p-value = 0.999), respectively. The p-value shows the probability that an SCV of this size (or higher) would be observed by chance if only random variation existed and therefore, no evidence of geographical variation beyond what would be expected at random. No adjustment has been made for area-specific demographic characteristics that may impact the rates of registration to the transplant list and transplantation such as age and sex. Therefore, these results should be interpreted with caution.

Figure 2.6 Comparison of pancreas registration rates (pmp) with transplant rates (pmp) by recipient country/NHS region of residence




Table 2.1 Pancreas and islet registration and transplant rates per million population (pmp) in the UK, 1 April 2021 - 31 March 2022, by Country/NHS region

Country/NHS region	Registrations (pmp)		Transplants (pmp)	
North East and Yorkshire	35	(4.1)	19	(2.2)
North West	43	(6.1)	13	(1.8)
Midlands	45	(4.2)	23	(2.2)
East of England	27	(4.1)	13	(2)
London	31	(3.4)	17	(1.9)
South East	27	(3)	25	(2.8)
South West	21	(3.7)	12	(2.1)
England	229	(4)	122	(2.2)
Isle of Man	0	-	0	-
Channel Islands	1	(5.9)	0	-
Wales	18	(5.7)	10	(3.2)
Scotland	43	(7.9)	17	(3.1)
Northern Ireland	4	(2.1)	4	(2.1)
TOTAL	298	(4.4)	153¹	(2.3)

¹ Registrations include 3 recipient whose postcode was unknown

Pancreas transplant list

3.1 Patients on the pancreas transplant list as at 31 March, 2013 – 2022

Figure 3.1 shows the number of patients on the pancreas [transplant list](#) at 31 March each year from 2013. The number of patients actively waiting for a pancreas transplant was the highest at 252 in 2022 an increase of 21% from 208 in 2019 prior to the COVID-19 pandemic.




Figure 3.2 shows the number of patients on the active pancreas [transplant list](#) at 31 March 2022 by centre. Manchester had the largest proportion of the [transplant list](#) (29%), closely followed by Oxford with 27%.





Figure 3.3 shows the number of patients on the pancreas [transplant list](#) at 31 March each year from 2013 by transplant centre. The number of patients actively waiting for a pancreas transplant at Manchester has increased in the last three years.



3.2 Post-registration outcomes, 1 April 2018 – 31 March 2019

An indication of outcomes for patients listed for a pancreas transplant is summarised in **Figure 3.4**. This shows the proportion of patients transplanted or still waiting one and three years after joining the list. It also shows the proportion removed from the [transplant list](#) (typically because they become too unwell for transplant) and who died while on the [transplant list](#).

35% of patients registered between 1 April 2018 and 31 March 2019 were transplanted within one year, while three years after listing 73% of patients had received a transplant. There were 2% of patients who had died waiting for a transplant within one year of listing and 6% within three years of listing. It is important to note that the three-year period after registration for these patients included two years impacted by the COVID-19 pandemic.





Figure 3.5 shows the proportion of patients transplanted or still waiting three years after joining the list by centre. Three years after listing, Cambridge had transplanted 88% of their patients while Guy's had transplanted 56% and 8% had died waiting.



3.3 Demographic characteristics, 1 April 2021 – 31 March 2022

The sex, ethnicity, age group, [sensitisation](#) group ([cRF%](#)) and [matchability points score](#) group of patients registered on the pancreas [transplant list](#) in 2021/22 are shown by centre and overall for the UK in **Figures 3.6, 3.7, 3.8, 3.9** and **3.10** respectively. Note that all percentages quoted are based only on data where relevant information was available.

Overall, 261 patients were registered on the pancreas transplant list, 250 (96%) were waiting for a SPK transplant. Of these 250, 58% were male, 89% were White, the median age was 40 years and the median [cRF](#) was 0%.

Of the 11 (4%) patients on the pancreas only transplant list, 27% were male, all were White, the median age was 37 years and the median [cRF](#) was 0%.




Figure 3.7 Registrations on the active pancreas transplant list, 1 April 2021 - 31 March 2022, by centre and ethnicity




Figure 3.8 Registrations on the active pancreas transplant list, 1 April 2021 - 31 March 2022, by centre and age group




Figure 3.9 Registrations on the active pancreas transplant list, 1 April 2021 - 31 March 2022, by centre and sensitisation group (cRF%)






Figure 3.10 Registrations on the active pancreas transplant list, 1 April 2021 - 31 March 2022, by centre and matchability group



3.4 Patient waiting times for those currently on the list, 31 March 2022

Figure 3.11 shows the length of time active patients have been waiting on the pancreas [transplant list](#) at 31 March 2022 by centre. The majority of patients currently listed have been waiting less than one year. However, three highly sensitised ([cRF](#) 100%) patients have been waiting more than 7 years for a pancreas transplant: one SPK at WLRTC, one SPK at Manchester and one pancreas alone patient at Oxford.



3.5 Median active waiting time to transplant, 1 April 2016 - 31 March 2020

The length of time a patient waits for a pancreas transplant varies across the UK. The [median](#) active waiting time for deceased donor pancreas transplantation is calculated using the [Kaplan-Meier method](#) and is shown in **Figure 3.12** and **Table 3.1** for patients registered at each individual centre.

The [median](#) active waiting time to transplant for patients registered on the pancreas [transplant list](#) between 1 April 2016 and 31 March 2020 is 359 days. This ranged from 167 days at Cambridge to 546 days at Edinburgh.

Figure 3.12 Median active waiting time to deceased donor transplant for patients registered on the pancreas transplant list, 1 April 2016 - 31 March 2020




Table 3.1 Median active waiting time to pancreas transplant in the UK, for patients registered 1 April 2016 - 31 March 2020

Transplant centre	Number of patients registered	Waiting time (days)	
		Median	95% Confidence interval
Cambridge	88	167	128 - 206
Cardiff	46	393	276 - 510
Edinburgh	93	546	486 - 606
Guy's	134	374	300 - 448
Manchester	173	272	196 - 348
Newcastle	36	430	202 - 658
Oxford	286	386	352 - 420
WLRTC	54	322	142 - 502
UK	910	359	330 - 388

Response to pancreas offers

4.1 Offer decline rates, 1 April 2019 – 31 March 2022

Pancreas offers from [DBD](#) and [DCD](#) donors whose pancreas was retrieved, offered directly on behalf of a named individual patient and resulted in transplantation were analysed separately. Any offers of pancreases declined for transplantation, pancreases offered for [multi-organ](#) or small bowel transplant were excluded, as were offers made through the fast track scheme or the reallocation of the pancreas.

[Funnel plots](#) are used to compare centre specific offer decline rates and indicate how consistent the rates of the individual transplant centres are with the national rate. Person [case mix](#) is known to influence the number of offers a centre may receive. In this analysis however, only individual offers for named patients were considered which excluded any [ABO](#)- and [HLA](#)-incompatible patients. For this reason, it was decided not to risk adjust for known centre differences in person [case mix](#).

Figure 4.1 compares individual centre offer [DBD](#) decline rates with the national rate over the time period, 1 April 2019 and 31 March 2022. Centres can be identified by the information shown in **Table 4.1**. Cambridge had offer decline rates significantly better than the national rate and was the only centre to remain open during 2020/21. Edinburgh had a higher decline rate than the national average.




Table 4.1 compares individual centre [DBD](#) offer decline rates over time by financial year. The overall offer decline rate increased from 40% in 2020/21 to 66% in 2021/22.

Centre	Code	2019/20 N (%)	2020/21 N (%)	2021/22 N (%)	Overall N (%)
Cambridge	A	6 (67)	11 (0)	3 (33)	20 (25)
Cardiff	B	19 (68)	6 (50)	7 (71)	32 (66)
Edinburgh	C	54 (63)	16 (56)	41 (88)	111 (71)
Guy's	D	39 (49)	3 (33)	32 (59)	74 (53)
Manchester	E	41 (51)	19 (63)	46 (65)	106 (59)
Newcastle	F	10 (40)	4 (75)	10 (50)	24 (50)
Oxford	G	98 (62)	28 (25)	79 (59)	205 (56)
WLRTC	H	13 (38)	1 (0)	11 (64)	25 (48)
UK		280 (58)	88 (40)	229 (66)	597 (58)

	Centre has reached the upper 99.8% confidence limit
	Centre has reached the upper 95% confidence limit
	Centre has reached the lower 95% confidence limit
	Centre has reached the lower 99.8% confidence limit

Figure 4.2 compares individual centre offer [DCD](#) decline rates with the national rate over the time period, 1 April 2019 and 31 March 2022. Centres can be identified by the information shown in **Table 4.2**.




Table 4.2 compares individual [DCD](#) centre offer decline rates over time by financial year. In 2021/22, Cambridge had an offer decline rate better than the national rate, whilst Newcastle had a much higher decline rate than the national average.

Table 4.2 DCD donor pancreas offer decline rates by transplant centre, 1 April 2019 and 31 March 2022

Centre	Code	2019/20		2020/21		2021/22		Overall	
		N	(%)	N	(%)	N	(%)	N	(%)
Cambridge	A	9	(33)	3	(0)	7	(29)	19	(26)
Cardiff	B	5	(40)	1	(100)	4	(50)	10	(50)
Edinburgh	C	4	(25)	4	(25)	7	(57)	15	(40)
Guy's	D	11	(36)	3	(0)	16	(63)	30	(47)
Manchester	E	15	(53)	6	(67)	10	(40)	31	(52)
Newcastle	F	4	(100)	1	(100)	3	(100)	8	(100)
Oxford	G	18	(50)	4	(50)	21	(62)	43	(56)
WLRTC	H	2	(0)	0	(0)	5	(80)	7	(57)
UK		68	(46)	22	(41)	73	(58)	163	(50)
		Centre has reached the upper 99.8% confidence limit							
		Centre has reached the upper 95% confidence limit							
		Centre has reached the lower 95% confidence limit							
		Centre has reached the lower 99.8% confidence limit							

Pancreas transplants

5.1 Pancreas transplants, 1 April 2012 – 31 March 2022

Figure 5.1 shows the total number of pancreas transplants performed in the last ten financial years, by type of donor. The first [DCD](#) pancreas transplant was performed in 2005/06 and by 2012/13 there were 38 [DCD](#) transplants (19%). The number of [DCD](#) transplants performed reached a peak of 60 in 2014/15 but, within the last two financial years, has dropped to 46 in 2019/20 and 37 in 2021/22 although this still accounts for around a quarter of all pancreas transplants.

In 2013/14 the number of [DBD](#) transplants peaked at 175 (82%), however, this has decreased in the last seven years to 94 [DBD](#) transplants in 2021/22 due to the COVID-19 pandemic.




Figure 5.2 shows the total number of pancreas transplants performed in 2021/22, by centre and type of donor. The same information is presented in **Figure 5.3** but this shows the proportion of [DBD](#) and [DCD](#) transplants performed at each centre. Oxford performed the most [DBD](#) and [DCD](#) transplants (49), however Cambridge had the largest proportion of [DCD](#) transplants (55%). Cardiff, Newcastle and WLRTC performed the lowest number of transplants (six each, including 2 [DCD](#) at Cardiff and 1 [DCD](#) at WLRTC), in the last financial year.

Figure 5.2 Pancreas transplants, 1 April 2021 - 31 March 2022, by centre and donor type




Figure 5.3 Pancreas transplants, 1 April 2021 - 31 March 2022, by type





Figure 5.4 shows the total number of pancreas transplants performed in last ten financial years, by centre and type of donor. Oxford have consistently performed a large number of pancreas transplants including a number of [DCD](#) transplants over the last ten years. However, the number of transplants performed at Oxford has been steadily decreasing over the time period. Edinburgh and Newcastle have not performed many [DCD](#) transplants over the ten year period.



5.2 Demographic characteristics, 1 April 2021 - 31 March 2022

The sex, ethnicity, age group, [sensitisation](#) group ([cRF%](#)) and [matchability points score](#) group of transplant recipients that received a pancreas transplant in 2021/22 are shown by centre in **Figures 5.5, 5.6, 5.7** and **5.8** respectively. Note that all percentages quoted are based only on data where relevant information was available.

Overall, 131 patients were transplanted on the pancreas transplant list, 120 (92%) were SPK transplants. Of which 55% were male, 82% were white, the [median](#) age was 43 years, the [median cRF](#) was 0% and 4% were in the difficult match group. Of the 11 (8%) patients transplanted as a pancreas only transplant, 27% were male, 82% were white, the [median](#) age was 39 years, the [median cRF](#) was 25% and 27% were in the difficult match group.





Figure 5.7 Pancreas transplants, 1 April 2021 - 31 March 2022, by centre and age group




Figure 5.8 Pancreas transplants, 1 April 2021 - 31 March 2022, by centre and sensitisation (cRF%) group





Figure 5.9 Pancreas transplants, 1 April 2021 - 31 March 2022, by centre and matchability group



5.3 Cold ischaemia time, 1 April 2012 – 31 March 2022

Median cold ischaemia times (CIT) are shown in addition to inter-quartile ranges in **Figures 5.10 to 5.15**. Fifty percent of the transplants have a CIT within the inter-quartile range (indicated by a box). Where there is only one observation to report, the single data point is represented by a circle and the median for multiple observations is represented by a line. There is some variation in average (median) CIT between different transplant centres although all centres continually try to reduce this time.

The cold ischaemia times used for all donors, is as reported on the pancreas transplant record form and may include periods of machine perfusion; no adjustment has been made for this.

Figure 5.10 shows the median cold ischaemia time in DBD donor pancreas transplants over the last ten years. During this time period the overall median cold ischaemia time has steadily decreased from 12 hours in 2012/13 to 11 hours in 2021/22.





Figure 5.11 shows the median cold ischaemia time in DBD donor pancreas transplants in 2021/22 for each transplant centre. Please note the small numbers used in the calculations for each centre and interpret with caution. **Figure 5.12** shows the median cold ischaemia time in DBD donor pancreas transplants over the last ten years for each transplant centre.



**Figure 5.12 Median cold ischaemia time in DBD donor pancreas transplants,
1 April 2012 - 31 March 2022**




Figure 5.13 shows the [median](#) cold ischaemia time in [DCD](#) donor pancreas transplants over the last ten years and overall has predominately been 10 hours.




Figure 5.14 shows the [median](#) cold ischaemia time in [DCD](#) donor pancreas transplants in 2021/22 for each transplant centre. Newcastle did not perform any DCD donor transplants in the time period and are not presented in the graph. Please note the small numbers used in the calculations for each centre and interpret with caution.





Figure 5.15 shows the [median](#) cold ischaemia time in [DCD](#) donor pancreas transplants for each transplant centre over the last ten years. The [median](#) cold ischaemia time has fluctuated in centres over the time period.



Pancreas outcomes

6.1 Deceased donor graft and patient survival for first SPK transplant

[Funnel plots](#) are used to compare centre specific [risk-adjusted patient](#) and [graft](#) survival rates and indicate how consistent these rates are with the national survival rates. Note that some patients return to local renal units for follow-up care after their transplant and although survival is reported according to transplant unit, patients may in fact be followed up quite distantly from their transplant centre. It is important to note that adjusting for patient mix through the use of risk-adjustment models may not account for all possible causes of centre differences. There may be other factors that are not taken into account in the risk-adjustment process that may affect the survival rate of a particular centre.

The survival data used for these analyses is reported to NHSBT via follow-up forms and to ensure validity of the survival rates, it is essential these follow-up forms are returned. Follow-up form return rates by centre, for forms issued during the 2021 calendar year, are presented in [Section 8](#).

Figures 6.1 and 6.2 compare individual centre survival estimates with the national rates for one-year [patient](#) and [graft](#) survival for deceased donor first SPK transplants. **Figures 6.3 and 6.4** compare five-year survival estimates. The [funnel plots](#) show that, for the most part, the centres lie within the [confidence limits](#). Some of the [funnel plots](#) show some centres to be above the upper 95% [confidence limit](#). This suggests that these centres may have survival rates that are considerably higher than the national rate. Centres can be identified by the information shown in **Tables 6.1 and 6.2** for patient and graft survival, respectively.




Figure 6.2 Risk-adjusted one year pancreas graft (death censored) survival rates for all deceased donor first SPK transplants, between 1 April 2017 and 31 March 2021





Figure 6.3 Risk-adjusted five year patient survival rates for deceased donor first SPK transplants, between 1 April 2013 and 31 March 2017






Table 6.1 Risk-adjusted one and five year patient survival for first SPK transplants using pancreases from deceased donors

Centre	Code	Patient survival					
		N	One-year*		N	Five-year**	
			%	(95% CI)		%	(95% CI)
Cambridge	A	85	98	(92 - 100)	94	96	(89 - 99)
Cardiff	B	28	100	N/A	24	84	(54 - 97)
Edinburgh	C	63	100	N/A	71	94	(83 - 99)
Guy's	D	85	97	(90 - 100)	104	94	(86 - 98)
Manchester	E	82	100	N/A	111	90	(81 - 95)
Newcastle	F	22	95	(75 - 100)	24	92	(73 - 99)
Oxford	G	160	98	(94 - 100)	205	90	(84 - 94)
WLRTC	H	23	100	N/A	27	87	(63 - 97)
UK		548	98	(97 - 99)	660	91	(89 - 93)

Centre has reached the lower 99.8% confidence limit
 Centre has reached the lower 95% confidence limit
 Centre has reached the upper 95% confidence limit
 Centre has reached the upper 99.8% confidence limit

* Includes transplants performed between 1 April 2017 - 31 March 2021

** Includes transplants performed between 1 April 2013 - 31 March 2017

Table 6.2 Risk-adjusted one and five year pancreas graft survival for first SPK transplants using pancreases from deceased donors							
Centre	Code	N	Pancreas graft survival				
			One-year*		Five-year**		
			%	(95% CI)	N	%	(95% CI)
Cambridge	A	85	96	(89 - 99)	99	80	(68 - 88)
Cardiff	B	29	89	(68 - 98)	25	82	(57 - 94)
Edinburgh	C	64	90	(79 - 96)	72	76	(61 - 86)
Guy's	D	86	89	(80 - 95)	107	75	(63 - 84)
Manchester	E	85	94	(85 - 98)	112	84	(74 - 91)
Newcastle	F	22	100	N/A	25	87	(62 - 97)
Oxford	G	162	95	(90 - 98)	210	85	(78 - 89)
WLRTC	H	24	100	N/A	28	93	(76 - 99)
UK		557	94	(91 - 95)	678	82	(79 - 85)
<div> <div></div> Centre has reached the lower 99.8% confidence limit <div></div> Centre has reached the lower 95% confidence limit <div></div> Centre has reached the upper 95% confidence limit <div></div> Centre has reached the upper 99.8% confidence limit </div>							
* Includes transplants performed between 1 April 2017 - 31 March 2021 ** Includes transplants performed between 1 April 2013 - 31 March 2017							

6.2 Deceased donor graft and patient survival for first PO transplants

Individual centre unadjusted survival estimates and national rates for one-year and five-year [patient](#) and pancreas [graft](#) survival for deceased donor first pancreas only (PO) transplants are shown in **Tables 6.3** and **6.4**, respectively. Centre specific estimates of these rates must be interpreted with caution due to the small number of transplants upon which they are based.

Table 6.3 Unadjusted one and five year patient survival for first PO transplants using pancreases from deceased donors							
Centre	Code	N	Patient survival				
			One-year*		Five-year**		
			%	(95% CI)	N	%	(95% CI)
Cambridge	A	0	-	-	0	-	-
Cardiff	B	2	-	-	6	-	-
Edinburgh	C	0	-	-	0	-	-
Guy's	D	0	-	-	2	-	-
Manchester	E	2	-	-	3	-	-
Newcastle	F	0	-	-	2	-	-
Oxford	G	8	-	-	26	87	(66 - 96)
WLRTC	H	3	-	-	0	-	-
UK		15	100	-	39	88	(71 - 95)
* Includes transplants performed between 1 April 2017 - 31 March 2021 ** Includes transplants performed between 1 April 2013 - 31 March 2017 - Data not presented where less than 10 transplants included							

Table 6.4 Unadjusted one and five year pancreas graft survival for first PO transplants using pancreases from deceased donors

Centre	Code	N	Pancreas graft survival				
			One-year*		Five-year**		
			%	(95% CI)	N	%	(95% CI)
Cambridge	A	0	-	-	2	-	-
Cardiff	B	3	-	-	10	48	(16 - 74)
Edinburgh	C	0	-	-	1	-	-
Guy's	D	0	-	-	2	-	-
Manchester	E	9	-	-	6	-	-
Newcastle	F	2	-	-	6	-	-
Oxford	G	12	92	(54 - 99)	36	67	(48 - 80)
WLRTC	H	4	-	-	2	-	-
UK		30	87	(68 - 95)	65	66	(52 - 76)

* Includes transplants performed between 1 April 2017 - 31 March 2021
 ** Includes transplants performed between 1 April 2013 - 31 March 2017
 - Data not presented where less than 10 transplants included

Survival from listing

7.1 Patient survival from listing for SPK transplant

Survival from listing was analysed for all adult (≥ 18 years) patients registered for the first time for SPK between 1 January 2010 and 31 December 2021. Patients registered for a pancreas only or islet transplant have been excluded from this analysis. Survival time was defined as the time from joining the [transplant list](#) to death, regardless of the length of time on the [transplant list](#), whether or not the patient was transplanted and any factors associated with such a transplant e.g. donor type. Survival time was censored at either the date of removal from the list, or at the last known follow up date post-transplant when no death date was recorded, or at the time of analysis if the patient was still active on the [transplant list](#).

The [funnel plot](#) shown in **Figure 7.1**, compares centre specific ten-year [risk-adjusted](#) patient survival rates from the point SPK transplant listing and indicates how consistent the rates of the individual transplant centres are with the national rate. All centres survival rates were very similar to the national rate of 77%. Centres can be identified by the information shown in **Table 7.1**, which also shows one and five-year [risk-adjusted](#) survival rates from the point of transplant listing. Note that all rates (at one, five and ten years) were calculated from the same cohort of patients, and the number of patients remaining at risk of death after each time horizon (i.e. not already censored or deceased) is included in **Table 7.1** for reference.




Table 7.1 Risk-adjusted 1, 5 and 10 year patient survival from listing for first deceased donor SPK transplant in patients registered between 1 January 2010 and 31 December 2021

Centre	Code	Number at risk at day 0	One year		Five year		Ten year	
			Survival rate (%) (95% CI)	Number at risk ¹	Survival rate (%) (95% CI)	Number at risk ¹	Survival rate (%) (95% CI)	Number at risk ¹
Cambridge	A	270	97 (95-98)	242	87 (82-90)	132	78 (70-84)	27
Cardiff	B	108	97 (95-98)	97	87 (83-90)	48	78 (72-82)	11
Edinburgh	C	252	97 (95-98)	229	87 (82-91)	114	78 (71-84)	17
Guy's	D	354	97 (96-97)	317	87 (84-89)	157	77 (70-82)	18
Manchester	E	403	97 (95-98)	360	87 (85-89)	155	77 (73-80)	29
Newcastle	F	89	97 (95-98)	77	87 (85-89)	35	77 (73-81)	5
Oxford	G	679	96 (95-98)	610	87 (83-90)	286	77 (72-81)	35
WLRTC	H	102	97 (95-98)	105	86 (85-89)	55	77 (72-81)	13
UK		2257	97 (96-97)	2037	86 (85-88)	982	77 (74-80)	155
<div style="display: flex; align-items: center;"> <div style="width: 20px; height: 10px; background-color: red; margin-right: 5px;"></div> <div>Centre has reached the lower 99.8% confidence limit</div> </div> <div style="display: flex; align-items: center;"> <div style="width: 20px; height: 10px; background-color: #f08080; margin-right: 5px;"></div> <div>Centre has reached the lower 95% confidence limit</div> </div> <div style="display: flex; align-items: center;"> <div style="width: 20px; height: 10px; background-color: #90ee90; margin-right: 5px;"></div> <div>Centre has reached the upper 95% confidence limit</div> </div> <div style="display: flex; align-items: center;"> <div style="width: 20px; height: 10px; background-color: green; margin-right: 5px;"></div> <div>Centre has reached the upper 99.8% confidence limit</div> </div>								

¹ Number of patients with reported follow-up beyond this time point

Form return rates

8.1 Pancreas form return rates, 1 January – 31 December 2021

Form return rates are reported in Table 8.1 for the pancreas transplant record, three month and one year follow up form, along with lifetime follow up (more than two years). These include all pancreas transplants performed between 1 January and 31 December 2021 for the transplant record, and all requests for follow up forms issued in this time period. Centres highlighted are transplant centres. Overall, 86% of transplant record forms issued and 74% of lifetime follow-up forms issued have been returned. Of the transplant centres, WLRTC has the lowest lifetime follow-up form return rates with 59%. Data as on the database at 17th November 2022.

Table 8.1 Form return rates following pancreas transplantation, by centre, 1 January - 31 December 2021								
Centre	Transplant record		3 month follow-up		12 month follow-up		Lifetime follow-up	
	N	% returned	N	% returned	N	% returned	N	% returned
Aberdeen, Aberdeen Royal Infirmary	.	.	1	100	.	.	21	81
Airdrie, Monklands District General Hospital	6	50
Bangor, Ysbyty Gwynedd District General Hospital	8	88
Basildon, Basildon Hospital	5	100
Belfast, Antrim Hospital	3	0
Belfast, Belfast City Hospital	6	100
Belfast, The Ulster Hospital	2	0
Birmingham, Birmingham Heartlands Hospital	13	0
Birmingham, Queen Elizabeth Hospital Birmingham	1	100	50	40
Bradford, St Luke's Hospital	11	82
Brighton, Royal Sussex County Hospital	1	100	24	79
Bristol, Southmead Hospital	36	89
Cambridge, Addenbrookes Hospital	18	94	16	100	19	100	136	100
Canterbury, Kent And Canterbury Hospital	.	.	2	50	3	100	39	100
Cardiff, University Of Wales Hospital	3	67	3	100	6	100	69	100
Carlisle, Cumberland Infirmary	5	100
Carshalton, St Helier Hospital	18	83
Chelmsford, Broomfield Hospital	1	0
Chester, Countess Of Chester Hospital	3	0
County Down, Daisy Hill Hospital	7	86
Coventry, University Hospital (Walsgrave)	2	100	27	100
Derby, Royal Derby Hospital	16	88
Doncaster, Doncaster Royal Infirmary	6	100
Dorchester, Dorset County Hospital	37	49
Douglas, Nobles I-o-M Hospital	5	0
Dudley, Russell's Hall Hospital	4	0
Dulwich, Kings College	2	50
Dumfries, Dumfries And Galloway Royal Infirmary	3	100
Dundee, Ninewells Hospital	20	40
Edinburgh, Royal Infirmary Of Edinburgh	12	100	9	56	9	67	56	63
Exeter, Royal Devon And Exeter Hospital (Wonford)	27	41
Glasgow, Queen Elizabeth University Hospital	30	0
Gloucester, Gloucestershire Royal Hospital	2	100	16	31
Hereford, The County Hospital	5	100
Hull, The Hull Royal Infirmary	19	100
Inverness, Raigmore Hospital	15	73

**Table 8.1 Form return rates following pancreas transplantation, by centre,
1 January - 31 December 2021**

Centre	Transplant record		3 month follow-up		12 month follow-up		Lifetime follow-up	
	N	% returned	N	% returned	N	% returned	N	% returned
Ipswich, Ipswich Hospital	3	100
Kilmarnock, Crosshouse Hospital	8	100
Kirkcaldy, Victoria Hospital	3	100
Larbert, Forth Valley Royal Hospital	5	0
Leeds, St James's University Hospital	21	100
Leicester, Leicester General Hospital	.	.	2	100	2	100	24	88
Lincoln, Lincoln County Hospital	1	100	4	100
Liverpool, Royal Liverpool University Hospital	7	100
Liverpool, University Hospital Aintree	1	100
London, Guys Hospital	15	93	11	45	4	100	145	84
London, Kings College Hospital	.	.	1	100	1	0	2	0
London, St Georges Hospital	6	17
London, The Royal Free Hospital	46	70
London, The Royal London Hospital (Whitechapel)	.	.	3	33	3	100	15	67
Manchester, Manchester Royal Infirmary	25	76	25	76	9	67	104	74
Middlesbrough, The James Cook University Hospital	13	85
Newcastle, Freeman Hospital	6	100	5	80	2	100	60	100
Northampton, Northampton General Hospital	1	100	18	50
Norwich, Norfolk And Norwich University Hospital	1	100	32	100
Nottingham, Nottingham University Hospitals City Campus	1	0	34	0
Omagh, Tyrone County Hospital	.	.	1	0	.	.	2	0
Oxford, Churchill Hospital	37	78	26	92	20	95	156	83
Peterborough, Peterborough City Hospital	5	0
Plymouth, Derriford Hospital	.	.	1	100	1	100	25	76
Portsmouth, Queen Alexandra Hospital	56	100
Portsmouth, St Marys Hospital	.	.	3	100	2	100	6	100
Preston, Royal Preston Hospital	28	71
Reading, Royal Berkshire Hospital	30	63
Rhyl, Royal Alexandra Hospital	2	100
Salford, Salford Royal	17	88
Sheffield, Northern General Hospital	11	73
Shrewsbury, Royal Shrewsbury Hospital	6	83
St Helier, Jersey General Hospital	1	0
Stevenage, Lister Hospital	8	75
Stoke-on-trent, Royal Stoke University Hospital	14	100
Sunderland, Sunderland Royal Hospital	1	0
Swansea, Morriston Hospital	21	71
Truro, Royal Cornwall Hospital (Treliske)	1	0	23	0
West London Renal Transplant Centre	5	100	4	50	3	67	98	59
Westcliff On Sea, Southend Hospital	4	0
Wirral, Arrowe Park Hospital	1	100	4	25
Wolverhampton, New Cross Hospital	1	100	29	93
Wrexham, Maelor General Hospital	13	77
York, York District Hospital	14	100
Overall	121	86	113	78	97	89	1880	74

Islet transplant list

9.1 Patients on the islet transplant list as at 31 March, 2013 – 2022

Figure 9.1 shows the number of patients on the islet [transplant list](#) at 31 March each year. The number of patients active on the islet [transplant list](#) has increased by 37% from 19 on 31 March 2021 to 26 on 31 March 2022. Of the 26, 31% (8) patients were registered for an SIK transplant.




Figure 9.2 shows the number of patients on the active islet [transplant list](#) at 31 March 2022 by centre. Of the 26 patients on the active [transplant list](#) 38% were registered at Manchester, of which six were SIK, 19% at Edinburgh (one SIK) and 12% at Oxford (one SIK). Bristol had no patients registered at 31 March 2022 and are no longer an islet transplant centre.





Figure 9.3 shows the number of patients on the islet [transplant list](#) at 31 March each year between 2013 and 2022 for each transplant centre. There have been very few patients registered at Bristol, King's College or the Royal Free, in the time period.



9.2 Post-registration outcomes, 1 April 2018 – 31 March 2019

An indication of outcomes for patients listed for an islet transplant is summarised in **Figure 9.4**. This shows the proportion of patients transplanted or still waiting one and three years after joining the list. It also shows the proportion removed from the [transplant list](#) (typically because they become too unwell for transplant) and those who died while on the [transplant list](#).

43% of patients were transplanted within one year, while three years after listing 53% of patients had received a transplant and 25% were removed from the list. It is important to note that the three-year period after registration for these patients included two-years impacted by the COVID-19 pandemic.






Figure 9.5 shows the proportion of patients transplanted or still waiting three years after joining the list by centre. Over half the centres have very small numbers of patient registrations in this time period so the figures should be interpreted with caution. Of those centres with patients registered in this time period, the majority transplanted 39% or more of their patients within three years, neither of the two patients registered at the Royal Free were transplanted. 14% and 13% of patients died whilst waiting for an islet transplant at Oxford and Manchester, respectively. Bristol had no patients registered in the time period.




9.3 Demographic characteristics, 1 April 2021 – 31 March 2022

The sex and age group of patients registered on the islet [transplant list](#) during 2021/22 are shown by centre in **Figures 9.6** and **9.7**. Note that all percentages quoted are based only on data where relevant information was available.


Overall, the majority of patients registered on the islet transplant list were female (57%) and the median age was 51 years.





9.4 Patient waiting times for those currently on the list, 31 March 2022

Figure 9.8 shows the length of time patients have been waiting on the islet [transplant list](#) at 31 March 2022 by centre.



9.5 Median active waiting time to transplant, 1 April 2016 - 31 March 2020

The length of time a person waits for routine islet transplant varies across the UK. The [median](#) active waiting time for deceased donor islet transplantation is calculated using the [Kaplan-Meier method](#) and is shown in **Figure 9.9** and **Table 9.1** for patients registered at each individual unit.

The [median](#) active waiting time to transplant for patients registered on the islet [transplant list](#) between 1 April 2016 and 31 March 2020 is 336 days (around 11 months). The median active waiting time is not shown where less than 10 patients are registered.




Table 9.1 Median active waiting time to islet transplant in the UK, for patients registered 1 April 2016 - 31 March 2020

Transplant centre	Number of patients registered	Waiting time (days)	
		Median	95% Confidence interval
Bristol	0	-	-
Edinburgh	41	209	91 - 327
King's	4	-	-
Manchester	46	567	334 - 800
Newcastle	17	467	21 - 913
Oxford	16	317	264 - 370
Royal Free	3	-	-
UK	127	336	217 - 455

- Data not presented when less than 10 patients registered

Response to islet offers

10.1 Offer decline rates, 1 April 2019 – 31 March 2022

Islet offers from [DBD](#) donors whose pancreas was retrieved, offered directly on behalf of a named individual person and resulted in islet transplantation are included in the analysis. Any offers of islets declined for transplantation or [DCD](#) offers were excluded, as were offers made through the fast track scheme or the reallocation of the pancreas.

Individual centre offer decline rates by financial year, 1 April 2019 and 31 March 2022 are shown in **Table 10.1**. King's College had the lowest overall decline rate (0%) whilst Royal Free had the highest decline rate (100%), although both rates are based on less than five offers. Edinburgh had an offer decline rate better than the national average. Bristol had no patients registered and received no offers in this time period.




Table 10.1 DBD donor islet offer decline rates by transplant centre, 1 April 2019 and 31 March 2022

Centre	Code	2019/20 N (%)	2020/21 N (%)	2021/22 N (%)	Overall N (%)
Edinburgh	C	7 (29)	5 (40)	7 (0)	19 (21)
King's	J	1 (0)	0 -	0 -	1 (0)
Manchester	E	8 (38)	4 (75)	7 (71)	19 (58)
Newcastle	F	8 (63)	6 (33)	7 (57)	21 (52)
Oxford	G	4 (50)	2 (0)	7 (86)	13 (62)
Royal Free	K	4 (100)	0 -	0 -	4 (100)
UK		32 (50)	17 (41)	28 (54)	77 (49)

	Centre has reached the upper 99.8% confidence limit
	Centre has reached the upper 95% confidence limit
	Centre has reached the lower 95% confidence limit
	Centre has reached the lower 99.8% confidence limit

Islet transplants

11.1 Islet transplants, 1 April 2012 – 31 March 2022

Figure 11.1 shows the total number of islet transplants performed in the last ten financial years, by type of donor. Since 2012/13, the number of islet transplants has fluctuated around 30 each year apart from the last two years, due to the COVID-19 pandemic.




Figure 11.2 shows the total number of islet transplants performed in 2021/22, by centre and type of donor. The same information is presented in **Figure 11.3** but this shows the proportion of **DBD** and **DCD** transplants performed at each centre. Edinburgh performed the most islet transplants in 2021/22 (13), followed by Manchester (four). Edinburgh, Manchester and Oxford were the only centres to perform **DCD** as well as **DBD** transplants. Royal Free, King's and Bristol did not perform any islet transplants in 2021/22.






Figure 11.4 shows the total number of islet transplants performed in last ten years, by centre and type of donor. Oxford and Manchester have consistently performed a number of islet transplants each year. Edinburgh have consistency performed the most transplants each year. Bristol has performed very few transplants over the ten year period and none in the last five years. Royal Free have performed no islet transplant in the last seven years.



11.2 Demographic characteristics, 1 April 2021 - 31 March 2022

The sex and age group of patients that received an islet transplant in 2021/22 are shown by centre in **Figures 11.5** and **11.6** respectively. Note that all percentages quoted are based only on data where relevant information was available. Overall, 22 patients were transplanted on the islet transplant list, the [median](#) age was 54 years and the majority were female 12 (55%).





Figure 11.7 Islet transplants, 1 April 2021 - 31 March 2022, by centre and sensitisation (cRF%) group





Figure 11.8 Islet transplants, 1 April 2021 - 31 March 2022, by centre and matchability group




Islet outcomes

12.1 Outcome measures for routine islet transplants

Key measures of islet outcome include [graft survival](#), annual rate of severe [hypoglycaemic](#) events, [HbA1c](#) and insulin requirements. This section includes outcomes reported to NHS Blood and Transplant for islet transplants between 1 April 2012 and 31 March 2021.


A one-year [Kaplan-Meier graft survival](#) plot for islet transplants between 1 April 2017 – 31 March 2021 is shown in **Figure 12.1**. Estimated one-year [graft survival](#) following a routine islet transplant is 76% with 95% confidence interval (CI) (55-88%). This includes patients who received only a routine graft and those patients who additionally received a priority graft.

Figure 12.1 One-year graft survival following first routine islet transplant between 1 April 2017 and 31 March 2021



A five-year [Kaplan-Meier graft survival](#) plot for islet transplants between 1 April 2012 – 31 March 2021 is shown in **Figure 12.2**. Estimated five-year [graft survival](#) following a routine islet transplant is 57% with 95% CI (46-66%). This includes patients who received only a routine graft and those who additionally received a priority graft.


Figure 12.2 Five-year graft survival following first routine islet transplant between 1 April 2012 and 31 March 2021



Further, five-year [Kaplan-Meier graft survival](#) plots by type of graft are shown in **Figure 12.3** and **12.4**, for islet transplants between 1 April 2012 – 31 March 2021. **Figure 12.4** only includes routine grafts (routine only or routine followed by a priority) that still were functioning at one year post-transplant. In order to receive a priority (top-up) graft the patient's routine graft must still be functioning and the priority graft should be given within the first 12 months post routine transplant. Therefore, to accurately compare the two groups, i.e. those receiving a routine graft alone and those receiving a routine and subsequent priority graft, the survival estimate is conditional on one-year graft survival in both groups.

Estimated five-year [graft survival](#) (for all islet transplants) is 38% for routine only grafts, 95% CI (22-54%) and for routine followed by priority grafts is 67%, 95% CI (54-78%). This difference was statistically significant, $p=0.0009$.

Figure 12.3 Five-year graft survival following routine islet transplantation, by type of graft, between 1 April 2012 and 31 March 2021



Estimated five-year [graft survival](#) (for islet transplant, where the routine graft was functioning at one year) is 57% for routine only grafts, 95% CI (33-75%) and for routine followed by priority grafts is 73%, 95% CI (58-83%). This difference was not statistically significant, $p=0.2677$.

Figure 12.4 Five-year graft survival following routine islet transplantation, where the routine graft was functioning at one year, between 1 April 2012 and 31 March 2021




Figure 12.5 shows, for routine islet only transplants between 1 April 2017 – 31 March 2021, the [median](#) annual rate of severe [hypoglycaemic](#) events, at registration, prior to transplant (reported as number of events between registration and transplant) and at one-year post-transplant. Of the 19 patients where the number of severe hypoglycaemic events at one-year post-transplant was available, 17 (89%) experienced no severe [hypoglycaemic](#) events, two (11%) experienced one event.

Figure 12.5 Median annual rate of severe hypoglycaemic events for routine islet only transplants, 1 April 2017 to 31 March 2021




Figure 12.6 shows the reduction in [median HbA1c](#) (mmol/mol) for routine islet only transplants between 1 April 2017 – 31 March 2021. [Median HbA1c](#) dropped from 68mmol/mol prior to transplant to 56mmol/mol at one-year post-transplant. Of those 23 patients with HbA1c reported at one-year, 9 (39%) had an [HbA1c](#) less than 53mmol/mol.

Figure 12.6 Median HbA1c (mmol/mol) for routine islet only transplants, 1 April 2017 to 31 March 2021





Figure 12.7 shows the [median](#) insulin dose per kilo recipient body weight at three-months, six-months and one-year post-transplant, for routine islet only transplants between 1 April 2017 – 31 March 2021. Prior to transplant the [median](#) insulin dose is 0.43 units/kg, by six-months the [median](#) dose has dropped to 0.24 units/kg and then increased slightly at one-year post-transplant with a [median](#) dose of 0.26 units/kg. Following islet transplantation, of the 17 patients where information was reported, 9 (53%) achieved insulin independence at some point during their first year post-transplant.

Figure 12.7 Median insulin dose per kilo of recipient weight for routine islet only transplants, 1 April 2017 to 31 March 2021



Form return rates

13.1 Islet form return rates, 1 January – 31 December 2021

Form return rates are reported in **Table 13.1** for the islet transplant record, three month and one year follow-up form, along with lifetime follow-up (more than two years). These include all islet transplants performed between 1 January and 31 December 2021 for the transplant record, and all requests for follow-up forms issued in this time period. Centres highlighted are transplant centres. There were 52% of transplant record and 69% of lifetime follow-up forms returned. 63% of 3-month and 100% of 12-month follow-up forms were returned. Of the transplant centres, London, Kings College Hospital had the lowest lifetime follow-up return rate of 0%.

Table 13.1 Form return rates following islet transplantation, by centre, 1 January - 31 December 2021								
Centre	Transplant record		3 month follow-up		12 month follow-up		Lifetime follow-up	
	N	% returned	N	% returned	N	% returned	N	% returned
Bristol, Southmead Hospital	2	100
Edinburgh, Royal Infirmary of Edinburgh	12	25	8	63	4	100	39	74
Glasgow, Queen Elizabeth University Hospital	1	0
London, Kings College Hospital	6	0
London, The Royal Free Hospital	5	80
Manchester, Manchester Royal Infirmary	4	100	4	50	2	100	13	85
Newcastle, Freeman Hospital	5	100	4	75	2	100	14	71
Oxford, Churchill Hospital	2	0	3	67	.	.	18	67
Overall	23	52	19	63	8	100	98	69

Appendix

A1 Glossary of terms

ABO

The most important human blood group system for transplantation is the ABO system. Every human being is of blood group O, A, B, AB, or one of the minor variants of these four groups. ABO blood groups are present on other tissues and, unless special precautions are taken, a blood group A pancreas transplanted to a blood group O patient will be rapidly rejected.

Active transplant list

When a patient is registered for a transplant, they are registered on what is called the 'active' transplant list. This means that when a donor pancreas becomes available, the patient is included among those who are matched against the donor to determine whether or not the pancreas is suitable for them. It may sometimes be necessary to take a patient off the transplant list, either temporarily or permanently. This may be done, for example, if someone becomes too ill to receive a transplant. The patient is told about the decision to suspend them from the list and is informed whether the suspension is temporary or permanent. If a patient is suspended from the list, they are not included in the matching of any donor pancreases that become available.

Calculated Reaction Frequency (cRF)

For a given patient with detectable [HLA](#) antibodies, the proportion blood group identical donors from a pool of 10,000 against which the recipient has [HLA](#) specific antibodies is calculated. This percentage of donors is termed the 'calculated Reaction Frequency' (cRF), more commonly referred to as the [sensitisation](#) level. Patients with no detectable [HLA](#) antibodies will have 0 [sensitisation](#) (0% cRF).

Case mix

The types of patients treated at a unit for a common condition. This can vary across units depending on the facilities available at the unit as well as the types of people in the catchment area of the unit. The definition of what type of patient a person is depends on the patient characteristics that influence the outcome of the treatment. For example, the case mix for patients registered for a pancreas transplant is defined in terms of various factors such as the blood group, tissue type and age of the patient. These factors have an influence on the chance of a patient receiving a transplant.

Cold ischaemia time (CIT)

The length of time that elapses between a pancreas being removed from the donor to its transplantation into the recipient is called the Cold Ischaemia Time (CIT). Generally, the shorter this time, the more likely the pancreas is to work immediately and the better the long-term outcome. The factors which determine CIT include a) transportation of the pancreas from the retrieval hospital to the hospital where the transplant is performed, b) the need to tissue type the donor and [cross-match](#) the donor and potential recipients, c) the occasional necessity of moving the pancreas to another hospital if a transplant cannot go ahead, d) contacting and preparing the recipient for the transplant, and e) access to the operating theatre.

Confidence interval (CI)

When an estimate of a quantity such as a survival rate is obtained from data, the value of the estimate depends on the set of patients whose data were used. If, by chance, data from a different set of patients had been used, the value of the estimate may have been different. There is therefore some uncertainty linked with any estimate. A confidence

interval is a range of values whose width gives an indication of the uncertainty or precision of an estimate. The number of transplants or patients analysed influences the width of a confidence interval. Smaller data sets tend to lead to wider confidence intervals compared to larger data sets. Estimates from larger data sets are therefore more precise than those from smaller data sets. Confidence intervals are calculated with a stated probability, usually 95%. We then say that there is a 95% chance that the confidence interval includes the true value of the quantity we wish to estimate.

Confidence limit

The upper and lower bounds of a confidence interval.

Cox Proportional Hazards model

A statistical model that relates the instantaneous risk (hazard) of an event occurring at a given time point to the [risk factors](#) that influence the length of time it takes for the event to occur. This model can be used to compare the hazard of an event of interest, such as graft failure or patient death, across different groups of patients.

Cross-match

A cross-match is a test for patient antibodies against donor antigens. A positive cross-match shows that the donor and patient are incompatible. A negative cross-match means there is no reaction between donor and patient and that the transplant may proceed.

Donor after brain death

Donation after brainstem death (DBD) means donation which takes place following the diagnosis of death using neurological criteria

Donor after circulatory death

Donation after circulatory death (DCD) means donation which takes place following the diagnosis of death using circulatory criteria.

Funnel plot

A graphical method that shows how consistent the survival rates of the different transplant units are compared to the national rate. The graph shows for each unit, a survival rate plotted against the number of transplants undertaken, with the national rate and confidence limits around this national rate superimposed. In this report, 95% and 99.8% confidence limits were used. Units that lie within the confidence limits have survival rates that are statistically consistent with the national rate. When a unit is close to or outside the limits, this is an indication that the centre may have a rate that is considerably different from the national rate.

Graft survival rate

The percentage of patients whose grafts are still functioning. This is usually specified for a given time period after transplant. For example, a five-year transplant survival rate is the percentage of transplants still functioning five years after transplant. For the purposes of pancreas transplantation, graft failure is defined as a return to permanent insulin dependence while for islet transplantation graft failure is defined as a C-peptide less than 50 pmol/l.

HbA1c

HbA1c refers to glycated haemoglobin which is measured by clinicians to obtain an overall picture of an individual's average blood sugar levels over a particular period. HbA1c is a valuable indicator of diabetes control.

HLA mismatch

Human Leucocyte Antigens (HLA) are carried on many cells in the body and the immune system can distinguish between those that can be recognised as 'self' (belonging to you or identical to your own) and those that can be recognised as 'nonself'. The normal response of the immune system is to attack foreign/non-self material by producing antibodies against the foreign material. This is one of the mechanisms that provide protection against infection. This is unfortunate from the point of view of transplantation as the immune system will see the graft as just another 'infection' to be destroyed, produce antibodies against the graft and rejection of the grafted organ will take place. To help overcome this response, it is recognised that 'matching' the recipient and donor on the basis of HLA (and blood group) reduces the chances of acute rejection and, with the added use of immunosuppressive drugs, very much improves the chances of graft survival. 'Matching' refers to the similarity of the recipient HLA type and donor HLA type. HLA mismatch refers to the number of mismatches between the donor and the recipient at the A, B and DR (HLA) loci. There can only be a total of two mismatches at each locus. For example, an HLA mismatch value of 000, means that the donor and recipient are identical at all three loci, while an HLA mismatch value of 210 means that the donor and recipient differ completely at the A locus, are partly the same at the B locus and are identical at the DR locus.

Hypoglycaemia

Hypoglycaemia occurs when the level of glucose present in the blood falls below a set point and is the most common complication of insulin therapy. Severe hypoglycaemia is defined as having low blood glucose levels that requires third party assistance to treat and is classed as a diabetic emergency.

Inter-quartile range

The values between which the middle 50% of the data fall. The lower boundary is the lower quartile, the upper boundary the upper quartile.

Kaplan-Meier method

A method that allows patients with incomplete follow-up information to be included in estimating survival rates. For example, in a cohort for estimating one year patient survival rates, a patient was followed up for only nine months before they relocated. If we calculated a crude survival estimate using the number of patients who survived for at least a year, this patient would have to be excluded as it is not known whether or not the patient was still alive at one year after transplant. The Kaplan-Meier method allows information about such patients to be used for the length of time that they are followed-up, when this information would otherwise be discarded. Such instances of incomplete follow-up are not uncommon and the Kaplan-Meier method allows the computation of estimates that are more meaningful in these cases.

Matchability points score

Matchability points score is a score between 1 and 10 reflecting the difficulty with which a well-matched HLA compatible organ can be found and takes into account sensitisation and rareness of HLA type. Scores are updated annually such that 10% of waiting list patients who are easiest to match have score=1 and 10% who are most difficult to match have a score=10.

Median

The midpoint in a series of numbers, so that half the data values are larger than the median, and half are smaller.

Multi-organ transplant

A transplant in which the patient receives more than one organ. For example, a patient may undergo a transplant of a pancreas and liver. Intestinal transplants involving a pancreas are excluded from the whole report.

National Pancreas Offering Scheme

A nationally agreed set of rules for sharing and allocating deceased donor pancreases for pancreas or islet transplant between transplant centres in the UK. The scheme was introduced on 1 December 2010, revised on 11 September 2019 and is administered by NHS Blood and Transplant. Prior to December 2010 deceased donors were allocated on a centre basis.

The Pancreas Offering Scheme, from September 2019, prioritises difficult to match (100% [sensitisation](#) or [matchability points score](#)=10) and long-waiting patients in a top tier. The second tier includes all other blood group eligible patients and assigns an individual point score to all patients based on a number of clinically relevant donor, recipient and transplant related factors. The individual points score assigns more points to patients with lower levels of [HLA mismatch](#), longer waiting times, higher levels of patient [sensitisation](#), short travel times between retrieval to transplant centre, longer duration of dialysis and better donor to recipient age matching. In addition, donors with a lower BMI are clinically desirable for pancreas transplantation whereas donors with a higher BMI are preferable for islet transplantation. As a result, where the donor has a low BMI more points are awarded for patients waiting for a pancreas transplant and where the donor has a high BMI more points are awarded to islet patients. Patients listed nationally for either a pancreas or islet transplant are then ranked by their total points score and the pancreas is offered preferentially to the patient with the highest total number of points, no matter where in the UK they receive their treatment or whether they are waiting for a pancreas or islet transplant.

Patient survival rate

The percentage of patients who are still alive (whether the graft is still functioning or not). This is usually specified for a given time period after transplant. For example, a five-year patient survival rate is the percentage of patients who are still alive five years after their first transplant.

p value

In the context of comparing survival rates across centres, the *p* value is the probability that the differences observed in the rates across centres occurred by chance. As this is a probability, it takes values between 0 and 1. If the *p* value is small, say less than 0.05, this implies that the differences are unlikely to be due to chance and there may be some

identifiable cause for these differences. If the p value is large, say greater than 0.1, then it is quite likely that any differences seen are due to chance.

Risk-adjusted survival rate

Some transplants have a higher chance than others of failing at any given time. The differences in expected survival times arise due to differences in certain factors, the [risk factors](#), among patients. A risk-adjusted survival rate for a centre is the expected survival rate for that centre given the case mix of their patients. Adjusting for case mix in estimating centre-specific survival rates allows valid comparison of these rates across centres and to the national rate.

Risk factors

These are the characteristics of a patient, transplant or donor that influence the length of time that a graft is likely to function or a patient is likely to survive following a transplant. For example, when all else is equal, a transplant from a younger donor is expected to survive longer than that from an older donor and so donor age is a risk factor.

Sensitisation

Potential recipients can develop a number of different [HLA](#) antibodies as a result of exposure to the different [HLA](#) through blood transfusion, previous transplants and pregnancy. Many patients however, have no detectable [HLA](#) antibodies. If a potential recipient has an antibody to an [HLA](#) then they cannot receive a transplant from a donor with that [HLA](#), thus restricting the pool of potential donors. Patients who are clinically incompatible with the donor are excluded from the offering sequence by the [Pancreas Offering Scheme](#).

Unadjusted survival rate

Unadjusted survival rates do not take account of [risk factors](#) and are based only on the number of transplants at a given centre and the number and timing of those that fail within the post-transplant period of interest. In this case, unlike for risk-adjusted rates, all transplants are assumed to be equally likely to fail at any given time. However, some centres may have lower unadjusted survival rates than others simply because they tend to undertake transplants that have increased risks of failure. Comparison of unadjusted survival rates across centres and to the national rate is therefore inappropriate.

A2 Methods

Statistical methodology and risk-adjustment for survival rate estimation

[Unadjusted](#) and [risk-adjusted](#) estimates of [patient](#) and [graft](#) survival for pancreas and simultaneous pancreas and kidney (SPK) transplant are given for each centre.

[Unadjusted](#) rates give an estimate of what the survival rate at a centre is, assuming that all patients at the centre have the same chance of surviving a given length of time after transplant. In reality, patients differ and a [risk-adjusted](#) rate that allows for these differences would give a more meaningful estimate of survival.

Computing unadjusted survival rates

[Unadjusted](#) survival rates were calculated using the [Kaplan-Meier](#) method, which allows patients with incomplete follow-up information to be included in the computation. For example, in a cohort for estimating one-year [patient](#) survival rates, a patient was followed up for only nine months before they relocated. If we calculated a crude survival estimate using the number of patients who survived for at least a year, this patient would have to be excluded, as it is not known whether or not the patient was still alive one year after transplant. The [Kaplan-Meier](#) method allows information about such patients to be used for the length of time that they are followed-up, when this information would otherwise be discarded. Such instances of incomplete follow-up are not uncommon in the analysis of survival data and the [Kaplan-Meier](#) method therefore allows the computation of survival estimates that are more meaningful.

Computing risk-adjusted survival rates

A [risk-adjusted](#) survival rate is an estimate of what the survival rate at a centre would have been if they had had the same mix of patients as that seen nationally. The [risk-adjusted](#) rate therefore presents estimates in which differences in patient mix across centres have been removed as much as possible. For that reason, it is valid to only compare centres using [risk-adjusted](#) rather than [unadjusted](#) rates, as differences among the latter can be attributed to differences in patient mix.

[Risk-adjusted](#) survival estimates were obtained through indirect standardisation. A [Cox](#) Proportional Hazards model was used to determine the probability of survival for each patient based on their individual risk factor values. The sum of these probabilities for all patients at a centre gives the number, E, of patients or grafts expected to survive at least one year or five years after transplant at that centre. The number of patients who actually survive the given time period is given by O. The [risk-adjusted](#) estimate is then calculated by multiplying the ratio O/E by the overall [unadjusted](#) survival rate across all centres. The risk-adjustment models used were based on results from previous studies that looked at factors affecting the survival rates of interest. The factors included in the models are shown in the table below.

First transplants from deceased donors

Simultaneous pancreas and kidney (SPK) survival

1 and 5 year [patient](#) and [graft](#) survival Donor age, donor type, donor BMI and waiting time

Funnel plots for comparing risk-adjusted survival rates

The [funnel plot](#) is a graphical method to show how consistent the survival rates of the different transplant centres are compared to the national rate. The graph shows for each

centre, a survival rate plotted against the number of transplants undertaken, with the national rate and [confidence limits](#) around this national rate superimposed. In this report, 95% and 99.8% [confidence limits](#) were used. Units that lie within the [confidence limits](#) have survival rates that are statistically consistent with the national rate. When a unit is close to or outside the limits, this is an indication that the centre may have a rate that is considerably different from the national rate.

A fundamentally similar method was used to conduct the survival from listing analysis. The [risk factors](#) used are detailed in the table below.

<p>First registrations for simultaneous pancreas and kidney (SPK) transplant</p> <p>1, 5 and 10 year patient Age, sex, grouped registration year, ethnicity, blood group, cRF>85% survival from listing</p>

Systematic Component of Variation

For a given individual who is a resident in a given NHS region registration to the transplant list is modelled as a Bernoulli trial. At the whole area level, this becomes a Binomial process which can be approximated by a Poisson distribution when rare events are modelled. Transplant counts follow similar assumptions.

To allow for the possibility that, even after allowing for area-specific Poisson rates, area differences remain, introduce an additional multiplicative rate factor which varies from area to area. Postulate a non-parametric distribution for the multiplicative factor, with variance σ^2 . If the factor is one for all areas, then area differences are fully explained by the area-specific Poisson rate. If the factor varies with a nonzero variance, σ^2 , then we conclude that there are unexplained area differences.

The systematic component of variation (SCV; McPherson et al., N Engl J Med 1982, 307: 1310-4) is the moment estimator of σ^2 . Under the null hypothesis of homogeneity across areas, the SCV would be zero. The SCV, therefore, allows us to detect variability across areas beyond that expected by chance; the larger the SCV, the greater the evidence of systematic variation across areas.

A one-sided p-value for the hypothesis that the SCV is greater than zero versus the null hypothesis that the SCV is equal to zero was derived using a parametric bootstrap where data were simulated from the Poisson distribution that would be consistent with the null hypothesis (multiplicative rate factor is equal to one in all areas and σ^2 equal to zero). The observed SCV was then compared against this simulated data to calculate the probability that an SCV of at least this size would be observed due to chance if the null hypothesis were true.

10,000 bootstrap samples of size 7 (number of areas) were simulated, where the registration/transplant count in each area was drawn from a Poisson distribution with its expected value being the area-specific expected count (the rate of transplants/registrations in the total population multiplied by the population of the area) . The SCV was then calculated in each of the 10,000 samples and a bootstrap p-value for the SCV in the observed data was estimated as:

$$P_{boot} = \frac{1 + \#\{SCV_{sim} \geq SCV_{obs}\}}{10000 + 1}$$

where $\#\{SCV_{sim} \geq SCV_{obs}\}$ is the number of SCV values in the simulated datasets which are greater than or equal to the SCV in the observed data. This follows the simulation method given in Ibanez et al., BMC Health Services Research, 2009, 9:60. No adjustment was made for area-specific demographic characteristics that may impact the rates of registration to the transplant list and transplantation such as age and sex.

Prepared by:

Statistics and Clinical Research, NHS Blood and Transplant

Mr Joseph Parsons

Mrs Claire Counter