

Detailed Report Actual and Potential Deceased Organ Donation 1 April 2022 - 31 March 2023

South Central Organ Donation Services Team





Table of Contents

1. Donor outcomes

2. Key rates in potential for organ donation

3. Best quality of care in organ donation

- 3.1 Neurological death testing
- 3.2 Referral to Organ Donation Service
- 3.3 Contraindications
- 3.4 SNOD presence
- 3.5 Consent
- 3.6 Solid organ donation

4. Comparative data

- 4.1 Neurological death testing
- 4.2 Referral to Organ Donation Service
- 4.3 SNOD presence
- 4.4 Consent

5. PDA data by hospital and unit

6. Paediatric ICU data

- 6.1 Key numbers for PICUs
- 6.2 Neurological death testing in PICUs
- 6.3 Referral to Organ Donation Service in PICUs
- 6.4 Contraindications in PICUs
- 6.5 SNOD presence for patients in PICUs
- 6.6 Consent for patients in PICUs
- 6.7 Solid organ donation in PICUs

7. Emergency Department data

- 7.1 Referral to Organ Donation Service
- 7.2 Organ donation discussions

8. Additional Data and Figures

- 8.1 Trust/Board Level Benchmarking
- 8.2 Comparative data for DBD and DCD deceased donors

Appendices

- A.1 Definitions
- A.2 Data description
- A.3 Table and figure description

Further Information

- Appendix A.1 contains definitions of terms and abbreviations used throughout this report and summarises the main changes made to the PDA over time.
- The latest Organ Donation and Transplantation Activity Report is available at
- https://www.organdonation.nhs.uk/supporting-my-decision/statistics-about-organ-donation/transplant-activity-report/
- The latest PDA Annual Report and our Power BI reports with up to date metrics are available at https://www.odt.nhs.uk/statistics-and-reports/potential-donor-audit-report/.
- Please refer any queries or requests for further information to your local Specialist Nurse Organ Donation (SNOD)

Source

NHS Blood and Transplant: UK Transplant Registry (UKTR), Potential Donor Audit (PDA) and Referral Record. Issued May 2023 based on data meeting PDA criteria reported at 9 May 2023.



1. Donor Outcomes

A summary of the number of donors, patients transplanted, average number of organs donated per donor and organs donated.


Data in this section is obtained from the UK Transplant Registry


Between 1 April 2022 and 31 March 2023, the South Central Organ Donation Services Team facilitated 135 deceased solid organ donors, resulting in 340 patients receiving a transplant. Additional information is shown in Tables 1.1 and 1.2, along with comparison data for 2021/22. Figure 1.1 shows the number of donors and patients transplanted for the previous ten periods for comparison.

Table 1.1 Donors, patients transplanted and organs per donor,1 April 2022 - 31 March 2023 (1 April 2021 - 31 March 2022 for comparison)									
Donor type	Number of donors	Number of patients transplanted	Average number donated per South Central	0					
DBD DCD DBD and DCD	64 (65) 71 (66) 135 (131)	179 (190) 161 (131) 340 (321)	3.3 (3.7) 3.2 (2.8) 3.2 (3.2)	3.5 (3.4) 2.9 (2.7) 3.2 (3.1)					

In addition to the 135 proceeding donors there were 39 additional consented donors that did not proceed, 6 where DBD organ donation was being facilitated and 33 where DCD organ donation was being facilitated.

Table 1.2 Organs transplanted by type, 1 April 2022 - 31 March 2023 (1 April 2021 - 31 March 2022 for comparison)									
Donor type	Kidney	Numb Pancreas	Number of organs transplanted by Is Liver Heart		oy type Lung	Small bowel			
DBD DCD DBD and DCD	106 (108) 115 (105) 221 (213)	10 (14) 6 (5) 16 (19)	50 (51) 29 (20) 79 (71)	10 (11) 10 (4) 20 (15)	11 (23) 14 (4) 25 (27)	2 (2) 0 (0) 2 (2)			







2. Key Rates in

Potential for Organ Donation


A summary of the key rates on the potential for organ donation


Data in this section is obtained from the National Potential Donor Audit (PDA)

This section presents specific percentage measures of potential donation activity for the South Central Organ Donation Services Team.

Performance in the team has been compared with UK performance in both Figure 2.1 and Table 2.1 using funnel plot boundaries and the Gold, Silver, Bronze, Amber, and Red (GoSBAR) colour scheme. When compared with UK performance, gold represents exceptional, silver represents good, bronze represents average, amber represents below average, and red represents poor performance. See Appendix A.3 for funnel plot ranges used.

It is acknowledged that the PDA does not capture all activity. There may be some patients referred in 2021/22 who are not included in this section onwards because they were either over 80 years of age or did not die in a unit participating in the PDA.












Table 2.1 Key numbers, rates and comparison with national rates,1 April 2022 - 31 March 2023

	s	DBI	כ	S	DCI outh	D	_	eceased outh	donors
	c	entral	UK	Ce	entral	UK	-	entral	UK
Patients meeting organ donation referral criteria ¹		129	1980		452	5307		561	6910
Referred to Organ Donation Service		129	1965		424	4886		533	6482
Referral rate %	G	100%	99%	В	94%	92%	В	95%	94%
Neurological death tested		108	1556						
Testing rate %	В	84%	79%						
Eligible donors ²		102	1439		321	3467		423	4906
Family approached		93	1244		150	1691		243	2935
Family approached and SNOD present		86	1190		131	1526		217	2716
% of approaches where SNOD present	В	92%	96%	В	87%	90%	В	89%	93%
Consent ascertained		67	846		90	959		157	1805
Consent rate %	В	72%	68%	В	60%	57%	В	65%	61%
- Expressed opt in		37	476		57	578		94	1054
- Expressed opt in %		100%	95%		86%	84%		91%	89%
- Deemed Consent		24	284		27	306		51	590
- Deemed Consent %		77%	63%		52%	52%		61%	57%
- Other*		6	86		6	74		12	160
- Other* %		67%	60%		55%	38%		60%	47%
Actual donors (PDA data)		61	783		70	636		131	1419
% of consented donors that became actual donors		91%	93%		78%	66%		83%	79%

¹ DBD - A patient with suspected neurological death

DCD - A patient in whom imminent death is anticipated, ie a patient receiving assisted ventilation, a clinical decision to withdraw treatment has been made and death is anticipated within 4 hours

² DBD - Death confirmed by neurological tests and no absolute contraindications to solid organ donation DCD - Imminent death anticipated and treatment withdrawn with no absolute contraindications to solid organ donation

* Includes patients where nation specific deemed criteria are not met and the patient has not expressed a donation decision in accordance with relevant legislation

Note that a patient that meets both the referral criteria for DBD and DCD organ donation is featured in both the DBD and DCD data but will only be counted once in the deceased donors total

Gold Silver Bronze Amber Red



3. Best quality of care

in organ donation

Key stages in best quality of care in organ donation


Data in this section is obtained from the National Potential Donor Audit (PDA)

This section provides information on the quality of care in the South Central Organ Donation Services Team at the key stages of organ donation. The ambition is that the team misses no opportunity to make a transplant happen and that opportunities are maximised at every stage.

3.1 Neurological death testing

Goal: neurological death tests are performed wherever possible.

Figure 3.1 Number of patients with suspected neurological death, 1 April 2018 - 31 March 2023



Patients not tested Patients tested

Table 3.1 Reasons given for neurological death tests not being performed,1 April 2022 - 31 March 2023

Family declined donation Family pressure not to test Inability to test all reflexes Medical contraindication to donation Other Patient had previously expressed a wish not to donate Patient haemodynamically unstable Pressure of ICU beds SN-OD advised that donor not suitable Treatment withdrawn	2 2 - 1 2 - 7 - 1 -	6 28 48 20 5 43 2 151 1 8 18
Unknown	- - 21	18 3 424
	21	uire



3.2 Referral to Organ Donation Service

Goal: Every patient who meets the referral criteria should be identified and referred to the Organ Donation Service, as per NICE CG135¹ and NHS Blood and Transplant (NHSBT) Best Practice Guidance on timely identification and referral of potential organ donors².

Aim: There should be no purple on the following charts.




Figure 3.2 Number of patients meeting referral criteria, 1 April 2018 - 31 March 2023

Table 3.2 Reasons given why patient not referred to SNOD,1 April 2022 - 31 March 2023

	DBD South		DCD South	
	Central	UK	Central	υĸ
Clinician assessed that patient was unlikely to become asystolic within 4 hours	-	-	-	2
Family declined donation following decision to remove treatment	-	1	3	15
Family declined donation prior to neurological testing	-	1	-	1
Medical contraindications	-	-	1	28
Not identified as potential donor/organ donation not considered	-	6	19	271
Other	-	-	1	27
Patient had previously expressed a wish not to donate	-	-	-	3
Pressure on ICU beds	-	-	-	3
Reluctance to approach family	-	1	-	2
Thought to be medically unsuitable	-	1	4	53
Uncontrolled death pre referral trigger	-	5	-	16
Total	-	15	28	421
If 'other', please contact your local SNOD or CLOD for more inform	mation, if requ	uired.		



3.3 Contraindications

In 2022/23 there were 95 potential donors in the South Central Organ Donation Services team with an ACI reported, 7 DBD and 93 DCD donors. Please note, the number of potential DBD and DCD donors with an ACI reported may not equal the total stated as a patient can meet potential donor criteria for both DBD and DCD donation.



3.4 SNOD presence

Goal: A SNOD should be present during the formal family approach as per NICE CG135¹ and NHS Blood and Transplant (NHSBT) Best Practice Guidance.³

Aim: There should be no purple on the following charts.

In the UK, in 2022/23, when a SNOD was not present for the approach to the family to discuss organ donation, DBD and DCD consent rates were 31% and 19%, respectively, compared with DBD and DCD consent rates of 70% and 61%, respectively, when a SNOD was present.

Within the Trusts in the team, when a SNOD was not present for the approach to the family to discuss organ donation, DBD and DCD consent rates were 29% and 21%, respectively, compared with DBD and DCD consent rates of 76% and 66%, respectively, when a SNOD was present.

Every approach to those close to the patient should be planned with the multidisciplinary team (MDT), should involve the SNOD and should be clearly planned taking into account the known wishes of the patient. The NHS Organ Donor Register (ODR) should be checked in all cases of potential donation and this information must be discussed with the family as it represents the eligible donor's legal consent to donation.




Figure 3.3 Number of families approached by SNOD presence, 1 April 2018 - 31 March 2023

NICE Clinical Guidelines - CG135 [accessed 9 May 2023]

² NHS Blood and Transplant, 2012. Timely Identification and Referral of Potential Organ Donors - A Strategy for Implementation of Best Practice [accessed 9 May 2023]

³ NHS Blood and Transplant, 2013. Approaching the Families of Potential Organ Donors - Best Practice Guidance


[accessed 9 May 2023]



3.5 Consent

In 2022/23 the DBD and DCD consent rates in the team were 72% and 60%, respectively.

Figure 3.4 Number of families approached, 1 April 2018 - 31 March 2023



	DBI	כ	DCI South	0
	South Central	UK	South Central	UK
amily believe patient's treatment may have been limited to	-	1	-	-
acilitate organ donation				
amily concerned donation may delay the funeral	-	2	-	1
amily concerned other people may disapprove/be offended	-	1	-	2 7
amily concerned that organs may not be transplantable	-	1	1	
amily did not believe in donation	-	4	-	12
amily did not want surgery to the body	-	38	1	51
amily divided over the decision	-	21	-	18
amily felt it was against their religious/cultural beliefs	1	40	3	24
amily felt patient had suffered enough	1	22	5	62
amily felt that the body should be buried whole (unrelated to eligious/cultural reasons)	2	20	1	13
amily felt the length of time for the donation process was too	-	17	4	126
amily had difficulty understanding/accepting neurological testing	1	3	-	-
amily wanted to stay with the patient after death	1	2	2	16
amily were not sure whether the patient would have agreed to onation	2	44	11	90
other	2	22	7	73
atient had previously expressed a wish not to donate	15	121	18	17
atient had registered a decision to Opt Out	1	22	3	31
trong refusal - probing not appropriate	-	17	4	31
otal	26	398	60	73



3.6 Solid organ donation

Goal: NHSBT is committed to supporting transplant units to ensure as many organs as possible are safely transplanted.

	DBD)	DCD	
	South		South	
	Central	UK	Central	UK
Clinical - Absolute contraindication to organ donation	-	10	-	8
Clinical - Cardiac arrest during referral	1	2	-	-
Clinical - Considered high risk donor	1	7	-	8
Clinical - DCD clinical exclusion	-	-	-	1
Clinical - No transplantable organ	1	6	1	12
Clinical - Organs deemed medically unsuitable by recipient	2	10	4	51
entres				
Clinical - Organs deemed medically unsuitable on surgical	-	7	-	3
nspection				
Clinical - Other	-	3	-	10
Clinical - PTA post WLST	-	-	14	16
Clinical - Patient actively dying	1	4	-	19
Clinical - Patient asystolic	-	1	-	-
Clinical - Patient's general medical condition	-	2	-	333
Clinical - Positive virology	-	1	-	3
Clinical - Predicted PTA therefore not attended	-	-	-	3
Consent / Auth - Coroner/Procurator fiscal refusal	-	5	-	1(
Consent / Auth - NOK withdraw consent / authorisation	-	5	1	24
ogistical - Other	-	-	-	3
otal	6	63	20	32



4. Comparative Data

A comparison of performance in your team with national data

Data in this section is obtained from the National Potential Donor Audit (PDA)


This section compares the quality of care in the key areas of organ donation in the South Central Organ Donation Services team with the UK rate using funnel plots. The UK rate is shown as a green dashed line and the funnel shape is formed by the 95% and 99.8% confidence limits around the UK rate. The confidence limits reflect the level of precision of the UK rate relative to the number of observations. Performance in the team is indicated by a black cross. The Gold, Silver, Bronze, Amber, and Red colour scheme is used to indicate whether performance in the team, when compared to UK performance, is exceptional (gold), good (silver), average (bronze), below average (amber) or poor (red).

It is important to note that the differences in patient mix have not been accounted for in these plots. Further to these, separate funnel plots for DBD and DCD rates are presented in Section 8.

4.1 Neurological death testing

Goal: neurological death tests are performed wherever possible.

Figure 4.1 Funnel plot of neurological death testing rate, 1 April 2022 - 31 March 2023





When compared with UK performance, the performance within the Trusts in the team was average (bronze) for neurological death testing.



4.2 Referral to Organ Donation Service

Goal: Every patient who meets the referral criteria should be identified and referred to NHSBT's Organ Donation Service, as per NICE CG135¹ and NHS Blood and Transplant (NHSBT) Best Practice Guidance on timely identification and referral of potential organ donors².





When compared with UK performance, the performance within the Trusts in the team was average (bronze) for referral of potential organ donors to NHS Blood and Transplant's Organ Donation Service.



4.3 SNOD presence

Goal: A SNOD should be present during the formal family approach as per NICE CG135¹ and NHS Blood and Transplant (NHSBT) Best Practice Guidance.³





Figure 4.3 Funnel plot of SNOD presence rate, 1 April 2022 - 31 March 2023

When compared with UK performance, the performance within the Trusts in the team was average (bronze) for Specialist Nurse presence when approaching families to discuss organ donation.



4.4 Consent

Figure 4.4 Funnel plot of consent rate, 1 April 2022 - 31 March 2023



When compared with UK performance, the consent rate within the Trusts in the team was average (bronze).



5. PDA data by hospital and unit

A summary of key numbers and rates from the PDA by hospital and unit where patient died

Data in this section is obtained from the National Potential Donor Audit (PDA)

Tables 5.1 and 5.2 show the key numbers and rates for patients who met the DBD and/or DCD referral criteria, respectively. Percentages have been excluded where numbers are less than 10.

Table 5.1 Patients who met the DBD referral criteria - key numbers and rates,1 April 2022 - 31 March 2023

Patients where neurological death was suspected	Patients tested	Neurological death testing rate (%)	Patients referred	DBD referral rate (%)	Patients confirmed dead by neurological testing	Eligible DBD donors	Eligible DBD donors whose family were approached	Approaches where SNOD involved	SNOD presence rate (%)	Consent ascertained	Consent rate (%)	Actual DBD and DCD donors from eligible DBD donors
Buckinghamsh 5	ire Healthca 4	are NHS Trust	5	_	4	4	2	2		2		2
		-		-	4	4	2	2	-	2	-	2
Gloucestershin 4	e Hospitals 4	NHS Foundatio	on Trust 4	-	3	3	3	3	-	2	-	2
		VHS Foundation										
4	4	-	4	-	4	4	4	4	-	3	-	3
Hampshire Ho 4	spitals NHS 3	Foundation Tru -	ust 4	-	3	3	3	3	-	3	-	3
Isle of Wight N 7	HS Trust 7	-	7	-	7	7	7	7	-	6	-	4
Milton Keynes 6	Hospital NF 6	IS Foundation 1	Trust 6	-	6	6	5	5	-	5	-	5
Northampton 0 16	General Hos 13	pital NHS Trust 81	16	100	12	11	10	9	90	4	40	3
Oxford Univers	sity Hospital 26	s NHS Trust 81	32	100	26	26	23	20	87	16	70	16
Portsmouth Ho	ospitals NHS 9	S Trust 75	12	100	9	7	6	5	-	4	-	4
Royal Berkshir 2	e NHS Fou 2	ndation Trust -	2	-	2	2	2	2	-	2	-	1
Salisbury NHS 1	Foundation	Trust	1	-	1	1	1	0	-	0	-	0
University Hos 36	pital Southa 29	mpton NHS Fo 81	undation Tr 36	ust 100	28	28	27	26	96	20	74	18

Table 5.2 Patients who met the DCD referral criteria - key numbers and rates,1 April 2022 - 31 March 2023

Patients for whom imminent death was anticipated	Patients referred	DCD referral rate (%)	Patients for whom treatment was withdrawn	Eligible DCD donors	Eligible DCD donors whose family were approached	Approaches where SNOD involved	SNOD presence rate (%)	Consent ascertained	Consent rate (%)	Actual DCD donors from eligible DCD donors
Buckinghamshire	e Healthcare	NHS Trust								
21	21	100	21	13	4	4	-	2	-	2
Gloucestershire	Gloucestershire Hospitals NHS Foundation Trust									
40	39	98	39	26	6	6		3	-	3



Table 5.2 Patients who met the DCD referral criteria - key numbers and rates,1 April 2022 - 31 March 2023

Patients for			Patients for		Eligible DCD					Actual DCD
whom imminent death was anticipated	Patients referred	DCD referral rate (%)	whom treatment was withdrawn	Eligible DCD donors	donors whose family were approached	Approaches where SNOD involved	SNOD presence rate (%)	Consent ascertained	Consent rate (%)	donors from eligible DCD donors
Great Western	Hospitals NH	S Foundation T	rust							
11	11	100	9	4	3	3	-	3	-	3
Hampshire Hos	spitals NHS F	oundation Trust								
25	25	100	22	14	6	6	-	5	-	3
Isle of Wight NI	HS Trust									
18	16	89	18	10	2	0	-	0	-	0
Milton Keynes	Hospital NHS	Foundation Tru	st							
7	7	-	6	5	1	1	-	1	-	0
Northampton G	eneral Hospit	al NHS Trust								
25	23	92	25	19	6	5	-	4	-	3
Oxford Univers	ity Hospitals I	NHS Trust								
103	91	88	103	80	53	46	87	23	43	19
Portsmouth Ho	spitals NHS 1	Frust								
48	45	94	48	36	13	10	77	9	69	7
Royal Berkshire	e NHS Found	ation Trust								
33	32	97	30	22	16	16	100	12	75	7
Salisbury NHS	Foundation T	rust								
11	11	100	11	10	2	2	-	1	-	1
University Hos	pital Southam	oton NHS Foun	dation Trust							
110	103	94	110	82	38	32	84	27	71	22

Tables 5.1 and 5.2 show the hospital where the patient died. However, it is acknowledged that there are some occasions where a patient is referred in an Emergency Department but moves to a critical care unit. In total for the team in 2022/23 there were 6 such patients. For more information regarding the Emergency Department please see Section 7.



6. Paediatric ICU data

A summary of key numbers for paediatric ICUs

Data in this section is obtained from the National Potential Donor Audit (PDA)

End of life care guidance and practice for paediatric patients does differ and care of the family unit as a whole is a core key principle. Paediatric Intensive Care Units (PICU) systems should never prevent families being offered the opportunity to donate if this is a possibility.

This section provides information on the quality of care for patients that died in PICUs in the South Central Organ Donation Services team at the key stages of organ donation. The ambition is that your PICU misses no opportunity to make a transplant happen and that opportunities are maximised at every stage.

6.1 Key numbers for PICUs

Table 6.1 PICU key numbers comparison with national rates,1 April 2022 - 31 March 2023

	DBI South	D	DCI South	כ	Deceased South	donors
	Central	UK	Central	UK	Central	UK
Patients meeting organ donation referral criteria ¹	6	88	20	219	25	270
Referred to Organ Donation Service	6	86	20	192	25	243
Referral rate %		98%		88%		90%
Neurological death tested	5	50				
Testing rate %		57%				
Eligible donors ²	5	46	19	173	24	219
Family approached	5	35	8	54	13	89
Family approached and SNOD present	5	27	5	44	10	71
% of approaches where SNOD present		77%		81%		80%
Consent ascertained	3	20	4	17	7	37
Consent rate %		57%		31%		42%
Actual donors (PDA data)	3	19	3	13	6	32
% of consented donors that became actual donors		95%		76%		86%

¹ DBD - A patient with suspected neurological death

DCD - A patient in whom imminent death is anticipated, ie a patient receiving assisted ventilation, a clinical decision to withdraw treatment has been made and death is anticipated within 4 hours

² DBD - Death confirmed by neurological tests and no absolute contraindications to solid organ donation DCD - Imminent death anticipated and treatment withdrawn with no absolute contraindications to solid organ donation


Note that a patient that meets both the referral criteria for DBD and DCD organ donation is featured in both the DBD and DCD data but will only be counted once in the deceased donors total



6.2 Neurological death testing in PICUs

Goal: neurological death tests are performed wherever possible.

Figure 6.1 Number of patients with suspected neurological death in PICUs, 1 April 2018 - 31 March 2023



Patients not tested Patients tested

Table 6.2 Reasons given for neurological death tests no PICUs,	t being performe	d in
1 April 2022 - 31 March 2023		
	South Central	UK
Biochemical/endocrine abnormality	-	3
Clinical reason/Clinician's decision Family declined donation	-	3 5 4
Family pressure not to test	1	11
Inability to test all reflexes	-	2
Other	-	4
Patient haemodynamically unstable	-	5
Treatment withdrawn	-	4
Total	1	38
If 'other', please contact your local SNOD or CLOD for more i	nformation, if requ	iired.



6.3 Referral to Organ Donation Service in PICUs

Goal: Every patient who meets the referral criteria should be identified and referred to the Organ Donation Service, as per NICE CG135¹ and NHS Blood and Transplant (NHSBT) Best Practice Guidance on timely identification and referral of potential organ donors².

Aim: There should be no purple on the following charts.




Figure 6.2 Number of patients meeting referral criteria in PICUs, 1 April 2018 - 31 March 2023

Table 6.3 Reasons given why patient not referred to Organ Donation Service in PICUs,1 April 2022 - 31 March 2023

	DBI South	D	DCI South	D
	Central	UK	Central	UK
Family declined donation following decision to remove treatment	-	1	-	1
Medical contraindications	-	-	-	1
Not identified as potential donor/organ donation not considered	-	1	-	17
Other	-	-	-	2
Thought to be medically unsuitable	-	-	-	4
Uncontrolled death pre referral trigger	-	-	-	2
Total	-	2	-	27
If 'other', please contact your local SNOD or CLOD for more inform	mation, if requ	iired.		



6.4 Contraindications in PICUs

In 2022/23 there were no potential donors in the South Central Organ Donation Services team with an ACI reported.




6.5 SNOD presence for patients in PICUs

Goal: A SNOD should be present during the formal family approach as per NICE CG135¹ and NHS Blood and Transplant (NHSBT) Best Practice Guidance.³

Aim: There should be no purple on the following charts.

Figure 6.3 Number of families of PICU patients approached by SNOD presence, 1 April 2018 - 31 March 2023





6.6 Consent for patients in PICUs

In 2022/23 less than 10 families of eligible donors, facilitated in the PICU, were approached to discuss organ donation in the team therefore consent rates are not presented.




Figure 6.4 Number of families of PICU patients approached, 1 April 2018 - 31 March 2023

Table 6.5 Reasons given why consent was not ascertained for PICU patients, 1 April 2022 - 31 March 2023

	DBI	D	DCD South		
	South Central	UK	Central	υĸ	
Family believe patient's treatment may have been limited to	-	1	-	-	
facilitate organ donation					
Family did not believe in donation	-	2	-	2 3	
Family did not want surgery to the body	-	-	-	3	
Family felt it was against their religious/cultural beliefs	-	-	1	4 5	
Family felt patient had suffered enough	-	2	-	5	
Family felt that the body should be buried whole (unrelated to	1	2	-	1	
religious/cultural reasons)					
Family felt the length of time for the donation process was too	-	-	-	4	
long					
Family had difficulty understanding/accepting neurological testing	1	1	-	-	
Family wanted to stay with the patient after death	-	-	1	5	
Family were not sure whether the patient would have agreed to	-	-	-	1	
donation					
Other	-	3 2	-	6	
Patient had previously expressed a wish not to donate	-	2	-	-	
Patient had registered a decision to Opt Out	-	-	1	1	
Strong refusal - probing not appropriate	-	2	1	5	
Total	2	15	4	37	
If 'other', please contact your local SNOD or CLOD for more inform	nation, if requ	iired.			



6.7 Solid organ donation in PICUs

Goal: NHSBT is committed to supporting transplant units to ensure as many organs as possible are safely transplanted.

1 April 2022 - 31 March 2023				
	DBI South	C	DCI South	D
	Central	UK	Central	UK
Clinical - Organs deemed medically unsuitable by recipient centres	-	-	-	1
Clinical - PTA post WLST	-	-	1	2
Consent / Auth - NOK withdraw consent / authorisation	-	1	-	1
Total	-	1	1	4



7. Emergency Department data

A summary of key numbers for Emergency Departments

Data in this section is obtained from the National Potential Donor Audit (PDA)

Most patients who go on to become organ donors start their journey in the emergency department (ED). Deceased donation is important, not just for those people waiting on the transplant list, but also because many people in the UK have expressed a decision in life to become organ donors after their death. The overarching principle of the NHSBT Organ donation and Emergency Department strategy ⁴ is that best quality of care in organ donation should be followed irrespective of the location of the patient within the hospital at the time of death.

7.1 Referral to Organ Donation Service

Goal: No one dies in your ED meeting referral criteria and is not referred to NHSBT's Organ Donation Service. Aim: There should be no blue on the following chart.






Figure 7.1 Number of patients meeting referral criteria that died in the ED, 1 April 2018 - 31 March 2023

7.2 Organ donation discussions

Goal: No family is approached in ED regarding organ donation without a SNOD present. Aim: There should be no red on the following chart.





SNOD not present SNOD present

^a NHS Blood and Transplant, 2016. Organ Donation and the Emergency Department [accessed 9 May 2023]



8. Additional data and figures

Key numbers and rates on the potential for organ donation

Data in this section is obtained from the National Potential Donor Audit (PDA)

8.1 Trust/Board Level Benchmarking

Trust/Board levels were reallocated in July 2018 using the average number of donors in 2016/17 and 2017/18, Table 8.1 shows the criteria used and how many Trusts/Boards belong to each level.

Table 8.1 Trust/Board level categories					
		Number of Trusts Boards in each level			
Level 1	12 or more (\geq 12) proceeding donors per year	35			
Level 2	6 or more but less than 12 (\geq 6 to <12) proceeding donors per year	45			
Level 3	More than 3 but less than 6 (>3 to <6) proceeding donors per year	47			
Level 4	3 or less (\leq 3) proceeding donors per year	41			

Tables 8.2 and 8.3 show the national DBD and DCD key numbers and rates for the UK by Trust/Board level, to aid in comparison with equivalent Trusts/Boards. Note that percentages have been excluded where numbers are less than 10.

Tabl	Table 8.2 National DBD key numbers and rate by Trust/Board level, 1 April 2022 - 31 March 2023												
	Patients where neurological death was suspected	Patients tested	Neurological death testing rate (%)	Patients referred	DBD referral rate (%)	Patients confirmed dead by neurological testing	Eligible DBD donors	Eligible DBD donors whose family were approached	Approaches where SNOD present	SNOD presence rate (%)	Consent ascertained	Consent rate (%)	Actual DBD and DCD donors from eligible DBD donors
Level 1	1133	896	79	1124	99	879	831	714	677	95	474	66	438
Level 2	441	340	77	439	100	331	307	267	259	97	182	68	171
Level 3	287	229	80	283	99	224	216	188	184	98	135	72	124
Level 4	119	91	76	119	100	90	85	75	70	93	55	73	50

Table 8.3 National DCD key numbers and rate by Trust/Board level,1 April 2022 - 31 March 2023

Level 1	Patients for whom imminent death was anticipated 2564	Patients referred 2370	DCD referral rate (%) 92	Patients for whom treatment was withdrawn 2464	Eligible DCD donors 1772	Eligible DCD donors whose family were approached 941	Approaches where SNOD present 856	SNOD presence rate (%) 91	Consent ascertained 537	Consent rate (%) 57	Actual DCD donors from eligible DCD donors 369
Level 2	1346	1239	92	1313	841	373	333	89	209	56	132
Level 3	979	910	93	944	571	269	241	90	155	58	97
Level 4	418	367	88	408	283	108	96	89	58	54	38



8.2 Comparative data for DBD and DCD deceased donors

Funnel plots are presented in Section 4 showing performance in the team against the UK rate for deceased organ donation. The following funnel plots present data for DBD and DCD donors separately.




Figure 8.1 Funnel plots of referral rates, 1 April 2022 - 31 March 2023

When compared with UK performance, the performance within the Trusts in the team was exceptional (gold) for referral of potential DBD organ donors and average (bronze) for referral of potential DCD organ donors to NHS Blood and Transplant's Organ Donation Service.





Figure 8.2 Funnel plots of SNOD presence rates, 1 April 2022 - 31 March 2023

When compared with UK performance, the performance within the Trusts in the team was average (bronze) and average (bronze) for Specialist Nurse presence in approaches to families of eligible DBD and DCD donors, respectively.



Figure 8.3 Funnel plots of consent rates, 1 April 2022 - 31 March 2023



When compared with UK performance, the consent rate within the Trusts in the team was average (bronze) and average (bronze) for DBD and DCD donors, respectively.



Appendices

Appendix A.1 Definitions

Potential Donor Audit Definitions

Potential Donor Audit inclusion criteria	1 October 2009 – 31 March 2010 All deaths in critical care in patients aged 75 and under, excluding cardiothoracic intensive care units 1 April 2010 – 31 March 2013 All deaths in critical and emergency care in patients aged 75 and under, excluding cardiothoracic intensive care units 1 April 2013 onwards All deaths in critical and emergency care in patients aged 80 and under (prior to 81st birthday)
--	---

Donors after brain death (DBD) definitions

A patient who meets all of the following criteria: invasive ventilation, Glasgow Coma Scale 3 not explained by sedation, no respiratory effort, fixed pupils, no cough or gag reflex. Excluding those not tested due to reasons 'cardiac arrest despite resuscitation', 'brainstem reflexes returned', 'neonates – below 37 weeks corrected gestational age'. Previously referred to as brain death
Neurological death tests performed to confirm and diagnose death
A patient with suspected neurological death
A member of Organ Donation Services Team including: Team Manager, Specialist Nurse Organ Donation, Specialist Requester, Donor Family Care Nurse
A patient with suspected neurological death referred to a SNOD. A referral is the provision of information to determine organ donation suitability. NICE CG135 (England) : Triggers for clinicians to refer a potential donor are a plan to withdraw life sustaining treatment or a plan to perform neurological death tests
A patient with suspected neurological death
Absolute medical contraindications identified in assessment which clinically preclude organ donation as per NHSBT criteria (POL188) Absolute medical contraindications to donation are listed here: https://nhsbtdbe.blob.core.windows.net/umbraco-assets-corp/17160/ clinical-contraindications-to-approaching-families-for-possible-organ-donati on-pol188.pdf
A patient confirmed dead by neurological death tests, with no absolute medical contraindications to solid organ donation
Family of eligible DBD asked to make or support patient's organ donation decision - This includes clarifying an opt out decision
Family supported opt in decision, deemed consent/authorisation, or where applicable the family or nominated/appointed representative gave consent/authorisation for organ donation
Patients who became actual DBD donors following confirmation of neurological death, as reported through the PDA (80 years and below). At least one organ donated for the purpose of transplantation (includes organs retrieved for transplant however used for research)
Patients who became actual DCD donors following confirmation of neurological death, as reported through the PDA (80 years and below). At least one organ donated for the purpose of transplantation (includes organs retrieved for transplant however used for research)



Neurological death testing rate	Percentage of patients for whom neurological death was suspected who were tested
Referral rate	Percentage of patients for whom neurological death was suspected who were referred to the SNOD
Donation decision conversation rate	Percentage of eligible DBD families or nominated/appointed representatives who were asked to make or support an organ donation decision - This includes clarifying an opt out decision
Consent/Authorisation rate	Percentage of donation decision conversations where consent/authorisation was ascertained
SNOD presence rate	Percentage of donation decision conversations where a SNOD was present (includes telephone and video call conversations)
Consent/Authorisation rate where SNOD was present	Percentage of donation decision conversations where a SNOD was present and consent/authorisation for organ donation was ascertained (as above)

Donors after circulatory death (DCD) definitions

Γ

ventilatio made an	, not confirmed dead using neurological criteria, receiving invasive n, in whom a clinical decision to withdraw treatment has been d a controlled death is anticipated within a time frame to allow to occur (as determined at time of assessment)
	for whom imminent (controlled) death is anticipated following al of life sustaining treatment (as defined above)
	er of Organ Donation Services Team including: Team Manager, st Nurse Organ Donation, Specialist Requester, Donor Family Care
SNOD. A donation a potentia	for whom imminent death is anticipated who was referred to a A referral is the provision of information to determine organ suitability NICE CG135 (England) : Triggers for clinicians to refer al donor are a plan to withdraw life sustaining treatment or a plan m neurological death tests
	who had treatment withdrawn and imminent death was ed within a time frame to allow donation to occur.
preclude medical o https://nh	medical contraindications identified in assessment which clinically organ donation as per NHSBT criteria (POL188). Absolute contraindications to donation are listed here: hsbtdbe.blob.core.windows.net/umbraco-assets-corp/17160/ ontraindications-to-approaching-families-for-possible-organ-donati 8.pdf
	who had treatment withdrawn and imminent (controlled) death cipated, with no absolute medical contraindications to solid organ
there are	ecific criteria determine a patient's suitability to donation when no absolute medical contraindications (see absolute dications documentation above)
	by which an organ may be screened with a local and national nt centre to determine suitability of organs for transplantation
for donat	le DCD donor to be assessed considered to be medically suitable ion (i.e. no DCD exclusions and not deemed unsuitable by the g process)
	f medically suitable eligible DCD donor who were asked to make rt patient's organ donation decision - This includes clarifying an ecision.
applicabl	upported opt in decision, deemed consent/authorisation, or where le the family or nominated/appointed representative gave authorisation for organ donation



Actual DCD	DCD patients who became actual DCD as reported through the PDA (80 years and below). At least one organ donated for the purpose of transplantation (includes organs retrieved for transplant however used for research)
Referral rate	Percentage of patients for whom imminent (controlled) death was anticipated who were referred to the SNOD
Donation decision conversation rate	Percentage of medically suitable eligible DCD families or nominated/appointed representatives who were asked to make or support an organ donation decision - This includes clarifying an opt out decision
Consent/Authorisation rate	Percentage of donation decision conversations where consent/authorisation was ascertained.
SNOD presence rate	Percentage of donation decision conversations where a SNOD was present (includes telephone and video call conversations).
Consent/Authorisation rate where SNOD was present	Percentage of donation decision conversations where a SNOD was present and consent/authorisation for organ donation was ascertained (as above).

Deemed Consent/Authorisation

Deemed consent applies if a person who died in Wales, Jersey or England has not expressed an organ donation decision either to opt in or opt out or nominate/appoint a representative, is aged 18 or over, has lived in the country in which they died for longer than 12 months and is ordinarily resident there, and had the capacity to understand the notion of deemed consent for a significant period before their death.

Deemed authorisation applies if a person who died in Scotland has not expressed, in writing, an organ donation decision either to opt in or opt out, is aged 16 or over, has lived in Scotland for longer than 12 months and is ordinarily resident there, and had the capacity to understand the notion of deemed authorisation for a significant period before their death. Note that, in Scotland, a patient who has verbally expressed an opt in decision is included as a deemed authorisation, whereas a patient who has verbally expressed an opt out decision is not included.

Consent/Authorisation groups

Expressed opt in	Patient had expressed an opt in decision. Opt in decisions can be expressed in writing or via the ODR in all nations and verbal opt in decisions are also included in Wales, England and Jersey. Verbally expressed opt in decisions are not included in Scotland
Deemed consent/authorisation	Patient meets deemed criteria specific to each nation as described above. In Scotland, this includes patients who have verbally expressed a decision to opt in
Expressed opt out	Patient had expressed an opt out decision. Opt out decisions can be expressed verbally, in writing or via the ODR in all nations
Other	Patient has expressed no decision or deemed criteria are not met. Paediatric patients are included in this group

UK Transplant Registry (UKTR) definitions

Donor type	Type of donor: Donation after brain death (DBD) or donation after circulatory death (DCD)	
Number of actual donors	Total number of donors reported to the UKTR	
Number of patients transplanted	Total number of patients transplanted from these donors	
Organs per donor	Number of organs donated divided by the number of donors.	
Number of organs transplanted	Total number of organs transplanted by organ type	



Appendix A.2 Data Description

This report provides a summary of data relating to potential and actual organ donors as recorded by NHS Blood and Transplant via the Potential Donor Audit (PDA), the accompanying Referral Record, and the UK Transplant Registry (UKTR) for the specified Trust, Board, Organ Donation Services Team, or nation.

This report is provided for information and to facilitate case based discussion about organ donation by the Organ Donation Committees and Trusts/Boards.

As part of the PDA, patients over 80 years of age and those who did not die on a critical care unit or emergency department are not audited nationally and are therefore excluded from the majority of this report. Data from neonatal intensive care units (ICU) have also been excluded from this report. In addition, some information may be outstanding due to late reporting and difficulties obtaining patient notes. Donations not captured by the PDA will still be included in the data supplied from the accompanying Referral Record or from the UKTR, as appropriate.



٦

Appendix A.3 Table and Figure Description

For the purposes of this report please note that Trust/Board is equivalent to team.

1 Donor outcomes	
Table 1.1	The number of actual donors, the resulting number of patients transplanted and the average number of organs donated per donor have been obtained from the UK Transplant Registry (UKTR) for your Trust/Board. Results have been displayed separately for donors after brain death (DBD) and donors after circulatory death (DCD).
Table 1.2	The number of organs transplanted by type from donors at your Trust/Board has been obtained from the UKTR. Further information can be obtained from your local Specialist Nurse – Organ Donation (SNOD), specifically regarding organs that were not transplanted. Results have been displayed separately for DBD and DCD.
Figure 1.1	The number of actual donors and the resulting number of patients transplanted obtained from the UKTR for your Trust/Board for the past 10 equivalent time periods are presented on a line chart.

Figure 2.1	Key percentage measures of DBD and DCD potential donation activity for your Trust/Board are presented in a bar chart, using data from the Potential Donor Audit (PDA). The comparative UK rate, for the same time period, is illustrated by the pink line. The key rates labels are coloured using the gold, silver, bronze, amber, and red (GoSBAR) colour scheme to show the performance of your Trust/Board, relative to the UK rate, as reflected in the funnel plots (see description for Figure 4.1 below.
Figure 2.2	Trends in the key percentage measures of DBD and DCD potential donation activity for your Trust/Board are presented for the past five equivalent time periods, using data from the PDA.
Table 2.1	A summary of DBD, DCD and deceased donor data and key numbers have been obtained from the PDA. A UK comparison is also provided. Note that caution should be applied when interpreting percentages based on small numbers. Appendix A.1 gives a fuller explanation of terms used. The key rates are highlighted using the gold, silver, bronze, amber, and red (GoSBAR) colour scheme to show the performance of your Trust/Board, relative to the UK rate, as reflected in the funnel plots (see description for Figure 4.1 below).

3 Best quality of care in organ d	onation
Figure 3.1	A stacked bar chart displays the number of patients with suspected neurological death who were tested and the number who were not tested in your Trust/Board for the past five equivalent time periods.
Table 3.1	The reasons given for neurological death tests not being performed in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.
Figure 3.2	Stacked bar charts display the number of DBD and DCD patients meeting referral criteria who were referred to the Organ Donation Service and the number who were not referred in your Trust/Board for the past five equivalent time periods.
Table 3.2	The reasons given for not referring patients to the Organ Donation Service in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.
Table 3.3	The primary absolute medical contraindications to solid organ donation for DBD and DCD patients have been obtained from the PDA, if applicable. A UK comparison is also provided.
Figure 3.3	Stacked bar charts display the number of families of DBD and DCD patients approached where a SNOD was present and the number approached where a SNOD was not present in your Trust/Board for the past five equivalent time periods.



Figure 3.4	Stacked bar charts display the number of families of DBD and DCD patients approached where consent/authorisation for organ donation was ascertained and the number approached where consent/authorisation was not ascertained in your Trust/Board for the past five equivalent time periods.
Table 3.4	The reasons why consent/authorisation was not ascertained for solid organ donation in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.
Table 3.5	The reasons why solid organ donation did not occur in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.

4 Comparative data	
Figure 4.1	A funnel plot of the neurological death testing rate is displayed using data obtained from the PDA. Each Trust/Board, of the same level, is represented on the plot as a blue dot, although one dot may represent more than one Trust/Board. The UK rate is shown on the plot as a green horizontal dashed line, together with 95% and 99.8% confidence limits for this rate. These limits form a 'funnel', which is shaded using the gold, silver, bronze, amber, and red (GoSBAR) colour scheme. Graphs obtained in this way are known as funnel plots. If a Trust/Board lies within the 95% limits, shaded bronze, then that Trust/Board has a rate that is statistically consistent with the UK rate (average performance). If a Trust/Board lies outside the 95% confidence limits, shaded silver (good performance) or amber (below average performance), this serves as an alert that the Trust/Board may have a rate that is significantly different from the UK rate. When a Trust/Board lies above the upper 99.8% limit, shaded gold, this indicates a rate that is significantly higher than the UK rate (exceptional performance), while a Trust/Board that lies below the lower limit, shaded red, has a rate that is significantly lower than the UK rate (poor performance). It is important to note that differences in patient mix have not been accounted for in these plots. Your Trust/Board is shown on the plot as the large black cross. If there is no large black cross on the plot, your Trust/Board did not report any patients of the type presented. The funnel plots can also be used to identify the maximum rates currently being achieved by Trusts/Boards with similar donor potential.
Figure 4.2	A funnel plot of the deceased donor referral rate is displayed using data obtained from the PDA. See description for Figure 4.1 above.
Figure 4.3	A funnel plot of the deceased donor SNOD presence rate is displayed using data obtained from the PDA. See description for Figure 4.1 above.
Figure 4.4	A funnel plot of the deceased donor consent/authorisation rate is displayed using data obtained from the PDA. See description for Figure 4.1 above.

5 PDA data by hospital and unit	
Table 5.1	DBD key numbers and rates by unit where the patient died have been obtained from the PDA. Percentages have been excluded where numbers are less than 10.
Table 5.2	DCD key numbers and rates by unit where the patient died have been obtained from the PDA. Percentages have been excluded where numbers are less than 10.



6 Paediatric ICU data		
Table 6.1	A summary of DBD, DCD and deceased donor data and key numbers for paediatric ICUs have been obtained from the PDA. A UK comparison is also provided. Note that caution should be applied when interpreting percentages based on small numbers. Appendix A.1 gives a fuller explanation of terms used.	
Figure 6.1	A stacked bar chart displays the number of paediatric ICU patients with suspected neurological death who were tested and the number who were not tested in your Trust/Board for the past five equivalent time periods.	
Table 6.2	The reasons given for neurological death tests not being performed for paediatric ICU patients in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.	
Figure 6.2	Stacked bar charts display the number of DBD and DCD paediatric ICU patients meeting referral criteria who were referred to the Organ Donation Service and the number who were not referred in your Trust/Board for the past five equivalent time periods.	
Table 6.3	The reasons given for not referring paediatric ICU patients to the Organ Donation Service in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.	
Table 6.4	The primary absolute medical contraindications to solid organ donation for DBD and DCD paediatric ICU patients have been obtained from the PDA, if applicable. A UK comparison is also provided.	
Figure 6.3	Stacked bar charts display the number of families of DBD and DCD paediatric ICU patients approached where a SNOD was present and the number approached where a SNOD was not present in your Trust/Board for the past five equivalent time periods.	
Figure 6.4	Stacked bar charts display the number of families of DBD and DCD paediatric ICU patients approached where consent/authorisation for organ donation was ascertained and the number approached where consent/authorisation was not ascertained in your Trust/Board for the past five equivalent time periods.	
Table 6.5	The reasons why consent/authorisation was not ascertained for solid organ donation in paediatric ICU patients in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.	
Table 6.6	The reasons why solid organ donation did not occur in paediatric ICU patients in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.	

Γ

7 Emergency department data	
Figure 7.1	Stacked bar charts display the number of patients that died in the emergency department (ED) who met the referral criteria and were referred to the Organ Donation Service and the number who were not referred in your Trust/Board for the past five equivalent time periods.
Figure 7.2	Stacked bar charts display the number of families of patients in ED approached where a SNOD was present and the number approached where a SNOD was not present in your Trust/Board for the past five equivalent time periods.

8 Additional data and figures	
Table 8.1	A summary of deceased donor, transplant, transplant list and ODR opt-in registration data for your region have been obtained from the UKTR. Your region has been defined as per former Strategic Health Authority. A UK comparison is also provided.
Table 8.2	Trust/board level categories and the relevant expected number of proceeding donors per year are provided for information.
Table 8.3	National DBD key numbers and rates for level 1, 2, 3 and 4 Trusts/Boards are displayed alongside your local data to aid comparison with equivalent Trusts/Boards. Percentages have been excluded where numbers are less than 10.



Table 8.4	National DCD key numbers and rates for level 1, 2, 3 and 4 Trusts/Boards are displayed alongside your local data to aid comparison with equivalent Trusts/Boards. Percentages have been excluded where numbers are less than 10.
Figure 8.1	A funnel plot of the DBD and DCD referral rates are displayed using data obtained from the PDA. See description for Figure 4.1 above.
Figure 8.2	A funnel plot of the DBD and DCD SNOD presence rates are displayed using data obtained from the PDA. See description for Figure 4.1 above.
Figure 8.3	A funnel plot of the DBD and DCD consent/authorisation rates are displayed using data obtained from the PDA. See description for Figure 4.1 above.