



UK Health  
Security  
Agency



**Blood and Transplant**

# Data sources and methods 2022

NHS Blood and Transplant and UK Health Security Agency

# Contents

Summary .....	3
Blood donor surveillance.....	4
Transfusion Transmitted Infections.....	12
Bacterial screening of platelets.....	14
Tissue and cord blood donor surveillance.....	16
Solid organ donor surveillance.....	18
Risk estimates for blood donors.....	19
Emerging infections.....	21

The aim of the NHS Blood and Transplant (NHSBT) and UK Health Security Agency (UKHSA) Epidemiology Unit is to monitor infections in blood, tissue and cell, and organ donors and recipients and to use this information to inform blood safety.

This document provides more detailed information on the data sources and methods used in each of the surveillance schemes run by the NHSBT and UKHSA Epidemiology Unit. It can be read in conjunction with the annual review.

The surveillance schemes can be modified to incorporate any changes to testing, such as the introduction of a new test, or when new data need to be gathered. These schemes are continually monitored to ensure high quality data is consistently collected.

Data is regularly published reports for different groups. The main reports include:

- monthly blood donation testing reports for UKBS
- monthly confirmed positive charts for NHSBT
- monthly emerging infections report (EIR) for SACTTI/JPAC
- monthly and quarterly bacterial screening of platelets report
- quarterly positive donors report for Transfusion Microbiology Clinical Group
- transfusion transmitted infections chapter in the Serious Hazards of Transfusion (SHOT) annual report
- blood donor sections for the UKHSA annual reports on HBV, HCV, HIV and syphilis
- annual residual risk estimates for Joint United Kingdom Blood Transfusion Services Professional Advisory Committee (JPAC)
- annual NHSBT/UKHSA review (all surveillance schemes)

The surveillance schemes are described on the UKHSA website - [Blood, tissue and organ donors: surveillance schemes - GOV.UK \(www.gov.uk\)](#)

Donated blood, tissues and cells are collected by the UK Blood Services from volunteer adults who do not acknowledge any medical condition, travel history or behaviour that is known to be associated with an increased risk of blood borne infections. The donor selection guidelines are available on the [Transfusion Guidelines website](#)

Document last updated 25 October 2023

UKHSA is an executive agency of the Department of Health and Social Care. It is responsible for planning, preventing and responding to external health threats, and providing intellectual, scientific and operational leadership at national and local level, as well as on the global stage.

[Return to Table of Contents](#)

# Blood donor surveillance

Blood donor data is collected by the NHSBT and UKHSA Epidemiology Unit through two parallel schemes: blood donation testing and confirmed positive blood donors. Both surveillance schemes started in October 1995. The information is used to monitor donation testing reactive rates and confirmed markers of infection detected in blood donors. Additional denominator data providing more detailed information on the profile of all blood donors making a donation by year is also gathered.

## Blood donation testing

### Background

Blood donations have been tested for infections since the 1940s when testing for markers of treponemes indicating syphilis (or other disease such as yaws or pinta that are also caused by *Treponema*) first began. Since then, screening for other blood-borne infections has been introduced as follows:

#### Hepatitis B virus (HBV)

Screening donations for hepatitis B surface antigen (HBsAg) began in 1972. HBV DNA testing began on 1 April 2009 in Filton as a by-product of the introduction of triplex NAT testing. HBV DNA testing was subsequently introduced at Manchester on 10 August 2009 and Colindale on 3 December 2009, when all donations in England were HBV DNA tested. HBV DNA testing began on 27 April 2009 in the Republic of Ireland and 1 June 2009 in Wales. Scotland and Northern Ireland began HBV DNA testing on 22 March 2010. Hepatitis B core antibody screening began in 2022 across the UK to supplement current screening procedures.

#### Human Immunodeficiency Virus (HIV)

Donation screening for antibodies to HIV (anti-HIV) began in 1985. Some England centres started using combined antigen-antibody screening for HIV (HIV Ag/Ab) in 2001 with all using HIV Ag/Ab by the end of 2005. Northern Ireland and Republic of Ireland Blood Transfusion Services introduced combined Ab/Ag testing in 2011 and 2016 respectively. Although not mandatory, HIV RNA testing has been introduced as a by-product of duplex and subsequently triplex assays. HIV RNA testing was introduced in Scotland and Northern Ireland in 2002 and some parts of England and Wales from November 2003 but did not become universal until 2007.

#### Hepatitis C virus (HCV)

Donation screening for antibodies to hepatitis C virus (anti-HCV) began in 1991. HCV RNA testing was introduced on a pilot basis in April 1999 and became a mandatory test carried out on all blood donations from 2002.

### Human T-cell lymphotropic virus I and II (anti-HTLV)

Testing for antibodies to human T-cell lymphotropic virus I and II (anti-HTLV) was piloted in Scotland in 2000 and started UK-wide in 2002. Testing in the UK was conducted in pools until 2013 when singleton testing was implemented in England from 27 February 2013. Scotland switched to singleton testing in 2015, Northern Ireland in 2016 and Wales switched in 2018. Republic of Ireland used singleton testing from 2002. In January 2017, NHSBT tested donations from new donors and donations used for non leucodepleted blood components. In 2022, SNBTS also changed to testing new donors and donations used for non-leucodepleted products for HTLV. In addition any repeat donors who advise of a sexual contact with a HTLV positive person since their initial HTLV test. NIBTS and WBS test every donation for HTLV.

### Hepatitis E virus (HEV)

Donation screening for HEV RNA started in 2016 on a proportion of donations in order to supply HEV-screened components for specific patient groups and extended in 2017 to all blood donations: starting with SNBTS on March 1st 2017, NHSBT in April 2017, and NIBTS on 30th May 2017 in pools of 24. WBS began individual donation NAT for HEV on apheresis platelets in November 2022.

Table A: Tests performed on blood donations within the UK blood services in 2022 by country

2022 blood services tested for	England	Northern Ireland	Scotland	Wales	Republic of Ireland	Guernsey	Isle of Man
HBsAg	yes	yes	yes	yes	yes	yes	yes
HBV DNA	yes	yes	yes	yes	yes	no	no
Anti-HBc	yes	yes	yes	yes	yes	no	no
Anti-HCV	yes	yes	yes	yes	yes	yes	yes
HCV RNA	yes	yes	yes	yes	yes	no	no
HEV RNA	pools 24	pools16	pools16	pools 16 or single for apheresis	singles	no	by NHSBT
Combined HIV Ab/Ag	yes	no	yes	yes	yes	yes	yes
HIV Ab only	no	yes	no	no	no	no	no
HIV RNA	yes	yes	yes	yes	yes	no	no
Anti-HTLV	new and non-leuco depleted	yes	yes	yes	yes	yes	yes
Anti-treponeme	yes	yes	yes	yes	yes	yes	yes

In 2022, all donations were screened for HBV using HBsAg and HBV DNA screening performed in pools of 24 for England, Northern Ireland, Scotland and Wales, and in singletons for Republic of Ireland. Anti-HBc screening was rolled out in 2022 for England (31 May), Northern Ireland (29 May), Scotland (5 April) and Wales (27 May) with the aim of screening all donors at least once. : Scotland had capacity from the start to screen all current donors once then donations from new and returning donors; Northern Ireland and Wales screen all donations.; England's capacity to screen all donations increased with time; donations with repeat reactive anti-HBc and anti-HBs over 100 IU/L on screening were discarded without additional confirmatory testing; all screen anti-HBc reactivities had confirmatory tests from March 2023; and functionality to allow screening of all donors once only rather than testing at every donation started from May 2023. Republic of Ireland had already been screening all donations for anti-HBc prior to 2022.

In 2022, all donations were screened for HCV using anti-HCV and HCV RNA screening performed in pools of 24 for England, Northern Ireland, Scotland and Wales, and in singletons for Republic of Ireland

In 2022, all donations were screened for HIV using combined HIV Ab/Ag assay, except Northern Ireland where HIV Ab assay is used and HIV RNA screening performed in pools of 24 for England, Northern Ireland, Scotland and Wales, and singletons for Republic of Ireland

In 2022, donations were screened for HTLV antibodies as follows: anti-HTLV screening performed on new and non-leucodepleted donations in England (from 2017) and Scotland (from 5 April 2022); anti-HTLV screening performed on all donations in Wales, Northern Ireland, Republic of Ireland, Guernsey and Isle of Man.

In 2022, all donations were screened for treponemal antibodies indicating syphilis including treated past infections as well as some non-sexually acquired treponemal infections such as Yaws and Pinta, rarely seen in the UK.

In 2022, all donations were screened for HEV RNA performed in pools of 24 for England, in pools of 16 for Northern Ireland, Scotland and Wales, and in singletons for Republic of Ireland and also for Wales apheresis donations from 30 November 2022 Wales do not perform confirmatory testing for HEV RNA. England performed HEV RNA screening for the Isle of Man from 24 July 2020.

In 2020 and 2021 the UK blood services collected convalescent plasma from recovered coronavirus patients for use in treatment trials. These donations were subject to the same screening and additional testing requirements as blood and apheresis donations with additional SARS-CoV-2 antibody testing. Donations in England were mainly collected via plasmapheresis.

NHSBT began collecting plasma for medicine from April 2021. These donations were subject to the same donation screening and additional testing requirements as blood and apheresis donations. They will also be screened for Hepatitis A and Parvovirus B19.

Jersey outsourced their donation testing in 2018 due to laboratory refurbishment and have not resumed reporting.

### Additional testing

Other additional (discretionary) tests may be performed including the detection of antibodies to hepatitis B core antigen (anti-HBc), malaria (2001) and *Trypanosoma cruzi* (Chagas disease, 1998) and nucleic acid testing (NAT) for West Nile virus (WNV, 2012). These tests are only performed if information given by the donor suggests that they may have been at risk for these infections.

Anti-HBc testing was previously performed for donations from donors reporting a recent piercing (for example acupuncture, ear, or body piercing and or tattooing), inoculation incident, flexible endoscopy, or history of jaundice or HBV infection. At the end of November 2017, anti-HBc testing ceased for donors with recent endoscopy, piercing and complementary therapies (for example acupuncture). All donors are now screened at least once for anti-HBc from April 2022 with further discretionary anti-HBc performed as required, such as donor reporting an uncertain history of HBV infection or exposure to someone with or recovered from HBV.

Malaria testing is performed where the donor reports a relevant travel history, residency or past infection. Donors are deferred for 4 months post-travel before testing up to 12 months (the deferral was shortened from 6 to 4 months in 2017). Donors are deferred for 4 months post residency before testing at any time. Donors are deferred for 3 years post treatment or malaria illness before testing at any time. Donations repeat reactive for malarial antibodies have been sent for confirmatory reference antibody testing since 30 August 2007. Donations confirmed antibody positive have been tested for malarial DNA by PCR from April 2010.

*T. cruzi* antibody testing was introduced in 1998. There is a 6-month deferral for rural travel, transfusion in South America or because they are of South American origin followed by testing. The deferral was shortened to 4 months in January 2021.

In 2012, WNV NAT testing of donations from donors returning from WNV affected areas was introduced in pools of 6 and replaced the temporary deferral of these donors. Testing continues each year between 01 May and 30 November.

Additional testing indications can be found in the donor selection guidelines on the [Transfusion Guidelines website](#)

### Data collection

Aggregate data on the number of blood donations tested and the number of donations initially reactive is reported to the Epidemiology Unit donation testing scheme each month by the UK blood service's testing centres throughout the UK and the Republic of Ireland and via Scottish

National Blood Transfusion Services (SNBTS) for Scottish blood centres. Disaggregate data on the number of donations repeat reactive with confirmatory testing results are also reported by centres outside England and by the NHSBT Microbiology Services surveillance team for centres in England.

The classification of new and repeat donors used in the Epidemiology Unit donation testing scheme is made by testing centres. This classification of donations tested by donor type is used in particular to estimate the frequency of infection and give an overview of the donations tested.

1. New donors are first time donors who were not known to have ever donated blood in the UK. New donors in the UK (excluding Scotland) and the Republic of Ireland may include 'lapsed' donors for instance repeat donors who have not donated for more than 2 years. In Scotland, all lapsed donors are counted as repeat donors.
2. Repeat donors for most UK centres (excluding Scotland, who include lapsed donors) and the Republic of Ireland, are donors known to have previously donated blood in the UK in the last 2 years, although not all previous donations have necessarily been tested for all markers of infection (for example anti-HTLV testing was first introduced in 2002).

Data on additional (discretionary) testing performed by NHSBT is reported monthly via the following sources:

1. an electronic report from PULSE (the NHSBT national donor database) of the monthly aggregate number of blood donations given an additional test by marker
2. an electronic line listing of donations sent for anti-HBc testing along with screen results
3. an electronic list of the repeat reactive and confirmed positive malaria and *T. cruzi* cases from Microbiology Services surveillance
4. an electronic list of the reference results for samples sent for confirmatory anti-HBc testing from the Microbiology Service Laboratory (MSL)
5. characteristics of donors sent for confirmatory anti-HBc testing are collected via the infected donor scheme and from PULSE

## Infected blood donors

When a marker of infection is detected in a blood donation, the donor is offered a post-test discussion, which may be held in a blood centre or more commonly by telephone. The donor is informed of their positive test results and the clinician explains what these test results mean and ascertains a likely source or risk factor for the infection, if possible. The clinician also discusses any infection control measures, testing and treatment of contacts and advises the donor that they will no longer be able to donate blood. Where appropriate, the donor is referred to the appropriate services for specialist care. Clinicians in blood centres in the UK (excluding Scotland) and Republic of Ireland pass anonymised information about infected blood donors to the Epidemiology Unit infected blood donor surveillance scheme using a standard electronic proforma. This information includes the characteristics of the infected donors (date of birth, gender, first part of



postcode), details of their donating history (if any, with details of their most recent previous donation) and any behaviour that could be associated with the donor's infection. Infected donors are classified by the Epidemiology Unit as newly tested and previously tested for the marker they are found positive for according to detailed information provided by blood centres about all or any previous donations in the UK.

Data from Scotland is supplied on an annual basis.

The classification of infected donors as newly or previously tested is done by the NHSBT and UKHSA Epidemiology Unit:

1. a newly tested donor is one who has not been previously tested for the marker under consideration by the blood transfusion services included in this surveillance
2. a previously tested donor is one who has been previously tested for the marker under consideration by the blood transfusion services included in this surveillance

Note that this classification differs to that used in the donation testing scheme and donor profile data sources (described above) where the donations are classified according to whether the donor has (or has not) donated blood in the last 3 years.

The classification of a seroconverter is made by the NHSBT and UKHSA Epidemiology Unit as a confirmed positive repeat donor with either a previous negative donation within 1 year, or microbiological and or clinical evidence of recent infection AND a previous negative donation within 3 years.

Surveillance data is being continually updated as new or additional information is received. Therefore, some changes between reports may be identified. For example, the seroconverter definition was changed from previously tested within 3 years to within 1 year from 2016 onwards.

Infection was assigned by the unit as recently acquired if there was a previous negative donation within 12 months, or if HBV NAT pick up and/or acute HBV is indicated by confirmatory tests, or if HIV NAT pick up, or HIV avidity test indicated acquired in the last 4 to 5 months or if HCV NAT pick up. For treponema, the clinical history was also reviewed, along with a previous negative donation within 12 months and/or IgM-positive result.

Convalescent plasma donors confirmed to have markers for infection were differentiated by a log number in order to be able to count them separately if required, CP for donations and CPS for samples. Similarly, plasma for medicine donors were differentiated by PFM in the log number.

## Donor profile

Information about individuals donating blood at NHSBT centres (England and north Wales to 1 April 2016) is stored on PULSE, the NHSBT computerised donor information database. Since 1996, this information has been available to the Epidemiology Unit via two sources:

## 1. NHSBT testing centres

Between 1996 and 2000, the number of donations made each month by gender and age group (17 to 24, 25 to 34, 35 to 44, 45 to 54, 55 years and over) for new and repeat donors was reported to the Epidemiology Unit by 7 of the 14 testing centres in England and north Wales. The breakdown from these centres was applied to the total number of donations tested by blood centres in the UK (excluding Scotland) and Republic of Ireland each year to derive the distribution of donations by gender and age group.

## 2. NHSBT Donor Insight

Between 2001 and 2006, aggregate information about individuals donating blood was available from NHSBT Donor Insight (the Market Research and Analysis Department). This information included the proportion of donations made by new and repeat donors by gender, age group (as above) and ethnic group (Asian Other, Black African, Black Caribbean, Black Other, Chinese Indian or Pakistani or Bangladeshi, Mixed, White British, White Other, Other and Not Known). This proportion was applied to the total number of donations tested by blood centres in the UK (excluding Scotland) and Republic of Ireland each year to derive the distribution of donations by gender, age and ethnic group. The data from Market Research and Analysis between 2001 and 2006 were based on a random sample of approximately 1.5 million donation records stored on PULSE.

For 2007 to 2009, data extracts were received for the month of September. These extracts were a complete dataset of every donor who donated during that month at NHSBT. Data included gender, age, donation date, date of most recent previous donation, ethnicity, postcode and new or repeat status. Region was identified using the donor postcode and mapped to strategic health authority regions, to provide more detailed geographical information.

For 2010 onwards, a PULSE extract of all individuals donating during the full year was made available to the NHSBT and UKHSA Epidemiology Unit. These data were used to determine a gender, age, and ethnicity profile of all blood donors. As all blood donors are included these data are able to reflect accurately the characteristics of new and repeat blood donors. In 2021 the method was adapted to include a field for donation made to differentiate PFM from whole blood and apheresis donations.

For both 1 and 2 above, individuals were classified as new donors if they had donated blood for the first time ever or if more than 2 years had elapsed since their last donation.

The NHSBT statistics unit provided data on the CVP donations and samples given on a monthly basis by male and female donors and a total age group and ethnicity breakdown for April 2020 to March 2021.

[Return to Table of Contents](#)

# Transfusion Transmitted Infections

Blood centres in England, Wales and Northern Ireland report investigations of suspected transfusion transmitted infections (TTIs) annually to the NHSBT and UKHSA Epidemiology Unit. For each report, information on the recipient, the recipient's infection, the implicated transfusion, and findings of the investigation are provided using a detailed proforma. Blood centres in Scotland report all incidents to the Microbiology Reference Unit of the SNBTS, and the details and conclusion of each case are passed to the surveillance system annually. NHSBT and UKHSA Epidemiology Unit data is reconciled with the Serious Hazards of Transfusion (SHOT) and all blood service investigations with outcomes are included in the TTI chapter in the SHOT annual report.

## Definition of a transfusion-transmitted Infection

A report of an infection suspected to be due to transfusion was classified as a TTI if the following criteria were met at the end of the investigation:

- the recipient(s) had evidence of infection following transfusion of blood components, and there was no evidence of infection prior to transfusion and no evidence of an alternative source of infection

AND

**Either** at least one component received by the infected recipient was donated by a donor who had evidence of the same infection.

**Or** at least one component received by the infected recipient was shown to contain the agent of infection.

These may be identified because of infection in the recipient where transfusion is the suspected source, and a post-transfusion infection reported to the Blood Services. Alternatively, an infection in a recipient may be identified from lookback investigations which are initiated when a donation from a repeat donor is identified as having markers of infection. Archive samples are retrieved for retrospective testing, which may find a previous donation to also be positive but with markers of infection below the detection level of routine screening. In this case further work will be carried out to identify recipients.

An incident should NOT be reported if:

- the incident involved HCV, HIV or HTLV in recipients who had received transfusions in the UK prior to routine testing (September 1991 for anti-HCV, October 1985 for anti-HIV, August 2002 for anti-HTLV)
- the incident involved HTLV in a recipient identified through the HTLV National Lookback

- the incident involved a transfusion outside UK
- the incident was identified as part of the HEV study

Please note

1. the blood services are rarely able to conduct follow-up investigation of all untested donors implicated in post-transfusion HCV or HIV incidents, and these cases do not contribute to knowledge of the current infection transmission risks of blood transfusions
2. any post-transfusion HTLV infections identified through the HTLV National Lookback are excluded but will be collated, analysed, and published elsewhere, as was done previously with HCV 'lookback'

Data is published annually in the SHOT Annual Report, the UKHSA's Health Protection Report and in the NHSBT and UKHSA Epidemiology Unit annual review. The SHOT Annual Reports can be found on the SHOT website.

[Return to Table of Contents](#)

# Bacterial screening of platelets

Platelets may be manufactured as pooled platelets from whole blood donors (a platelet pool from four donors) or as apheresis platelets where between 1 and 3 apheresis platelet packs may be manufactured from one component donor.

Testing of apheresis platelets by NHSBT began in February 2011 and testing of pooled platelets in June 2011. NHSBT was the last of the UK blood services to introduce bacterial screening of platelets. Bacterial screening was already in place in the other UK blood services. The BacTALERT culture system is used for bacterial screening across all four services but with slightly different sampling methods (see Table A). A surveillance system was put in place within NHSBT in 2011 to report the number of confirmed and indeterminate reactions each month and bacterial species on a quarterly basis.

NHSBT introduced the use of Platelet Additive Solution (PAS) for pooled platelets in 2015; by the end of June 2015 all pooled platelets were manufactured in PAS.

NHSBT produces and screens the largest number of platelets of the four services. Platelets are held for a minimum of 36 hours' post-donation before being sampled. An 8ml sample is inoculated into an aerobic and an anaerobic bottle and placed on the BacTALERT system. If samples are negative after a minimum six-hour incubation, the associated platelet donation can be released as negative-to-date. Platelets are released to stock with a seven-day shelf life since time of donation; prior to the introduction of bacterial screening platelets had a five-day shelf life. Table A shows the sampling volumes and incubation conditions used for the four UK blood services

Table A: Bacterial screening methods used by the UK blood services

UK Service	Time of sampling (hours)	Volume sampled (mls)	Apheresis sample	Time at release (hours)	Length of screening
NHSBT (NHS Blood and Transplant)	36	2 X 8	Post-split	6	Day 7
NIBTS (Northern Ireland Blood Transfusion Service)	36	16	Pre-split	6	Day 7
SNBTS (Scottish National Blood Transfusion Service)	36	2 X 8	Pre-split	6	Day 7
WBS (Welsh Blood Service)	36	2 X 8	Post-split	12	Day 7

Please note:

- Screening methods in Wales changed mid-2018 from testing on day 1 and day 4 to testing on day 2 only

BacTALERT uses a colourimetric system that detects change in pH. A decrease in pH results in an initial reactive report. In some cases, units will have already been transfused. All associated

units will be recalled, and the BacTALERT bottle and any associated packs will be re-tested, and a final result released. Platelet donations are reported as confirmed positive if both the initial screen bottle and the index and or at least 1 associated pack is positive, and the same organism is identified in both. An indeterminate positive result is reported if bacteria are detected in only the initial screen bottle plus no index or associated packs are received or no index packs are received and there's no growth in any associated pack. For indeterminate negative packs, there is no growth from the initial reactive bottle, but a negative result cannot be confirmed because the index pack is not available. Units are confirmed negative if no organism is isolated from the initial reactive bottle and the index pack.

Bacteria isolated from the pack and the bottle are identified to the species level and an assessment will be made of any significance to the donor's health. If a unit has been transfused the transfusion laboratory is notified and asked about any reaction in the recipient.

# Tissue and cord blood donor surveillance

The NHSBT and UKHSA Epidemiology Unit tissue and cell donor scheme collects information on tissue and cord blood donations tested by NHSBT. Tissue donors include deceased and living donors who gave surgical bone. Corneas are donated by deceased donors, some of whom have also donated other tissues; those tested by NHSBT have been included in this scheme since 2012. Since 2013, deceased donors have been reported according to their type of donation for instance those who gave only corneas, those who gave multi-tissue donation including corneas and those who gave multi-tissue excluding corneas.

## Donations tested by NHSBT

### Data collection

Testing for the following mandatory markers of infection is carried out: on antibodies to HIV, HCV, HTLV and treponema; HBsAg and anti-HBc; HBV, HCV and HIV nucleic acid testing (NAT; singletons, triplex). Although not mandatory, donations are also tested for HEV RNA. Disaggregate data on the number of donations tested for mandatory markers of infection, as well as data on additional testing for malaria and *T. cruzi*, are extracted from PULSE (the NHSBT national donor database) on an annual basis. Donations are classified according to donor type (cord blood, living surgical bone or deceased donors [the latter including cornea donors]). The information extracted includes product or component type donated (for instance femoral head, left knee etc.) and gender and date of birth of the donors. Ethnicity of these donors is not currently recorded on PULSE and is therefore not available. In 2012 for the first time, the same data for cornea donors managed and tested by NHSBT were included in this scheme. Some data for stem cell and amnion donors are available but information is limited, thus these donors are not fully integrated into the surveillance scheme and are not reported upon.

### Changes to testing

- there have been several changes to tissue and cord blood donor testing within NHSBT since surveillance began in 2001. HCV, HIV, HBV and HEV NAT (on single samples NOT pools) have been introduced for different donor types at different times
- Cord Blood Donors: HIV NAT and HCV NAT introduced in November 2003. HBV NAT since April 2009. HEV NAT since October 2017
- Deceased Donors: HIV NAT and HCV NAT since 2001. HBV NAT since September 2008. HEV NAT since October 2017

Surgical Bone Donors: Triplex HBV, HCV, HIV NAT since September 2008 (Note: A small number of surgical bone donors require two serology samples [initial and 6-month] where there is

insufficient sample for NAT. Any follow-up samples are excluded from the count of number of donors tested, as they do not represent new donors). HEV NAT since October 2017

Anti-HBc testing has been mandatory for all tissue and cord blood donors since 2006 under EU Commission Directive 2006/17/EC.

#### Infected donors

Follow-up or risk exposure information is received for living surgical bone donors and, where possible, cord blood donors whose donations had markers of infection. As for blood donors, these donors are contacted and asked to telephone the blood centre to discuss their results. The post-test discussion commonly takes place over the telephone and, as for infected blood donors, a behaviour history is sought. For infections detected among deceased donors, an assessment is made to see if any family member or other individual is likely to be at risk before the donor's family is contacted. Risk exposures are not requested for deceased donors. These data are reported to the infected donor surveillance scheme by NHSBT clinicians using standard proformas.

[Return to Table of Contents](#)



# Deceased organ donor surveillance

## Data collection

Organ donor data are derived from the UK Transplant Registry (UKTR) and provided by NHSBT Organ Donation and Transplantation (ODT), and includes donor characteristics, cause of death, reactive test results, and a description of organs donated, and organs transplanted. This information is collected by the NHSBT and hospital staff and submitted to the UKTR either by paper or electronic form.

[Return to Table of Contents](#)

# Risk estimates for blood donors

The estimated risk that a donation entering the UK blood supply is potentially infectious for HBV, HCV or HIV, but not detected on screening because the donation is made during the infectious 'window period', is calculated annually for a rolling three-year period. This statistical process combines information about tests in use by the UK blood services, the infection itself, and data on characteristics of blood donors and donations to produce a point estimate for each infection. An infectious donation may not be detected if a blood donation is made during the infectious 'window period'. This is the period early in the course of infection when the tests in use will not detect the marker of infection. It is also possible that a false negative test result may arise because of issues relating to assay sensitivity other than window period or a blood donation may be erroneously issued as negative due to a sampling, processing, or issuing error. The contribution of these latter two elements is thought to be extremely small and is no longer estimated because of uncertainty around these values.

The estimates for HBV are for acute infections only and do not consider risk due to occult HBV. Hepatitis B core antibody screening for blood donations was rolled out across the UK in 2022 in response to a review carried out by SaBTO. This has already had an impact on increased detection of potentially transmissible hepatitis B virus from donors with occult hepatitis B, which have been removed from the blood supply.

HEV residual risk estimates are not routinely calculated. This is primarily because of uncertainty of the duration of the WP and the fluctuating incidence of HEV in the donor population. This means that the relevance of the traditional incidence WP method across 3-years, as used here for HBV, HCV or HIV, would be questionable for HEV. However, HEV risks have been calculated elsewhere for apheresis and whole blood donors in England between 2016-2020<sup>[1]</sup>.

[1] Harvala H, **Reynolds C, Brailsford S, Davison K**. Fulminant Transfusion-Associated Hepatitis E Virus Infection Despite Screening, England, 2016-2020. *Emerg Infect Dis*. 2022 Sep;28(9):1805-1813

The model used for HBV, HCV and HIV combines data collected in a number of the surveillance schemes. The data required includes:

- number of donations tested (from new and repeat donors), taken from monthly donation testing surveillance
- number of infections detected, by marker (for new and repeat donors), from infected donor surveillance
- number of seroconverting donors - from infected donor surveillance. A seroconverter is defined as an infected repeat donor with either a previous negative donation within one year, or microbiological and or clinical evidence of recent infection with 3-years of their previous screen negative donation. This is taken as 3-years to allow for archive testing to

confirm absence of infection (archives kept for up to 3- years in England). If more than 3- years, the donor is considered lapsed and for the purposes of RR calculations is categorised as a new donor.

- inter-donation interval (IDI) for all donors, taken from NHSBT Donor Insight department data

Parameters used in the model include:

- window period: from expert opinion and literature
- blood donor and donation characteristics: infected donors, NHSBT's Donor Insight department data

The window period risk is calculated as the incidence multiplied by the length of the window period and multiplied by one million. This gives an estimate of the number of potentially infectious donations in one million donations entering the blood supply. Incidence in repeat donors is calculated as the number of observed seroconversions divided by the number of person years exposed, which is estimated from the number of donations from repeat donors multiplied by the average inter-donation interval (IDI) in years. The IDI is created by calculating the difference in days between each donation and the most recent previous donation by the same donor. The average of all these differences is then taken. New donors who returned within the year to donate were excluded, as were donations with IDIs greater than 730 days (two years). Incidence in new donors is derived by adjusting the repeat donor incidence by the relative difference in acute or recent infections among the two groups.

Each year, 95% confidence intervals for the point estimates are calculated by a simulation approach. The same calculation of risk was repeated 1000 times each time using a different set of parameter values. Assumed sampling distributions were intended to reflect the degree of uncertainty in incidence, in average window period length and all other uncertainties in the calculation of the risk of a window period donation. From the resulting 1000 different estimates of risk, we then selected the 2.5 percentile and the 97.5 percentile as the lower and upper limits of a 95% confidence interval.

Parameters and estimates are reviewed annually by the NHSBT and UKHSA Epidemiology Unit. They are then approved annually by the joint UK Blood Transfusion and Tissue Transplantation Services Joint Professional Advisory Committee (JPAC) Standing Advisory Committee on Transfusion Transmitted Infection (SACTTI).

[Return to Table of Contents](#)

# Emerging infections

The Emerging Infections Report (EIR) produced by the NHSBT and UKHSA Epidemiology Unit is distributed monthly. A range of sources are checked for relevant infection issues relating to patient safety and or blood and tissue availability in the UK and collated into a monthly listing. Sources include outbreak alerts, various regular outbreak surveillance reports, journals, websites, and online news resources, listed in more detail below.

The EIR is sent to the Chair of the UK Standing Advisory Committee on Transfusion Transmitted Infection (SACTTI). The chair of SACTTI also receives early warning communications or other reports deemed urgent as they arise.

These monthly listings, alongside other sources of information are reviewed by SACTTI and may lead to further risk assessment and changes to the donor selection guidelines, or other blood safety measures, where necessary. Sources of information for the EIR include:

Outbreak alerts and reports used are listed below;

- ProMED website
- Travel health updates on the [National Travel Health Network and Centre \(NaTHNaC\) website](#).
- Daily and weekly reports from UKHSA EpiIntel with the weekly reports tailored to EIR needs.
- UKHSA All Hazards Situational Awareness Daily Dashboard
- The Infectious Disease Surveillance and Monitoring System for Animal and Human Health: Summary of notable events or incidents of public health significance is a monthly summary of new or emerging infectious disease events that could affect UK public health. It is published monthly by UKHSA, on behalf of the Human Animal Infections and Risk Surveillance group, and can be found on the [Government website](#).
- ECDC Weekly Communicable Disease Threats Report (CDTR) is a weekly bulletin summarising information gathered through epidemic intelligence by ECDC regarding communicable disease threats of concern to the European Union. It also provides updates on the global situation and changes in the epidemiology of communicable diseases with potential to affect Europe including diseases under elimination and is found on the [ECDC website](#) which is also scanned for other surveillance reports.
- Health Protection Report (HPR) is published weekly on the [Government website](#) with routine data and commentary reporting on infectious diseases
- (Currently paused) Health Protection Scotland eWeekly Report is Published weekly on the [Health Protection Scotland website](#) , it is the national health protection bulletin for Scotland and contains current news and articles as well as surveillance reports.
- Morbidity and Mortality Weekly Report (MMWR) is a weekly report prepared by and available on the [Centers for Disease Control and Prevention \(CDC\) website](#)

- Saved PubMed search is set to run monthly for transfusion transmitted infections, blood donors and infection and selected organisms such as hepatitis E, CJD and Chronic Wasting Disease, Babesia
- The [American Association of Blood Banks \(AABB\)](#) online news resource and emails are monitored.
- The Feedly news platform is used to monitor multiple sources, including Outbreak news and various public health resources. An algorithm using key words is used and monitored on a weekly basis.

Journal tables of contents on the following websites are checked:

- [American Journal of Transplantation](#)
- [Clinical Infectious Diseases](#)
- [Emerging Infectious Diseases](#)
- [Epidemiology and Infection](#)
- [Eurosurveillance](#)
- [Journal of Infectious Disease](#)
- [PLOS Pathogens](#)
- [PLOS Neglected Tropical Diseases](#)
- [Transfusion](#)
- [Transfusion medicine](#)
- [Transplant Infectious Disease - Wiley Online Library](#)
- [Vox Sanguinis](#)

Further information is sought from the UKHSA Tuberculosis, Acute Respiratory, Gastrointestinal, Emerging Zoonotic Infections, and Travel and Migrant Health Division (TARZET) where necessary.

The processes for producing the EIR and onward risk assessments are detailed in the JPAC position statements:

[Arrangements in place for monitoring threats to the UK blood supply from new and emerging infections \(May 2023\)](#)

[Horizon Scanning - Management Process Description \(January 2023\)](#)

[Return to Table of Contents](#)