

Detailed Report Actual and Potential Deceased Organ Donation 1 April 2021 - 31 March 2022

Scotland Organ Donation Services Team





Table of Contents

1. Donor outcomes

2. Key rates in potential for organ donation

3. Best quality of care in organ donation

- 3.1 Neurological death testing
- 3.2 Referral to Organ Donation Service
- 3.3 Contraindications
- 3.4 SNOD presence
- 3.5 Authorisation
- 3.6 Solid organ donation

4. Comparative data

- 4.1 Neurological death testing
- 4.2 Referral to Organ Donation Service
- 4.3 SNOD presence
- 4.4 Authorisation

5. PDA data by hospital and unit

6. Paediatric ICU data

- 6.1 Key numbers for PICUs
- 6.2 Neurological death testing in PICUs
- 6.3 Referral to Organ Donation Service in PICUs
- 6.4 Contraindications in PICUs
- 6.5 SNOD presence for patients in PICUs
- 6.6 Authorisation for patients in PICUs
- 6.7 Solid organ donation in PICUs

7. Emergency Department data

- 7.1 Referral to Organ Donation Service
- 7.2 Organ donation discussions

8. Additional Data and Figures

- 8.1 Trust/Board Level Benchmarking
- 8.2 Comparative data for DBD and DCD deceased donors

Appendices

- A.1 Definitions
- A.2 Data description
- A.3 Table and figure description

Further Information

- We acknowledge that the data presented includes the period most significantly impacted by COVID-19 and appreciate that the COVID-19 pandemic affected Trusts/Boards differently across the UK.
- Appendix A.1 contains definitions of terms and abbreviations used throughout this report and summarises the main changes made to the PDA over time.
- The latest Organ Donation and Transplantation Activity Report is available at
- https://www.organdonation.nhs.uk/supporting-my-decision/statistics-about-organ-donation/transplant-activity-report/
 - The latest PDA Annual Report is available at http://www.odt.nhs.uk/statistics-and-reports/potential-donor-audit/
- Please refer any queries or requests for further information to your local Specialist Nurse Organ Donation (SNOD)

Source

NHS Blood and Transplant: UK Transplant Registry (UKTR), Potential Donor Audit (PDA) and Referral Record. Issued May 2022 based on data meeting PDA criteria reported at 9 May 2022.



1. Donor Outcomes

A summary of the number of donors, patients transplanted, average number of organs donated per donor and organs donated.


Data in this section is obtained from the UK Transplant Registry


Between 1 April 2021 and 31 March 2022, the Scotland Organ Donation Services Team facilitated 81 deceased solid organ donors, resulting in 198 patients receiving a transplant. Additional information is shown in Tables 1.1 and 1.2, along with comparison data for 2020/21. Figure 1.1 shows the number of donors and patients transplanted for the previous ten periods for comparison.

Table 1.1 Donors, patie 1 April 2021 -			onor, arch 2021 for comp	arison)
Donor type	Number of donors	Number of patients transplanted	Average number donated per Scotland	•
DBD DCD DBD and DCD	47 (64) 34 (31) 81 (95)	137 (194) 61 (61) 198 (255)	3.3 (3.5) 2.6 (2.6) 3.0 (3.3)	3.5 (3.3) 2.8 (2.6) 3.2 (3.0)

In addition to the 81 proceeding donors there were 35 additional consented donors that did not proceed, 5 where DBD organ donation was being facilitated and 30 where DCD organ donation was being facilitated.

Table 1.2 Organ 1 Apri		l by type, rch 2022 (1 Ap	oril 2020 - 31	March 2021 fo	r compariso	1)	
Donor type	Kidney	Numt Pancreas	per of organs t Liver	transplanted b Heart	Small bowel		
DBD DCD DBD and DCD	84 (114) 47 (51) 131 (165)	1 (8) 0 (2) 1 (10)	38 (53) 13 (8) 51 (61)	9 (13) 1 (1) 10 (14)	7 (15) 2 (2) 9 (17)	0 (0) 0 (0) 0 (0)	







2. Key Rates in

Potential for Organ Donation

A summary of the key rates on the potential for organ donation


Data in this section is obtained from the National Potential Donor Audit (PDA)

This section presents specific percentage measures of potential donation activity for the Scotland Organ Donation Services Team.

Performance in the team has been compared with UK performance in both Figure 2.1 and Table 2.1 using funnel plot boundaries and the Gold, Silver, Bronze, Amber, and Red (GoSBAR) colour scheme. When compared with UK performance, gold represents exceptional, silver represents good, bronze represents average, amber represents below average, and red represents poor performance. See Appendix A.3 for funnel plot ranges used.

It is acknowledged that the PDA does not capture all activity. There may be some patients referred in 2021/22 who are not included in this section onwards because they were either over 80 years of age or did not die in a unit participating in the PDA.

Figure 2.1 Key rates on the potential for organ donation including UK comparison, 1 April 2021 - 31 March 2022











Table 2.1 Key numbers, rates and comparison with national rates,1 April 2021 - 31 March 2022

	Sc	DBI otland	UK	Sc	DCI otland	UK	_	eceased otland	UK
Patients meeting organ donation referral criteria ¹		108	1919 1894		193	5198 4700		287	6767 6258
Referred to Organ Donation Service		108	1894		187	4700		281	0208
Referral rate %	G	100%	99%	G	97%	90%	G	98%	92%
Neurological death tested		93	1530						
Testing rate %	В	86%	80%						
Eligible donors ²		85	1373		132	2972		217	4345
Family approached		75	1239		83	1445		158	2684
Family approached and SNOD present		68	1188		73	1306		141	2494
% of approaches where SNOD present	В	91%	96%	В	88%	90%	В	89%	93%
Authorisation ascertained		50	861		55	902		105	1763
Authorisation rate %	В	67%	69%	В	66%	62%	В	66%	66%
- Expressed opt in		26	522		28	550		54	1072
- Expressed opt in %		90%	95%		82%	90%		86%	92%
- Deemed Authorisation		19	260		23	267		42	527
- Deemed Authorisation %		83%	63%		77%	56%		79%	59%
- Other*		4	78		4	83		8	161
- Other* %		67%	66%		36%	47%		47%	55%
Actual donors (PDA data)		47	787		34	602		81	1389
% of authorised donors that became actual donors		94%	91%		62%	67%		77%	79%
¹ DBD - A patient with suspected neurological death DCD - A patient in whom imminent death is anticipa withdraw treatment has been made and death is anti		•		g ass	isted ven	tilation, a	clinica	al decisio	on to

² DBD - Death confirmed by neurological tests and no absolute contraindications to solid organ donation DCD - Imminent death anticipated and treatment withdrawn with no absolute contraindications to solid organ donation

* Includes patients where nation specific deemed criteria are not met and the patient has not expressed a donation decision in accordance with relevant legislation

Note that a patient that meets both the referral criteria for DBD and DCD organ donation is featured in both the DBD and DCD data but will only be counted once in the deceased donors total





3. Best quality of care

in organ donation

Key stages in best quality of care in organ donation


Data in this section is obtained from the National Potential Donor Audit (PDA)

This section provides information on the quality of care in the Scotland Organ Donation Services Team at the key stages of organ donation. The ambition is that the team misses no opportunity to make a transplant happen and that opportunities are maximised at every stage.

3.1 Neurological death testing

Goal: neurological death tests are performed wherever possible.

Figure 3.1 Number of patients with suspected neurological death, 1 April 2017 - 31 March 2022



Patients not tested Patients tested

Table 3.1 Reasons given for neurological death tests not being performed, 1 April 2021 - 31 March 2022

Biochemical/endocrine abnormality	Scotland	UK 21
Clinical reason/Clinician's decision	7	48
Continuing effects of sedatives	-	10
Family declined donation	-	20
Family pressure not to test	1	27
Hypothermia	-	2
Inability to test all reflexes	-	17
Medical contraindication to donation	-	7
Other	3	37
Patient had previously expressed a wish not to donate	1	1
Patient haemodynamically unstable	3	162
Pressure of ICU beds	-	8
SN-OD advised that donor not suitable	-	10
Treatment withdrawn	-	14
Unknown	-	5
Total	15	389
If 'other', please contact your local SNOD or CLOD for more info	rmation, if requ	iired.



3.2 Referral to Organ Donation Service

Goal: Every patient who meets the referral criteria should be identified and referred to the Organ Donation Service, as per NHS Blood and Transplant (NHSBT) Best Practice Guidance on timely identification and referral of potential organ donors.¹

Aim: There should be no purple on the following charts.




Figure 3.2 Number of patients meeting referral criteria, 1 April 2017 - 31 March 2022

Table 3.2 Reasons given why patient not referred to SNOD,1 April 2021 - 31 March 2022

	DBD		DCD	
	Scotland	UK	Scotland	UK
Clinician assessed that patient was unlikely to become asystolic within 4 hours	-	-	-	2
Coroner / Procurator Fiscal reason	-	_	-	1
Family declined donation after neurological testing	_	2	_	-
amily declined donation following decision to remove treatment	-	-	1	7
Family declined donation prior to neurological testing	-	1	1	1
Medical contraindications	-	3	-	78
Not identified as potential donor/organ donation not considered	-	12	3	275
Dther	-	1	1	51
Patient had previously expressed a wish not to donate	-	1	-	-
Pressure on ICU beds	-	-	-	5
Reluctance to approach family	-	-	-	4
Thought to be medically unsuitable	-	2	-	65
Jncontrolled death pre referral trigger	-	3	-	9
Total	-	25	6	498



3.3 Contraindications

In 2021/22 there were 29 potential donors in the Scotland Organ Donation Services team with an ACI reported, 7 DBD and 24 DCD donors. Please note, the number of potential DBD and DCD donors with an ACI reported may not equal the total stated as a patient can meet potential donor criteria for both DBD and DCD donation.



3.4 SNOD presence

Goal: A SNOD should be present during the formal family approach as per NHS Blood and Transplant (NHSBT) Best Practice Guidance.²

Aim: There should be no purple on the following charts.

In the UK, in 2021/22, when a SNOD was not present for the approach to the family to discuss organ donation, DBD and DCD authorisation rates were 35% and 19%, respectively, compared with DBD and DCD consent rates of 71% and 67%, respectively, when a SNOD was present.

Within the Boards in the team, when a SNOD was not present for the approach to the family to discuss organ donation, DBD and DCD authorisation rates were 29% and 10%, respectively, compared with DBD and DCD authorisation rates of 71% and 74%, respectively, when a SNOD was present.

Every approach to those close to the patient should be planned with the multidisciplinary team (MDT), should involve the SNOD and should be clearly planned taking into account the known wishes of the patient. The NHS Organ Donor Register (ODR) should be checked in all cases of potential donation and this information must be discussed with the family as it represents the eligible donor's legal consent to donation.




Figure 3.3 Number of families approached by SNOD presence, 1 April 2017 - 31 March 2022

¹ NHS Blood and Transplant, 2012.

Timely Identification and Referral of Potential Organ Donors - A Strategy for Implementation of Best Practice [accessed 9 May 2022]

² NHS Blood and Transplant, 2013.

Approaching the Families of Potential Organ Donors – Best Practice Guidance [accessed 9 May 2022]



3.5 Authorisation

In 2021/22 the DBD and DCD authorisation rates in the team were 67% and 66%, respectively.

Figure 3.4 Number of families approached, 1 April 2017 - 31 March 2022




Table 3.3 Reasons given why authorisation was not ascertained,1 April 2021 - 31 March 2022

	DBD D		DCD)
	Scotland	UK	Scotland	UK
Family concerned donation may delay the funeral	-	-	-	2
Family concerned other people may disapprove/be offended	1	3	-	1
Family concerned that organs may not be transplantable	-	1	2	4
Family did not believe in donation	-	10	2	13
Family did not want surgery to the body	-	35	3	46
Family divided over the decision	1	13	-	11
Family felt it was against their religious/cultural beliefs	-	39	-	24
Family felt patient had suffered enough	1	26	3	42
Family felt that the body should be buried whole (unrelated to	-	16	-	9
religious/cultural reasons)				
Family felt the length of time for the donation process was too	1	15	6	85
long				
Family had difficulty understanding/accepting neurological testing	-	2	-	-
Family wanted to stay with the patient after death	1	2	1	5
Family were not sure whether the patient would have agreed to	3	35	2	64
donation				
Other	1	20	1	45
Patient had previously expressed a wish not to donate	13	125	6	148
Patient had registered a decision to Opt Out	1	23	-	20
Strong refusal - probing not appropriate	2	13	2	23
Total	25	378	28	542
If 'other', please contact your local SNOD or CLOD for more infor	mation, if requ	ired.		



3.6 Solid organ donation

Goal: NHSBT is committed to supporting transplant units to ensure as many organs as possible are safely transplanted.

	DBD)	DCE	`
	Scotland	, UK		, UK
Clinical - Absolute contraindication to organ donation	-	4	-	
Clinical - Considered high risk donor	1	3	-	6 5
Clinical - No transplantable organ	-	5	4	21
Clinical - Organs deemed medically unsuitable by recipient	-	25	2	70
centres				
Clinical - Organs deemed medically unsuitable on surgical	-	8	2	4
nspection				
Clinical - Other	1	3	1	10
Clinical - PTA post WLST	-	-	12	13
Clinical - Patient actively dying	-	6	-	14
Clinical - Patient's general medical condition	-	-	-	6 5
Clinical - Positive virology	1	3	-	5
Consent / Auth - Coroner/Procurator fiscal refusal	-	11	-	1
Consent / Auth - Known wish not to donate	-	1	-	1
Consent / Auth - NOK withdraw consent / authorisation	-	5	-	8 2
Consent / Auth - Other	-	-	-	2
ogistical - No critical care bed available	-	-	-	1
ogistical - Other	-	-	-	1
Fotal	3	74	21	30



4. Comparative Data

A comparison of performance in your team with national data

Data in this section is obtained from the National Potential Donor Audit (PDA)


This section compares the quality of care in the key areas of organ donation in the Scotland Organ Donation Services team with the UK rate using funnel plots. The UK rate is shown as a green dashed line and the funnel shape is formed by the 95% and 99.8% confidence limits around the UK rate. The confidence limits reflect the level of precision of the UK rate relative to the number of observations. Performance in the team is indicated by a black cross. The Gold, Silver, Bronze, Amber, and Red colour scheme is used to indicate whether performance in the team, when compared to UK performance, is exceptional (gold), good (silver), average (bronze), below average (amber) or poor (red).

It is important to note that the differences in patient mix have not been accounted for in these plots. Further to these, separate funnel plots for DBD and DCD rates are presented in Section 8.

4.1 Neurological death testing

Goal: neurological death tests are performed wherever possible.

Figure 4.1 Funnel plot of neurological death testing rate, 1 April 2021 - 31 March 2022



When compared with UK performance, the performance within the Boards in the team was average (bronze) for neurological death testing.



4.2 Referral to Organ Donation Service

Goal: Every patient who meets the referral criteria should be identified and referred to NHSBT's Organ Donation Service, as per NHS Blood and Transplant (NHSBT) Best Practice Guidance on timely identification and referral of potential organ donors.¹




Figure 4.2 Funnel plot of deceased donor referral rate, 1 April 2021 - 31 March 2022

When compared with UK performance, the performance within the Boards in the team was exceptional (gold) for referral of potential organ donors to NHS Blood and Transplant's Organ Donation Service.



4.3 SNOD presence

Goal: A SNOD should be present during the formal family approach as per NHS Blood and Transplant (NHSBT) Best Practice Guidance.²





Figure 4.3 Funnel plot of SNOD presence rate, 1 April 2021 - 31 March 2022

When compared with UK performance, the performance within the Boards in the team was average (bronze) for Specialist Nurse presence when approaching families to discuss organ donation.



4.4 Authorisation

Figure 4.4 Funnel plot of authorisation rate, 1 April 2021 - 31 March 2022



When compared with UK performance, the authorisation rate within the Boards in the team was average (bronze).



5. PDA data by hospital and unit

A summary of key numbers and rates from the PDA by hospital and unit where patient died

Data in this section is obtained from the National Potential Donor Audit (PDA)

Tables 5.1 and 5.2 show the key numbers and rates for patients who met the DBD and/or DCD referral criteria, respectively. Percentages have been excluded where numbers are less than 10.

Table 5.1 Patients who met the DBD referral criteria - key numbers and rates,1 April 2021 - 31 March 2022

neur dea		^D atients tested	Neurological death testing rate (%)	Patients referred	DBD referral rate (%)	Patients confirmed dead by neurological testing	Eligible DBD donors	Eligible DBD donors whose family were approached	Approaches where SNOD involved	SNOD presence rate (%)	Authorisation ascertained	Authorisation rate (%)	Actual DBD and DCD donors from eligible DBD donors
Golde	en Jubilee N		lospital										
ł	1	0	-	1	-	0	0	0	0	-	0	-	0
NHS	Ayrshire ar			7		7	0		0		0		
ł	7	7	-	7	-	7	6	4	3	-	3	-	3
NHS	Borders 1	1		1		1	1	1	1		1		1
ł			-	I	-	I	I	I	I	-	I	-	1
NHS	Dumfries a	nd Gallov 3	vay -	3	_	3	2	2	2	_	2	_	2
i		5	-	5	-	5	2	2	2	-	2	-	2
NHS	Fife 9	7	_	9	_	7	7	7	7	_	5	-	5
i				0			•	,			0		Ŭ
	Forth Valle	y 7	58	12	100	7	7	4	4	-	2	-	2
i													
	<i>Grampian</i> 17	14	82	17	100	14	14	12	12	100	9	75	8
	Orrester Ol	0	Chuda										
NHS	Greater Gla 23	21 21	91	23	100	21	19	18	16	89	12	67	11
	Highland												
NIIS	4	4	-	4	-	4	4	4	2	-	2	-	1
NHS	Lanarkshire	•											
	12	, 11	92	12	100	11	11	11	10	91	7	64	7
NHS	Lothian												
	11	11	100	11	100	11	10	9	8	-	6	-	6
NHS	Tayside												
	8	7	-	8	-	7	4	3	3	-	1	-	1

Table 5.2 Patients who met the DCD referral criteria - key numbers and rates,1 April 2021 - 31 March 2022

Patients for whom imminent death was anticipated	Patients referred	DCD referral rate (%)	Patients for whom treatment was withdrawn	Eligible DCD donors	Eligible DCD donors whose family were approached	Approaches where SNOD involved	SNOD presence rate (%)	Authorisation ascertained	Authorisation rate (%)	Actual DCD donors from eligible DCD donors
Golden Jubilee	National Hos	pital								
6	6	-	6	6	4	4	-	4	-	1
NHS Ayrshire a	and Arran									
15	15	100	15	9	6	6	-	2	-	2



Table 5.2 Patients who met the DCD referral criteria - key numbers and rates, 1 April 2021 - 31 March 2022

Patients for whom imminent death was anticipated	Patients referred	DCD referral rate (%)	Patients for whom treatment was withdrawn	Eligible DCD donors	Eligible DCD donors whose family were approached	Approaches where SNOD involved	SNOD presence rate (%)	Authorisation ascertained	Authorisation rate (%)	Actual DCD donors from eligible DCD donors
NHS Borders 1	1	-	1	0	0	0	-	0	-	0
NHS Dumfries ar 4	nd Galloway 4	-	4	3	2	2	-	2	-	2
NHS Fife 17	17	100	16	16	6	6	-	5	-	4
NHS Forth Valley 13	, 13	100	13	8	3	3	-	0	-	0
NHS Grampian 23	21	91	21	19	12	10	83	8	67	4
NHS Greater Gla 45	sgow & Cly 44	rde 98	45	26	16	13	81	9	56	5
NHS Highland 5	5	-	5	3	1	1	-	1	-	1
NHS Lanarkshire 7	6	-	5	5	2	1	-	1	-	1
NHS Lothian 45	43	96	40	26	23	20	87	17	74	8
NHS Tayside 12	12	100	12	11	8	7	-	6	_	6

Tables 5.1 and 5.2 show the hospital where the patient died. However, it is acknowledged that there are some occasions where a patient is referred in an Emergency Department but moves to a critical care unit. In total for the team in 2021/22 there were 2 such patients. For more information regarding the Emergency Department please see Section 7.



6. Paediatric ICU data

A summary of key numbers for paediatric ICUs

Data in this section is obtained from the National Potential Donor Audit (PDA)

End of life care guidance and practice for paediatric patients does differ and care of the family unit as a whole is a core key principle. Paediatric Intensive Care Units (PICU) systems should never prevent families being offered the opportunity to donate if this is a possibility.

This section provides information on the quality of care for patients that died in PICUs in the Scotland Organ Donation Services team at the key stages of organ donation. The ambition is that your PICU misses no opportunity to make a transplant happen and that opportunities are maximised at every stage.

6.1 Key numbers for PICUs

Table 6.1 PICU key numbers comparison with national rates,1 April 2021 - 31 March 2022

	DBI)	DCE)	Deceased	donors
	Scotland	UK	Scotland	UK	Scotland	UK
Patients meeting organ donation referral criteria ¹	0	58	3	178	3	211
Referred to Organ Donation Service	0	56	2	156	2	187
Referral rate %		97%		88%		89%
Neurological death tested	0	33				
Testing rate %		57%				
Eligible donors ²	0	31	3	138	3	169
Family approached	0	25	3	53	3	78
Family approached and SNOD present	0	24	1	39	1	63
% of approaches where SNOD present		96%		74%		81%
Authorisation ascertained	0	15	1	21	1	36
Authorisation rate %		60%		40%		46%
Actual donors (PDA data)	0	15	1	15	1	30
% of authorised donors that became actual donors		100%		71%		83%

¹ DBD - A patient with suspected neurological death

DCD - A patient in whom imminent death is anticipated, ie a patient receiving assisted ventilation, a clinical decision to withdraw treatment has been made and death is anticipated within 4 hours

² DBD - Death confirmed by neurological tests and no absolute contraindications to solid organ donation DCD - Imminent death anticipated and treatment withdrawn with no absolute contraindications to solid organ donation


Note that a patient that meets both the referral criteria for DBD and DCD organ donation is featured in both the DBD and DCD data but will only be counted once in the deceased donors total



6.2 Neurological death testing in PICUs

Goal: neurological death tests are performed wherever possible.

Figure 6.1 Number of patients with suspected neurological death in PICUs, 1 April 2017 - 31 March 2022



Patients not tested Patients tested

Table 6.2 Reasons given for neurological death test PICUs, 1 April 2021 - 31 March 2022	s not being performed	d in
	Scotland	UK
Biochemical/endocrine abnormality	-	3
Clinical reason/Clinician's decision	-	2
Continuing effects of sedatives	-	1
Family declined donation	-	4
Family pressure not to test	-	4
Other	-	3
Patient haemodynamically unstable	-	6
SN-OD advised that donor not suitable	-	1
Treatment withdrawn	-	1
Total	-	25



6.3 Referral to Organ Donation Service in PICUs

Goal: Every patient who meets the referral criteria should be identified and referred to the Organ Donation Service, as per NHS Blood and Transplant (NHSBT) Best Practice Guidance on timely identification and referral of potential organ donors.¹

Aim: There should be no purple on the following charts.




Figure 6.2 Number of patients meeting referral criteria in PICUs, 1 April 2017 - 31 March 2022

Table 6.3 Reasons given why patient not referred to Organ Donation Service in PICUs,1 April 2021 - 31 March 2022

	DBD		DCD		
	Scotland	UK	Scotland	UK	
Coroner / Procurator Fiscal reason	-	-	-	1	
amily declined donation after neurological testing	-	1	-	-	
Family declined donation prior to neurological testing	-	1	1	1	
Not identified as potential donor/organ donation not considered	-	-	-	3	
Other	-	-	-	13	
Thought to be medically unsuitable	-	-	-	4	
Total	-	2	1	22	



6.4 Contraindications in PICUs

In 2021/22 there were no potential donors in the Scotland Organ Donation Services team with an ACI reported.




6.5 SNOD presence for patients in PICUs

Goal: A SNOD should be present during the formal family approach as per NHS Blood and Transplant (NHSBT) Best Practice Guidance.²

Aim: There should be no purple on the following charts.

Figure 6.3 Number of families of PICU patients approached by SNOD presence, 1 April 2017 - 31 March 2022







6.6 Authorisation for patients in PICUs

In 2021/22 less than 10 families of eligible donors, facilitated in the PICU, were approached to discuss organ donation in the team therefore authorisation rates are not presented.

Figure 6.4 Number of families of PICU patients approached, 1 April 2017 - 31 March 2022





Authorisation not ascertained Authorisation ascertained

Authorisation not ascertained Authorisation ascertained

Table 6.5 Reasons given why authorisation was not ascertained for PICU patients,1 April 2021 - 31 March 2022

	DBD		DCD	
	Scotland	UK	Scotland	UK
Family concerned donation may delay the funeral	-	-	-	1
Family did not believe in donation	-	1	1	2
Family did not want surgery to the body	-	3	-	7
Family felt it was against their religious/cultural beliefs	-	1	-	-
Family felt patient had suffered enough	-	1	-	9
Family felt that the body should be buried whole (unrelated to	-	1	-	-
religious/cultural reasons)				
Family felt the length of time for the donation process was too	-	-	-	2
long				
Family wanted to stay with the patient after death	-	_	_	1
Other	-	2	_	6
Patient had previously expressed a wish not to donate	_	1	_	-
Strong refusal - probing not appropriate	_	-	1	4
Total	_	10	2	32
			-	72
If 'other', please contact your local SNOD or CLOD for more info	rmation if requ	irod		
		neu.		



6.7 Solid organ donation in PICUs

Goal: NHSBT is committed to supporting transplant units to ensure as many organs as possible are safely transplanted.

1 April 2021 - 31 March 2022				
	DBC)	DCD)
	Scotland	UK	Scotland	UK
Clinical - Organs deemed medically unsuitable by recipient centres	-	-	-	2
Clinical - PTA post WLST	-	-	-	2
Consent / Auth - Coroner/Procurator fiscal refusal	-	-	-	1
Consent / Auth - NOK withdraw consent / authorisation	-	-	-	1
lotal	-	-	-	6



7. Emergency Department data

A summary of key numbers for Emergency Departments


Data in this section is obtained from the National Potential Donor Audit (PDA)

Most patients who go on to become organ donors start their journey in the emergency department (ED). Deceased donation is important, not just for those people waiting on the transplant list, but also because many people in the UK have expressed a wish in life to become organ donors after their death. The overarching principle of the NHSBT Organ donation and Emergency Department strategy ³ is that best quality of care in organ donation should be followed irrespective of the location of the patient within the hospital at the time of death.

7.1 Referral to Organ Donation Service


Goal: No one dies in your ED meeting referral criteria and is not referred to NHSBT's Organ Donation Service. Aim: There should be no blue on the following chart.






7.2 Organ donation discussions

Goal: No family is approached in ED regarding organ donation without a SNOD present. Aim: There should be no red on the following chart.





SNOD not present SNOD present

³ NHS Blood and Transplant, 2016. Organ Donation and the Emergency Department [accessed 9 May 2022]



8. Additional data and figures

Key numbers and rates on the potential for organ donation

Data in this section is obtained from the National Potential Donor Audit (PDA)

8.1 Trust/Board Level Benchmarking

Trust/Board levels were reallocated in July 2018 using the average number of donors in 2016/17 and 2017/18, Table 8.1 shows the criteria used and how many Trusts/Boards belong to each level.

Table 8.1	Trust/Board level categories	
		Number of Trusts Boards in each level
Level 1	12 or more (\geq 12) proceeding donors per year	35
Level 2	6 or more but less than 12 (\geq 6 to <12) proceeding donors per year	45
Level 3	More than 3 but less than 6 (>3 to <6) proceeding donors per year	47
Level 4	3 or less (\leq 3) proceeding donors per year	41

Tables 8.2 and 8.3 show the national DBD and DCD key numbers and rates for the UK by Trust/Board level, to aid in comparison with equivalent Trusts/Boards. Note that percentages have been excluded where numbers are less than 10.

			rch 202	22			oard level,					
Patients where neurological death was suspected	Patients tested	Neurological death testing rate (%)	Patients referred	DBD referral rate (%)	Patients confirmed dead by neurological testing	Eligible DBD donors	Eligible DBD donors whose family were approached	Approaches where SNOD present	SNOD presence rate (%)	Authorisation ascertained	Authorisation rate (%)	Actual DBD and DCD donors from eligible DBD donors
1044	840	80	1034	99	827	748	679	646	95	470	69	434
455	361	79	445	98	355	318	284	274	96	187	66	173
286	225	79	282	99	221	208	189	184	97	147	78	128
134	104	78	133	99	103	99	87	84	97	57	66	52
	where neurological death was suspected 1044 455 286	where neurological death was suspected 1044 840 455 361 286 225	where neurological death was suspected 255 361 79 286 225 79	where neurological death was suspectedNeurological death testing testedNeurological death testing rate (%)Patients referred10448408010344553617944528622579282	where neurological death was suspectedNeurological death testing testedDBD referral referred rate (%)10448408010349945536179445982862257928299	where neurological death was suspectedNeurological death testing rate (%)DBD patients referradConfirmed dead by neurological referrad10448408010349982745536179445983552862257928299221	where neurological death was suspectedNeurological death testing testedNeurological death testing rate (%)DBD Patients referred rate (%)Confirmed deadh testing neurological rate (%)Eligible DBD neurological neurological referred rate (%)DBD referral rate (%)Eligible testing donors10448408010349982774845536179445983553182862257928299221208	where neurological death was suspectedNeurological 	where neurological death was suspectedNeurological death testing rate (%)Neurological PatientsNeurological patientsDBD referral referral referral rate (%)Eligible referral referral referral rate (%)Eligible referral referral referral referralEligible referral reformal referralEligible referral reformal referral referralEligible referral reformal referral referralEligible referral reformal referral referralDBD referral referral referral referralEligible referral reformal referral<	where neurological death was suspectedNeurological death testing testedNeurological death testing rate (%)Neurological PatientsNeurological referred <th< td=""><td>where neurological death was suspectedNeurological death testing rate (%)Neurological Patients referred referred rate (%)DBD referred rate (%)Eligible referral referred restrict (%)Eligible referred restrict (%)DBD family were about esting restrict (%)Eligible referred restrict (%)NOD referred restrict (%)Approaches restrict (%)SNOD restrict esting restrict (%)10448408010349982774867964695470455361794459835531828427496187286225792829922120818918497147</td><td>where neurological death was suspectedNeurological death testing rate (%)Neurological Patients referred referred referredDBD referred referred referredEligible deadh y absolutionEligible DBD donors approached approachedSNOD presentAuthorisation rate (%)Authorisation rate (%)10448408010349982774867964695470694553617944598355318284274961876628622579282992212081891849714778</td></th<>	where neurological death was suspectedNeurological death testing rate (%)Neurological Patients referred referred rate (%)DBD referred rate (%)Eligible referral referred restrict (%)Eligible referred restrict (%)DBD family were about esting restrict (%)Eligible referred restrict (%)NOD referred restrict (%)Approaches restrict (%)SNOD restrict esting restrict (%)10448408010349982774867964695470455361794459835531828427496187286225792829922120818918497147	where neurological death was suspectedNeurological death testing rate (%)Neurological Patients referred referred referredDBD referred referred referredEligible deadh y absolutionEligible DBD donors approached approachedSNOD presentAuthorisation rate (%)Authorisation rate (%)10448408010349982774867964695470694553617944598355318284274961876628622579282992212081891849714778

Table 8.3 National DCD key numbers and rate by Trust/Board level,1 April 2021 - 31 March 2022

	Patients for whom imminent death was anticipated	Patients referred	DCD referral rate (%)	Patients for whom treatment was withdrawn	Eligible DCD donors	Eligible DCD donors whose family were approached	Approaches where SNOD present	SNOD presence rate (%)	Authorisation ascertained	Authorisation rate (%)	Actual DCD donors from eligible DCD donors
Level 1	2391	2224	93	2289	1498	818	728	89	513	63	347
Level 2	1451	1261	87	1383	750	335	310	93	197	59	137
Level 3	915	827	90	882	464	184	174	95	130	71	76
Level 4	441	388	88	425	260	108	94	87	62	57	42



8.2 Comparative data for DBD and DCD deceased donors

Funnel plots are presented in Section 4 showing performance in the team against the UK rate for deceased organ donation. The following funnel plots present data for DBD and DCD donors separately.





Figure 8.1 Funnel plots of referral rates, 1 April 2021 - 31 March 2022

When compared with UK performance, the performance within the Boards in the team was exceptional (gold) for referral of potential DBD organ donors and exceptional (gold) for referral of potential DCD organ donors to NHS Blood and Transplant's Organ Donation Service.





When compared with UK performance, the performance within the Boards in the team was average (bronze) and average (bronze) for Specialist Nurse presence in approaches to families of eligible DBD and DCD donors, respectively.






Figure 8.3 Funnel plots of authorisation rates, 1 April 2021 - 31 March 2022

When compared with UK performance, the authorisation rate within the Boards in the team was average (bronze) and average (bronze) for DBD and DCD donors, respectively.



Appendices

Appendix A.1 Definitions

Potential Donor Audit Definitions

Potential Donor Audit inclusion criteria	1 October 2009 – 31 March 2010 All deaths in critical care in patients aged 75 and under, excluding cardiothoracic intensive care units 1 April 2010 – 31 March 2013 All deaths in critical and emergency care in patients aged 75 and under, excluding cardiothoracic intensive care units 1 April 2013 onwards All deaths in critical and emergency care in patients aged 80 and under (prior to 81st birthday)	

Donors after brain death (DBD) definitions

Suspected Neurological Death	A patient who meets all of the following criteria: invasive ventilation, Glasgow Coma Scale 3 not explained by sedation, no respiratory effort, fixed pupils, no cough or gag reflex. Excluding those not tested due to reasons 'cardiac arrest despite resuscitation', 'brainstem reflexes returned', 'neonates – below 37 weeks corrected gestational age'. Previously referred to as brain death
Neurological death tested	Neurological death tests performed to confirm and diagnose death
DBD referral criteria	A patient with suspected neurological death
Specialist Nurse Organ Donation or Organ Donation Services Team Member (SNOD)	A member of Organ Donation Services Team including: Team Manager, Specialist Nurse Organ Donation, Specialist Requester, Donor Family Care Nurse
Referred to Specialist Nurse – Organ Donation	A patient with suspected neurological death referred to a SNOD. A referral is the provision of information to determine organ donation suitability. NICE CG135 (England) : Triggers for clinicians to refer a potential donor are a plan to withdraw life sustaining treatment or a plan to perform neurological death tests
Potential DBD donor	A patient with suspected neurological death
Absolute contraindications	Absolute medical contraindications identified in assessment which clinically preclude organ donation as per NHSBT criteria (POL188) Absolute medical contraindications to donation are listed here: https://nhsbtdbe.blob.core.windows.net/umbraco-assets-corp/17160/ clinical-contraindications-to-approaching-families-for-possible-organ-donati on-pol188.pdf
Eligible DBD donor	A patient confirmed dead by neurological death tests, with no absolute medical contraindications to solid organ donation
Donation decision conversation	Family of eligible DBD asked to make or support patient's organ donation decision - This includes clarifying an opt out decision
Consent/Authorisation ascertained	Family supported opt in decision, deemed consent/authorisation, or where applicable the family or nominated/appointed representative gave consent/authorisation for organ donation
Actual donors: DBD	Patients who became actual DBD donors following confirmation of neurological death, as reported through the PDA (80 years and below). At least one organ donated for the purpose of transplantation (includes organs retrieved for transplant however used for research)
Actual donors: DCD	Patients who became actual DCD donors following confirmation of neurological death, as reported through the PDA (80 years and below). At least one organ donated for the purpose of transplantation (includes organs retrieved for transplant however used for research)



Neurological death testing rate	Percentage of patients for whom neurological death was suspected who were tested
Referral rate	Percentage of patients for whom neurological death was suspected who were referred to the SNOD
Donation decision conversation rate	Percentage of eligible DBD families or nominated/appointed representatives who were asked to make or support an organ donation decision - This includes clarifying an opt out decision
Consent/Authorisation rate	Percentage of donation decision conversations where consent/authorisation was ascertained
SNOD presence rate	Percentage of donation decision conversations where a SNOD was present (includes telephone and video call conversations)
Consent/Authorisation rate where SNOD was present	Percentage of donation decision conversations where a SNOD was present and consent/authorisation for organ donation was ascertained (as above)

Donors after circulatory death (DCD) definitions

Γ

Imminent death anticipated	A patient, not confirmed dead using neurological criteria, receiving invasive ventilation, in whom a clinical decision to withdraw treatment has been made and a controlled death is anticipated within a time frame to allow donation to occur (as determined at time of assessment)
DCD referral criteria	A patient for whom imminent (controlled) death is anticipated following withdrawal of life sustaining treatment (as defined above)
Specialist Nurse Organ Donation or Organ Donation Services Team Member (SNOD)	A member of Organ Donation Services Team including: Team Manager, Specialist Nurse Organ Donation, Specialist Requester, Donor Family Care Nurse
Referred to SNOD	A patient for whom imminent death is anticipated who was referred to a SNOD. A referral is the provision of information to determine organ donation suitability NICE CG135 (England) : Triggers for clinicians to refer a potential donor are a plan to withdraw life sustaining treatment or a plan to perform neurological death tests
Potential DCD donor	A patient who had treatment withdrawn and imminent death was anticipated within a time frame to allow donation to occur.
Absolute contraindications	Absolute medical contraindications identified in assessment which clinically preclude organ donation as per NHSBT criteria (POL188). Absolute medical contraindications to donation are listed here: https://nhsbtdbe.blob.core.windows.net/umbraco-assets-corp/17160/ clinical-contraindications-to-approaching-families-for-possible-organ-donati on-pol188.pdf
Eligible DCD donor to be assessed	A patient who had treatment withdrawn and imminent (controlled) death was anticipated, with no absolute medical contraindications to solid organ donation.
DCD exclusion criteria	DCD specific criteria determine a patient's suitability to donation when there are no absolute medical contraindications (see absolute contraindications documentation above)
DCD screening process	Process by which an organ may be screened with a local and national transplant centre to determine suitability of organs for transplantation
Medically suitable eligible DCD donor	An eligible DCD donor to be assessed considered to be medically suitable for donation (i.e. no DCD exclusions and not deemed unsuitable by the screening process)
Donation decision conversation	Family of medically suitable eligible DCD donor who were asked to make or support patient's organ donation decision - This includes clarifying an opt out decision.
Consent/Authorisation ascertained	Family supported opt in decision, deemed consent/authorisation, or where applicable the family or nominated/appointed representative gave consent/authorisation for organ donation



Actual DCD	DCD patients who became actual DCD as reported through the PDA (80 years and below). At least one organ donated for the purpose of transplantation (includes organs retrieved for transplant however used for research)
Referral rate	Percentage of patients for whom imminent (controlled) death was anticipated who were referred to the SNOD
Donation decision conversation rate	Percentage of medically suitable eligible DCD families or nominated/appointed representatives who were asked to make or support an organ donation decision - This includes clarifying an opt out decision
Consent/Authorisation rate	Percentage of donation decision conversations where consent/authorisation was ascertained.
SNOD presence rate	Percentage of donation decision conversations where a SNOD was present (includes telephone and video call conversations).
Consent/Authorisation rate where SNOD was present	Percentage of donation decision conversations where a SNOD was present and consent/authorisation for organ donation was ascertained (as above).

Deemed Consent/Authorisation

Deemed consent applies if a person who died in Wales, Jersey or England has not expressed an organ donation decision either to opt in or opt out or nominate/appoint a representative, is aged 18 or over, has lived in the country in which they died for longer than 12 months and is ordinarily resident there, and had the capacity to understand the notion of deemed consent for a significant period before their death.

Deemed authorisation applies if a person who died in Scotland has not expressed, in writing, an organ donation decision either to opt in or opt out, is aged 16 or over, has lived in Scotland for longer than 12 months and is ordinarily resident there, and had the capacity to understand the notion of deemed authorisation for a significant period before their death. Note that, in Scotland, a patient who has verbally expressed an opt in decision is included as a deemed authorisation, whereas a patient who has verbally expressed an opt out decision is not included.

Consent/Authorisation groups

Expressed opt in	Patient had expressed an opt in decision. Opt in decisions can be expressed in writing or via the ODR in all nations and verbal opt in decisions are also included in Wales, England and Jersey. Verbally expressed opt in decisions are not included in Scotland
Deemed consent/authorisation	Patient meets deemed criteria specific to each nation as described above. In Scotland, this includes patients who have verbally expressed a decision to opt in
Expressed opt out	Patient had expressed an opt out decision. Opt out decisions can be expressed verbally, in writing or via the ODR in all nations
Other	Patient has expressed no decision or deemed criteria are not met. Paediatric patients are included in this group

UK Transplant Registry (UKTR) definitions

Donor type	Type of donor: Donation after brain death (DBD) or donation after circulatory death (DCD)
Number of actual donors	Total number of donors reported to the UKTR
Number of patients transplanted	Total number of patients transplanted from these donors
Organs per donor	Number of organs donated divided by the number of donors.
Number of organs transplanted	Total number of organs transplanted by organ type



Appendix A.2 Data Description

This report provides a summary of data relating to potential and actual organ donors as recorded by NHS Blood and Transplant via the Potential Donor Audit (PDA), the accompanying Referral Record, and the UK Transplant Registry (UKTR) for the specified Trust, Board, Organ Donation Services Team, or nation.

This report is provided for information and to facilitate case based discussion about organ donation by the Organ Donation Committees and Trusts/Boards.

As part of the PDA, patients over 80 years of age and those who did not die on a critical care unit or emergency department are not audited nationally and are therefore excluded from the majority of this report. Data from neonatal intensive care units (ICU) have also been excluded from this report. In addition, some information may be outstanding due to late reporting and difficulties obtaining patient notes. Donations not captured by the PDA will still be included in the data supplied from the accompanying Referral Record or from the UKTR, as appropriate.



Appendix A.3 Table and Figure Description

Г

For the purposes of this report please note that Trust/Board is equivalent to team.

1 Donor outcomes	
Table 1.1	The number of actual donors, the resulting number of patients transplanted and the average number of organs donated per donor have been obtained from the UK Transplant Registry (UKTR) for your Trust/Board. Results have been displayed separately for donors after brain death (DBD) and donors after circulatory death (DCD).
Table 1.2	The number of organs transplanted by type from donors at your Trust/Board has been obtained from the UKTR. Further information can be obtained from your local Specialist Nurse – Organ Donation (SNOD), specifically regarding organs that were not transplanted. Results have been displayed separately for DBD and DCD.
Figure 1.1	The number of actual donors and the resulting number of patients transplanted obtained from the UKTR for your Trust/Board for the past 10 equivalent time periods are presented on a line chart.

2 Key rates in potential for orga	an donation
Figure 2.1	Key percentage measures of DBD and DCD potential donation activity for your Trust/Board are presented in a bar chart, using data from the Potential Donor Audit (PDA). The comparative UK rate, for the same time period, is illustrated by the pink line. The key rates labels are coloured using the gold, silver, bronze, amber, and red (GoSBAR) colour scheme to show the performance of your Trust/Board, relative to the UK rate, as reflected in the funnel plots (see description for Figure 4.1 below.
Figure 2.2	Trends in the key percentage measures of DBD and DCD potential donation activity for your Trust/Board are presented for the past five equivalent time periods, using data from the PDA.
Table 2.1	A summary of DBD, DCD and deceased donor data and key numbers have been obtained from the PDA. A UK comparison is also provided. Note that caution should be applied when interpreting percentages based on small numbers. Appendix A.1 gives a fuller explanation of terms used. The key rates are highlighted using the gold, silver, bronze, amber, and red (GoSBAR) colour scheme to show the performance of your Trust/Board, relative to the UK rate, as reflected in the funnel plots (see description for Figure 4.1 below).

3 Best quality of care in organ do	onation
Figure 3.1	A stacked bar chart displays the number of patients with suspected neurological death who were tested and the number who were not tested in your Trust/Board for the past five equivalent time periods.
Table 3.1	The reasons given for neurological death tests not being performed in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.
Figure 3.2	Stacked bar charts display the number of DBD and DCD patients meeting referral criteria who were referred to the Organ Donation Service and the number who were not referred in your Trust/Board for the past five equivalent time periods.
Table 3.2	The reasons given for not referring patients to the Organ Donation Service in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.
Table 3.3	The primary absolute medical contraindications to solid organ donation for DBD and DCD patients have been obtained from the PDA, if applicable. A UK comparison is also provided.
Figure 3.3	Stacked bar charts display the number of families of DBD and DCD patients approached where a SNOD was present and the number approached where a SNOD was not present in your Trust/Board for the past five equivalent time periods.



Figure 3.4	Stacked bar charts display the number of families of DBD and DCD patients approached where consent/authorisation for organ donation was ascertained and the number approached where consent/authorisation was not ascertained in your Trust/Board for the past five equivalent time periods.
Table 3.4	The reasons why consent/authorisation was not ascertained for solid organ donation in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.
Table 3.5	The reasons why solid organ donation did not occur in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.

4 Comparative data	
Figure 4.1	A funnel plot of the neurological death testing rate is displayed using data obtained from the PDA. Each Trust/Board, of the same level, is represented on the plot as a blue dot, although one dot may represent more than one Trust/Board. The UK rate is shown on the plot as a green horizontal dashed line, together with 95% and 99.8% confidence limits for this rate. These limits form a 'funnel', which is shaded using the gold, silver, bronze, amber, and red (GoSBAR) colour scheme. Graphs obtained in this way are known as funnel plots. If a Trust/Board lies within the 95% limits, shaded bronze, then that Trust/Board has a rate that is statistically consistent with the UK rate (average performance). If a Trust/Board lies outside the 95% confidence limits, shaded silver (good performance) or amber (below average performance), this serves as an alert that the Trust/Board may have a rate that is significantly different from the UK rate. When a Trust/Board lies above the upper 99.8% limit, shaded gold, this indicates a rate that is significantly higher than the UK rate (exceptional performance), while a Trust/Board that lies below the lower limit, shaded red, has a rate that is significantly lower than the UK rate (poor performance). It is important to note that differences in patient mix have not been accounted for in these plots. Your Trust/Board is shown on the plot as the large black cross. If there is no large black cross on the plot, your Trust/Board did not report any patients of the type presented. The funnel plots can also be used to identify the maximum rates currently being achieved by Trusts/Boards with similar donor potential.
Figure 4.2	A funnel plot of the deceased donor referral rate is displayed using data obtained from the PDA. See description for Figure 4.1 above.
Figure 4.3	A funnel plot of the deceased donor SNOD presence rate is displayed using data obtained from the PDA. See description for Figure 4.1 above.
Figure 4.4	A funnel plot of the deceased donor consent/authorisation rate is displayed using data obtained from the PDA. See description for Figure 4.1 above.

5 PDA data by hospital and unit	
Table 5.1	DBD key numbers and rates by unit where the patient died have been obtained from the PDA. Percentages have been excluded where numbers are less than 10.
Table 5.2	DCD key numbers and rates by unit where the patient died have been obtained from the PDA. Percentages have been excluded where numbers are less than 10.



6 Paediatric ICU data	
Table 6.1	A summary of DBD, DCD and deceased donor data and key numbers for paediatric ICUs have been obtained from the PDA. A UK comparison is also provided. Note that caution should be applied when interpreting percentages based on small numbers. Appendix A.1 gives a fuller explanation of terms used.
Figure 6.1	A stacked bar chart displays the number of paediatric ICU patients with suspected neurological death who were tested and the number who were not tested in your Trust/Board for the past five equivalent time periods.
Table 6.2	The reasons given for neurological death tests not being performed for paediatric ICU patients in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.
Figure 6.2	Stacked bar charts display the number of DBD and DCD paediatric ICU patients meeting referral criteria who were referred to the Organ Donation Service and the number who were not referred in your Trust/Board for the past five equivalent time periods.
Table 6.3	The reasons given for not referring paediatric ICU patients to the Organ Donation Service in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.
Table 6.4	The primary absolute medical contraindications to solid organ donation for DBD and DCD paediatric ICU patients have been obtained from the PDA, if applicable. A UK comparison is also provided.
Figure 6.3	Stacked bar charts display the number of families of DBD and DCD paediatric ICU patients approached where a SNOD was present and the number approached where a SNOD was not present in your Trust/Board for the past five equivalent time periods.
Figure 6.4	Stacked bar charts display the number of families of DBD and DCD paediatric ICU patients approached where consent/authorisation for organ donation was ascertained and the number approached where consent/authorisation was not ascertained in your Trust/Board for the past five equivalent time periods.
Table 6.5	The reasons why consent/authorisation was not ascertained for solid organ donation in paediatric ICU patients in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.
Table 6.6	The reasons why solid organ donation did not occur in paediatric ICU patients in your Trust/Board, have been obtained from the PDA, if applicable. A UK comparison is also provided.

Γ

Г

7 Emergency department data	
Figure 7.1	Stacked bar charts display the number of patients that died in the emergency department (ED) who met the referral criteria and were referred to the Organ Donation Service and the number who were not referred in your Trust/Board for the past five equivalent time periods.
Figure 7.2	Stacked bar charts display the number of families of patients in ED approached where a SNOD was present and the number approached where a SNOD was not present in your Trust/Board for the past five equivalent time periods.

8 Additional data and figures	
Table 8.1	A summary of deceased donor, transplant, transplant list and ODR opt-in registration data for your region have been obtained from the UKTR. Your region has been defined as per former Strategic Health Authority. A UK comparison is also provided.
Table 8.2	Trust/board level categories and the relevant expected number of proceeding donors per year are provided for information.
Table 8.3	National DBD key numbers and rates for level 1, 2, 3 and 4 Trusts/Boards are displayed alongside your local data to aid comparison with equivalent Trusts/Boards. Percentages have been excluded where numbers are less than 10.



Table 8.4	National DCD key numbers and rates for level 1, 2, 3 and 4 Trusts/Boards are displayed alongside your local data to aid comparison with equivalent Trusts/Boards. Percentages have been excluded where numbers are less than 10.
Figure 8.1	A funnel plot of the DBD and DCD referral rates are displayed using data obtained from the PDA. See description for Figure 4.1 above.
Figure 8.2	A funnel plot of the DBD and DCD SNOD presence rates are displayed using data obtained from the PDA. See description for Figure 4.1 above.
Figure 8.3	A funnel plot of the DBD and DCD consent/authorisation rates are displayed using data obtained from the PDA. See description for Figure 4.1 above.