




ORIGINAL ARTICLE

Perioperative management of adult cadaveric and live donor renal transplantation in the UK: a survey of national practice

Clare M. Morkane ¹, Jez Fabes¹, Neal R. Banga², Peter D. Berry³ and Christopher J. Kirwan⁴

¹Division of Surgery and Interventional Science (University College London) & Royal Free Perioperative Research Group, Department of Anaesthesia, Royal Free Hospital, London, UK, ²Department of Renal Transplant Surgery, Royal Free Hospital, London, UK, ³Department of Anaesthesia, Royal Free Hospital, London, UK and ⁴Adult Critical Care Unit, Royal London Hospital, London, UK

Correspondence and offprint requests to: Christopher J. Kirwan; E-mail: christopher.kirwan@bartshealth.nhs.uk;

ABSTRACT

Background. There is a limited evidence base and no national consensus regarding the perioperative management of patients undergoing renal transplantation. We developed an electronic survey to capture an overview of renal transplant perioperative practice across UK renal transplant centres and determine the need for future guidelines on patient management.

Methods. A 29-question survey was developed to encompass the entire renal transplant perioperative pathway and input was sought from clinicians with expertise in renal transplant surgery, anaesthesia, nephrology and intensive care. The survey was sent to lead renal anaesthetists at each of the 23 transplant centres across the UK.

Results. A 96% response rate was achieved with 22 out of 23 centres returning complete responses. There was limited evidence of guideline-based approaches to preoperative workup. Questions regarding intraoperative fluid management, blood pressure targets, vasopressor administration and central venous pressure (CVP) monitoring identified a broad range of practice. Of note, the routine use of goal-directed fluid therapy based on cardiac output estimation was reported in six (27.3%) centres, while nine centres (40.9%) continue to target a specific CVP intraoperatively. In all, 12 (54.5%) centres perform transversus abdominis plane blocks with fentanyl-based patient-controlled analgesia as the most common mode of postoperative analgesia. A single centre reported a renal transplant-specific Enhanced Recovery after Surgery programme for cadaveric organ recipients.

Conclusions. This questionnaire highlighted a high degree of heterogeneity in current UK practice as regards the perioperative management of renal transplant recipients. Development of evidence-based national consensus guidelines to standardize the perioperative care of these patients is recommended in order to improve patient outcomes and focus areas of future research.

Keywords: anaesthesia, perioperative care, renal transplant, survey

Received: 31.10.2018; Editorial decision: 15.1.2019

© The Author(s) 2019. Published by Oxford University Press on behalf of ERA-EDTA.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

INTRODUCTION

Patient outcomes after major surgery can be improved considerably through the delivery of effective perioperative care [1]. Patients undergoing renal transplantation are increasing in age and multi-morbidity [2], with a unique pathophysiology that mandates effective evidence-based perioperative practice. Although significant research effort has been put into improving the management of patients undergoing major surgery, the majority of these studies exclude renal transplant recipients [3, 4]. This has resulted in a limited evidence base to inform the perioperative care of this complex patient cohort.

Currently, there is little national consensus on the perioperative management of patients undergoing renal transplantation and efforts to address this are hampered by the lack of a national body or representation to engender communication between transplant centres. To provide an evidence base to highlight the importance of this consensus, we developed a survey to generate an overview of current national practice across UK renal transplant units and identify areas of heterogeneity in management.

MATERIALS AND METHODS

An electronic survey was sent to the lead renal anaesthetist at each of the 23 centres performing adult cadaveric renal transplantation across the UK, between February 2018 and April 2018. Where this individual could not be identified, an anaesthetist with a renal transplant interest was sought.

The survey was developed to encompass the entire perioperative episode of patients undergoing renal transplantation. Input regarding content was sought from clinicians with expertise across this clinical spectrum, including transplant surgery, anaesthesia, nephrology and intensive care medicine. A total of 29 questions were included, divided into pre-, intra- and postoperative management. Closed questions designed to generate unambiguous responses were developed with free-text options to expand on answers given.

Information was requested regarding:

- (i) the presence of pre-assessment guidelines and the use of cardiac risk assessment testing;
- (ii) preoperative dialysis;
- (iii) intraoperative fluid management;
- (iv) intraoperative monitoring and haemodynamic targets;
- (v) vasoactive drugs;
- (vi) management of reperfusion; and
- (vii) analgesia.

All 29 questions are presented in [Appendix 1](#).

The survey was assessed via the National Research Ethics Service decision-making tool and did not meet current UK definitions for research requiring ethics committee approval [5]. The survey was registered with the local audit and clinical governance department.

RESULTS

Complete survey responses were obtained from 22 out of 23 adult renal transplant centres (96% response rate). All but one centre identified a site-specific guideline for the perioperative management of renal transplantation; these guidelines were predominately under the auspices of anaesthesia and nephrology.

Preoperative workup

Despite the presence of site-specific perioperative guidelines for the management of these patients there was significant heterogeneity in the indications for, and modality of, preoperative cardiorespiratory function testing performed. Of the 22 centres, 5 (22.7%) do not routinely perform cardiorespiratory investigations and where these are implemented they are performed on a case-by-case basis without defined indications or criteria. In the remaining centres, the selected testing modality showed broad variation based on local expertise and availability. The most commonly requested investigations include transthoracic echocardiography (TTE), cardiopulmonary exercise testing, pulmonary function and stress echo ([Table 1](#)). A single centre (4.5%) routinely requests cardiology review prior to transplantation, whereas the remaining centres identified common indications for cardiology referral as age >50 years and the presence of pre-existing diabetic and cardiovascular disease.

Perioperative management

Dialysis. The majority of centres (63.6%) dialyse preoperatively to the recipient's 'dry weight'. Three centres (13.6%) stated that they target a post-dialysis body weight 1–2 kg above the recipient's dry weight, whereas the others do not specify a preoperative dialysis target weight.

Grade of anaesthetist. The grade of the most senior anaesthetist managing a cadaveric renal transplant showed variation between centres and according to the time of day. During normal working hours, 81.8% of centres provide consultant-led care. Between 12 a.m. and 8 a.m. the most senior anaesthetist managing these cases was a post-fellowship registrar in 16 (72.7%) centres. In two (9.1%) centres, out-of-hours care was provided by a pre-fellowship registrar, with the four (18.2%) remaining centres providing consultant-led care at all times of the day.

Fluid management and cardiac output optimization. Survey responses identified a wide range of, and variability in, practice as regards intraoperative fluid administration. While most centres routinely use a balanced salt solution (plasmalyte, compound sodium lactate and the haemodialysis fluid 'hemosol'), eight (36.4%) centres reported the use of 0.9% saline as their primary intravenous fluid ([Figure 1](#)). One centre (4.5%) uses hypertonic 1.26% sodium bicarbonate as intraoperative maintenance. Fluid administration via a fixed-rate infusion pump was only used in one centre with all others opting for free-flow systems.

Goal-directed fluid therapy based on cardiac output (CO) estimation was described as routine practice in six (27.3%) centres. The remaining centres all used CO systems to guide fluid

Table 1. Responses to the question: which tests do you routinely request to assess preoperative cardiopulmonary function in patients to be listed for renal transplantation (select all that apply)?

Investigation	Number of centres (%)
TTE	10 (45.5)
Stress echo	4 (18.2)
Cardiopulmonary exercise test	2 (9.1)
Pulmonary function tests	2 (9.1)
Myocardial perfusion scan	1 (4.5)
6-min walk test	0 (0)
Shuttle test	0 (0)
Coronary angiogram	0 (0)




FIGURE 1: Responses to the question: which intravenous fluid is predominantly used intraoperatively during renal transplantation in your centre?

Table 2. Responses to the free-text question: if a specific MAP or SBP is targeted, please give details

Free-text responses

Patient dependent
 Above 100–110 mmHg systolic post-reperfusion
 Patient's baseline MAP
 MAP >80 mmHg
 Dependent on patient's preoperative blood pressure and surgical communication
 10% variability around the baseline MAP
 SBP >110 mmHg
 SBP >120 mmHg 20 min prior to reperfusion and onwards
 Within 10% of preoperative blood pressure
 Patient's usual SBP

administration on a patient-specific basis though no centres identified defined criteria for patient selection. The modalities of CO estimation employed were predominately arterial pressure wave analysis, such as LiDCOrapid and Flotrac (11 centres, 50%), and oesophageal Doppler (9 centres, 40.9%), with a single centre implementing non-invasive Finapres technology.

Central venous catheters. Three-quarters of centres regularly insert a central venous catheter (CVC) into renal transplant recipients and half of all centres would place a CVC in addition to an existing patent dialysis line. In all, nine centres (40.9%) target a specific central venous pressure (CVP) intraoperatively; the target CVP showed wide variation, with reported values from 7 to 17 cmH₂O.

Blood pressure management. A total of 13 centres (59.1%) target either a specific mean arterial pressure (MAP) or systolic blood pressure (SBP), but there was a wide variety in the targets described (Table 2). Quoted MAP targets varied between 10% and 20% of baseline recipient MAP, with two centres (9.1%) stating an MAP target >80 mmHg for all cases. A total of three centres (13.6%) target the recipient's preoperative SBP, aiming to maintain this throughout the case.

Two (9.1%) centres reported the routine insertion of arterial lines, with the remainder only doing so where arterial waveform methods of CO estimation were employed, as above. The vasoactive agents employed intraoperatively are predominately ephedrine and metaraminol in 11 and 15 centres, respectively (Figure 2). Phenylephrine and noradrenaline are reserved as secondary agents. One centre (4.5%) reported the frequent use of dopamine infusions.




FIGURE 2: Responses to the question: regarding vasoactive drugs, does your centre use the following frequently, rarely or never: dopamine, ephedrine, metaraminol, phenylephrine and noradrenaline.

Reperfusion. Forced diuresis with furosemide was reported as routine practice in 3 (13.6%) centres, whereas 11 centres (50% of responses) stated an institutional avoidance of its use for this indication. The administration of mannitol was more common, with routine use reported in nine (40.9%) centres, whereas only five centres (22.7%) stated that it was never used.

Analgesia. Transversus abdominis plane blocks are used routinely in 12 (54.5%) centres and of these two-thirds are sited by the anaesthetist (the remainder under direct vision by the operating surgeon). In all, two centres (9.1%) used a local anaesthetic infusion via a surgically inserted catheter between wound layers. Twenty (90.9%) centres provided patient-controlled analgesia postoperatively, with the majority of these (13 centres, 65%) using fentanyl as the first choice opioid, 5 (25%) using morphine and 2 (10%) oxycodone.

Postoperative care

A dedicated renal ward was the most common postoperative destination for renal transplant recipients (13 centres, 59.1%), while a renal- or transplant-specific high dependency unit provided postoperative care in the remaining centres. Several sites stipulated that high-risk recipients would be cared for in a general intensive care unit setting, decided on a case-by-case basis, although no predetermined criteria were used for patient selection.

A renal transplant-specific Enhanced Recovery after Surgery (ERAS) programme was in place in only one centre (4.5%) for cadaveric organ recipients and four centres (18.2%) for live donor recipients.

Practice according to transplant centre surgical volume

There was no evidence of standardization in practice between the nine largest volume transplant centres, which each performed >150 renal transplants in 2017/18 [6]. For example, regarding preoperative assessment, four of these nine centres (44.4%) reported a locally developed approach to preoperative investigation with no routine tests. Furthermore, one (11.1%) large-volume centre reported the routine use of CO estimation to guide fluid administration intraoperatively (oesophageal Doppler), while there was marked variation in implementing forced diuresis at reperfusion and vasopressor use. Of the four UK centres reporting the existence of an ERAS protocol for live-donor recipients, two were large-volume centres.

DISCUSSION

There is a clear heterogeneity in current UK practice regarding the management of renal transplant recipients throughout the entire perioperative period. Of note, this variation in practice is independent of transplant centre surgical volume, where it would be reasonable to expect defined protocols produced by experienced clinicians. Although the majority of centres have developed guidelines for the perioperative management of cadaveric renal transplantation, these are centre specific and disparate in their content.

There are some components of current practice identified in the survey for which there is evidence of an association with harm. The routine administration of chloride-rich solutions has been associated with impaired diuresis, decreased renal blood flow and development of metabolic acidosis compared with buffered salt solutions [7–9]. Hyperchloraemic acidosis is associated with acute kidney injury (AKI) following abdominal surgery, which may be attenuated by reducing the intraoperative chloride load [10]. It follows that the perioperative use of physiologically balanced replacement fluids is associated with a reduction in the need for dialysis and a reduced incidence of hyperkalaemia when compared with 0.9% saline in renal transplant surgery [11].

While renal transplant surgery generally involves much smaller fluid losses than major intra-peritoneal surgery [12], transplant recipients have an increased risk of fluid overload, especially in the anuric cohort. Despite this it remains common practice to give significant crystalloid volume expansion prior to renal vessel clamping with the aim of decreasing postoperative acute tubular necrosis. However, no published evidence supports the pursuit of positive fluid balance to improve renal outcomes [13], with retrospective data showing perioperative fluid administration of >2500 mL as strongly and independently associated with chronic allograft dysfunction [14]. This indiscriminate fluid administration approach is facilitated by ongoing debate regarding the utility of CO-based goal-directed fluid administration and use of approaches such as CVP targeting [15, 16].

Management of renal graft reperfusion is highly heterogeneous and not in keeping with the current evidence. There is no evidence in the general or post-renal transplantation population that forced diuresis with furosemide shortens AKI duration, reduces the need for dialysis or improves outcomes [17, 18]. The evidence base concerning the use of furosemide and mannitol is outdated, retrospective or contains a small number of patients, and thus their relevance to current practice is unclear [19, 20]. Similarly, the routine administration of mannitol must be weighed against the associated risks with accompanying hydration if the patient is anuric.

There is conflicting evidence regarding the effects of dopamine in the perioperative period. The majority of studies fail to show any significant benefit when administered to renal transplant recipients [21, 22]. Although recent evidence suggests that low-dose dopamine may increase renal plasma flow, urine flow rate and creatinine clearance [23], these findings have not been reproduced. Dopamine administered to live donor recipients incurs no outcome advantage in either recipient or donor [24].

We have shown that current out-of-hours care of these patients is often provided by more junior clinicians. Hence, a particular focus on trainee education, competence and confidence is required. Care of renal transplant recipients is poorly represented in the current anaesthetic curriculum and this requires redress.

Despite a growing evidence base within some specific domains of perioperative management of renal transplant recipients, controversy and variation in clinical practice persist. These debates are not unique to UK practice; indeed controversy over aspects of perioperative care for the renal transplant recipient is also evident in continental Europe, with several areas of management still under discussion and investigation [25, 26].

Though more up-to-date and robust research is required, significant improvements can be made to patient outcomes through better use and implementation of the evidence base already available through systematic review. Additional national-level consensus is also required and should be developed through the use of collaborative tools such as Delphi consensus methodology [27]. This should seek to generate guidelines covering the entire perioperative pathway of renal transplant recipients. Non-standardized, outdated practice has the potential to impact negatively on patient outcomes; the development of an ERAS programme is likely to resolve some of the uncontrolled variation in current practice and would be expected to enhance outcomes. The feasibility of an ERAS programme for deceased donor renal transplant recipients has been demonstrated [28], and there is emerging evidence pointing towards improved postoperative outcomes [29] alongside a wealth of existing literature to support the approach in other surgical specialties [30].

CONCLUSION

This survey has demonstrated a high degree of heterogeneity in current UK practice as regards the perioperative management of renal transplant recipients. It has also highlighted a number of non-standardized and potentially outdated practices. The formation of a working party and development of evidence-based national consensus guidelines are recommended to focus future research targets and standardize the perioperative care of these complex patients.

ACKNOWLEDGEMENTS

The authors would like to thank all the clinicians who took time to respond to this questionnaire.

CONFLICT OF INTEREST STATEMENT

None declared.

REFERENCES

1. Pearse RM, Ackland GL. Perioperative fluid therapy. *Br Med J* 2012; 344: e2865
2. Jassal SV, Schaubel DE, Fenton SS. Baseline comorbidity in kidney transplant recipients: a comparison of comorbidity indices. *Am J Kidney Dis* 2005; 46: 136–142
3. Pearse RM, Moreno RP, Bauer P et al. Mortality after surgery in Europe: a 7 day cohort study. *Lancet* 2012; 380: 1059–1065
4. Findlay G, Goodwin A, Protopappa K et al. *Knowing the Risk: A Review of the Perioperative Care of Surgical Patients*. London: National Confidential Enquiry into Patient Outcome and Death, 2011
5. Medical Research Council and NHS Health Research Authority. <http://www.hra-decisiontools.org.uk/research/index.html> (2 February 2019, date last accessed)
6. NHS Blood and Transplant and NHS England. *Annual Report on Kidney Transplantation. Report for 2017/2018, 2018*; <https://>

- nhsbtdeblob.core.windows.net/umbraco-assets-corp/12256/nhsbt-kidney-transplantation-annual-report-2017-2018.pdf (2 February 2019, date last accessed)
7. O'Malley CM, Frumento RJ, Hardy MA. A randomized, double-blind comparison of lactated Ringer's solution and 0.9% NaCl during renal transplantation. *Anesth Analg* 2005; 100: 1518–1524
 8. Chowdhury AH, Cox EF, Francis ST et al. A randomized, controlled, double-blind crossover study on the effects of 2-L infusions of 0.9% saline and plasma-lyte(R) 148 on renal blood flow velocity and renal cortical tissue perfusion in healthy volunteers. *Ann Surg* 2012; 256: 18–24
 9. Reid F, Lobo DN, Williams RN et al. (Ab)normal saline and physiological Hartmann's solution: a randomized double-blind crossover study. *Clin Sci* 2003; 104: 17–24
 10. Toyonaga Y, Kikura M. Hyperchloremic acidosis is associated with acute kidney injury after abdominal surgery. *Nephrology* 2017; 22: 720–727
 11. Adwaney A, Randall DW, Blunden MJ et al. Perioperative plasma-lyte use reduces the incidence of renal replacement therapy and hyperkalaemia following renal transplantation when compared with 0.9% saline: a retrospective cohort study. *Clin Kidney J* 2017; 10: 838–844
 12. Othman MM, Ismael AZ, Hammouda GE. The impact of timing of maximal crystalloid hydration on early graft function during kidney transplantation. *Anesth Analg* 2010; 110: 1440–1446
 13. Prowle JR, Echeverri JE, Ligabo EV et al. Fluid balance and acute kidney injury. *Nat Rev Nephrol* 2010; 6: 107–115
 14. Campos L, Parada B, Furriel F et al. Do intraoperative hemodynamic factors of the recipient influence renal graft function. *Transplant Proc* 2012; 44: 1800–1803
 15. De Gasperi A, Narcisi S, Mazza E et al. Perioperative fluid management in kidney transplantation: is volume overload still mandatory for graft function. *Transplant Proc* 2006; 38: 807–809
 16. Ferris RL, Kittur DS, Wilasrusmee C et al. Early hemodynamic changes after renal transplantation: determinants of low central venous pressure in the recipients and correlation with acute renal dysfunction. *Med Sci Monit* 2003; 9: CR61–CR66
 17. Ho KM, Sheridan DJ. Meta-analysis of frusemide to prevent or treat acute renal failure. *Br Med J* 2006; 333: 420
 18. Lachance SL, Barry JM. Effect of furosemide on dialysis requirement following cadaveric kidney transplantation. *J Urol* 1985; 133: 950–951
 19. van Valenberg PL, Hoitsma AJ, Tiggeler RG et al. Mannitol as an indispensable constituent of an intraoperative hydration protocol for the prevention of acute renal failure after renal cadaveric transplantation. *Transplantation* 1987; 44: 784–788
 20. Lauzurica R, Teixidó J, Serra A et al. Hydration and mannitol reduce the need for dialysis in cadaveric kidney transplant recipients treated with CyA. *Transplant Proc* 1992; 24: 46–47
 21. Grundmann R, Kindler J, Meider G et al. Dopamine treatment of human cadaver kidney graft recipients: a prospectively randomized trial. *Klin Wochenschr* 1982; 60: 193–197
 22. Sandberg J, Tyden G, Groth CG. Low-dose dopamine infusion following cadaveric renal transplantation: no effect on the incidence of ATN. *Transplant Proc* 1992; 24: 357
 23. Dalton RSJ, Webber JN, Cameron C et al. Physiologic impact of low-dose dopamine on renal function in the early post renal transplant period. *Transplantation* 2005; 79: 1561–1567
 24. O'Dair J, Evans L, Rigg KM et al. Routine use of renal-dose dopamine during living donor nephrectomy has no beneficial effect to either donor or recipient. *Transplant Proc* 2005; 37: 637–639
 25. Schmid S, Jungwirth B. Anaesthesia for renal transplant surgery: an update. *Eur J Anaesthesiol* 2012; 29: 552–558
 26. Breda A, Budde K, Figueiredo A et al. *European Association of Urology. EAU Guidelines on Renal Transplantation, 2017*; <https://uroweb.org/wp-content/uploads/Renal-Transplantation-2017-pocket.pdf> (2 February 2019, date last accessed)
 27. Dalkey NC. The Delphi method: an experimental study of group opinion. In: Dalkey NC, Rourke DL, Lewis R et al. (eds). *Studies in the Quality of Life: Delphi and Decision-making*. 1972. Lexington, MA: Lexington Books
 28. Kruszyna T, Niekowal B, Kraśnicka M et al. Enhanced recovery after kidney transplantation surgery. *Transplant Proc* 2016; 48: 1461–1465
 29. Espino KA, Narvaez JRF, Ott MC et al. Benefits of multimodal enhanced recovery pathway in patients undergoing kidney transplantation. *Clin Transplant* 2018; 32: e13173
 30. Lau CS, Chamberlain RS. Enhanced recovery after surgery programs improve patient outcomes and recovery: a meta-analysis. *World J Surg* 2017; 41: 899–913

Appendix 1

Table A1. Complete set of questions sent electronically to each transplant centre

Question number	Question details
1	Transplant unit (select option)
2	What is your job title?
3	Does your centre have a specific guideline for the perioperative management of renal transplantation? <ul style="list-style-type: none"> • Yes • No • If yes, which speciality(ies) is responsible for it? • Unsure/do not know
4	Which tests do you routinely request to assess preoperative cardiopulmonary function in patients to be listed for renal transplantation (select all that apply)? <ul style="list-style-type: none"> • Transthoracic echo (TTE) • Pulmonary function tests (PFTs) • Stress echo • Myocardial perfusion scan (MPS) • Cardiopulmonary exercise test • Coronary angiogram • 6-min walk • Shuttle test • Other (please specify)
5	Are patients routinely assessed by a cardiologist prior to transplantation? <ul style="list-style-type: none"> • Always • If previous history of cardiovascular disease • If patient >50 years old • Not routinely • Other (please specify)
6	For patients requiring haemodialysis prior to transplantation, does your centre aim to dialyse to dry weight pre-op? <ul style="list-style-type: none"> • Yes • No • If no, please give detail (i.e. 1 kg over dry weight)
7	Generally, what is the level of the most senior anaesthetist managing a cadaveric renal transplant in the following situations: <ul style="list-style-type: none"> • Normal working hours Monday–Friday (8 a.m.–6 p.m.) • Between the hours of 6 p.m. and 12 a.m. • Between the hours of 12 a.m. to 8 a.m.
8	Which intravenous fluid is predominantly used intraoperatively in your centre? <ul style="list-style-type: none"> • 0.9% saline • Plasmalyte • Compound sodium lactate • Hemosol (solution for haemodialysis) • Other (please specify)
9	What do you anticipate the average operative blood loss to be during renal transplant? <ul style="list-style-type: none"> • 0–250 mL • 250–500 mL • 500–750 mL • >750 mL
10	How do you manage intraoperative fluid administration: <ul style="list-style-type: none"> • Fixed continuous infusion with infusion pump • Continuous infusion without infusion pump • Boluses on top of infusion • Boluses only
11	Does your centre regularly use cardiac output (CO) monitoring to guide fluid management during renal transplantation? <ul style="list-style-type: none"> • Yes • No • Sometimes • Comments

(continued)