

Literature Review

Positive cultures of organ preservation fluid predict postoperative infections in solid organ transplantation recipients.

Yansouni CP; Dendukuri N; Liu G; Fernandez M; Frenette C; Paraskevas S; Sheppard DC.

Infection Control & Hospital Epidemiology. 33(7):672-80, 2012 Jul.

Bacterial contamination of organ graft preservation solution and infection after transplantation.

Sauget M; Verdy S; Slekovec C; Bertrand X; Talon D.

Transplant Infectious Disease. 13(4):331-4, 2011 Aug.

Perfusion fluid contamination in relation to recipient survival and acute cellular rejection in orthotopic liver transplantation: retrospective analysis.

Chaim FH; Boin IF; Ataide EC; Stucchi RS.

Transplantation Proceedings. 43(4):1313-5, 2011 May.

Impact of bacterial and fungal donor organ contamination in lung, heart-lung, heart and liver transplantation.

Mattner F; Kola A; Fischer S; Becker T; Haverich A; Simon A; Suerbaum S; Gastmeier P; Weissbrodt H; Struber M.

Infection. 36(3):207-12, 2008 Jun.

Outbreak of pseudomonas aeruginosa by multiple organ transplantation from a common donor.

Kumar D; Cattral MS; Robicsek A; Gaudreau C; Humar A.

Transplantation. 75(7):1053-5, 2003 Apr 15.

The use of contaminated donor organs in transplantation.

Zibari GB; Lipka J; Zizzi H; Abreo KD; Jacobbi L; McDonald JC.

Clinical Transplantation. 14(4 Pt 2):397-400, 2000 Aug.