



**Blood and Transplant**

# **ANNUAL REPORT ON INTESTINE TRANSPLANTATION**

**REPORT FOR 2025/2026  
(1 APRIL 2016 – 31 MARCH 2026)**

**PUBLISHED SEPTEMBER 2026**

**PRODUCED IN COLLABORATION WITH NHS ENGLAND**



# CONTENTS

## Contents

|                                                                       |           |
|-----------------------------------------------------------------------|-----------|
| <b>1. Executive summary</b>                                           | <b>4</b>  |
| <b>2. Introduction</b>                                                | <b>7</b>  |
| 2.1 Transplant list                                                   | 8         |
| 2.1.1 Post-registration outcomes, 1 April 2022 – 31 March 2024        | 11        |
| 2.1.2 Median waiting time to transplant, 1 April 2022 – 31 March 2025 | 12        |
| 2.2 Transplants                                                       | 13        |
| 2.3 Geographical variation in registration and transplant rates       | 15        |
| <b>ADULT INTESTINE TRANSPLANTATION</b>                                | <b>17</b> |
| 3. Transplant list                                                    | 18        |
| 3.1 Adult intestine transplant list as at 31 March, 2017 – 2026       | 18        |
| 4. Response to offers                                                 | 23        |
| 5. Transplants                                                        | 23        |
| 5.1 Adult intestine transplants, 1 April 2016 – 31 March 2026         | 23        |
| 5.2 Total preservation time, 1 April 2016 – 31 March 2026             | 27        |
| 6. Post-transplant survival                                           | 29        |
| <b>PAEDIATRIC INTESTINE TRANSPLANTATION</b>                           | <b>32</b> |
| 7. Transplant list                                                    | 33        |
| 7.1 Paediatric intestine transplant list as at 31 March, 2017 – 2026  | 33        |
| 8. Response to offers                                                 | 37        |
| 9. Transplants                                                        | 37        |
| 9.1 Paediatric intestine transplants, 1 April 2016 – 31 March 2026    | 37        |
| 9.2 Total preservation time, 1 April 2016 – 31 March 2026             | 41        |
| 10. Post-transplant survival                                          | 42        |
| 10.1 Survival by transplant type                                      | 42        |
| <b>11. Form return rates</b>                                          | <b>44</b> |
| 11.1 Form return rates, 1 January 2025 – 31 December 2025             | 45        |
| <b>APPENDIX</b>                                                       | <b>45</b> |
| A1: Number of patients analysed                                       | 46        |
| A2: Methods                                                           | 48        |
| A3: Glossary of terms                                                 | 50        |



# **EXECUTIVE SUMMARY**

## **EXECUTIVE SUMMARY**

This report presents key figures about intestine transplantation in the UK. The period reported covers 10 years of transplant data, from 1 April 2016 to 31 March 2026. The report presents information on patients on the transplant list, number of transplants, demographic characteristics of donors and transplant recipients, and patient survival after first intestine transplant, both on a national and a centre-specific basis. The unadjusted results on post-transplant survival should be regarded as guidance only due to the limited number of transplants performed.

## Key findings

- On 31 March 2026, there were 14 patients on the UK active intestine **transplant list**, an increase of 6 since the end of the previous year.
- Of those patients registered onto the transplant list in a recent two year period (1 April 2022 – 31 March 2024), 71% had received a transplant two years post-registration, while 10% died on the list (or were removed due to condition deteriorating), 7% were removed and 12% were still waiting at time of data analysis.
- **Median total waiting time** to elective intestine transplant between 1 April 2022 and 31 March 2025 was 136 days for **adult registrations** and 316 days for **paediatric registrations**.
- There were 201 intestine **transplants** performed in the UK over the 10 year period covered by this report. Of all transplants, 76% were in adult recipients while 24% were in paediatric recipients. A small proportion of these were re-transplants while the rest were primary transplants.
- In the last financial year, 2025/2026, 17 intestine **transplants** were performed, the same as 2024/2025. Of the 17 transplants, 15 (88%) were in adult recipients and 2 (12%) were in paediatric recipients.
- The national rates of **survival** ([unadjusted](#)) after first intestine transplantation for **adult patients** were estimated at 98%, 88%, and 72% at 90 days, one and five years post-transplant, respectively, for those **who did not receive the liver** and 92%, 78%, and 56% at 90 days, one and five years post-transplant, respectively, for those **who did receive the liver**. There was no evidence to suggest a difference in survival between those receiving and those not receiving the liver at 90 days and one year post-transplant. However, there was evidence to suggest a difference in survival rates at five years post-transplant (log-rank p-value=0.03). Please note that these survival rates are unadjusted and include registrations due to malignancy.
- The national rates of **survival** after first intestine transplantation for **paediatric patients** ([unadjusted](#)) were estimated at 100%, 100% and 80% at 90 days, one and five years post-transplant, respectively, for those **who did not receive the liver** and 96%, 84% and 68% at 90 days, one and five years post-transplant, respectively, for those **who did receive the liver**.

Use of the contents of this report should be acknowledged as follows: *Annual Report on Intestine Transplantation 2025/2026, NHS Blood and Transplant.*

# INTRODUCTION



## INTRODUCTION


This report presents information on the UK transplant list, transplant activity and transplant outcomes between 1 April 2016 and 31 March 2026, for all four designated centres performing intestine transplantation in the UK. Data were obtained from the UK Transplant Registry, at NHS Blood and Transplant, which holds information relating to donors, recipients and outcomes for all intestine transplants performed in the UK. Data were extracted on 08 July 2026.

The report is divided into two main sections; one for adult recipients (aged  $\geq 18$  years) and one for paediatric recipients (aged  $< 18$  years). Cambridge and Oxford are *adult* transplant centres, whereas Birmingham and King's College Hospital are *paediatric* transplant centres. Any transplants carried out at Cambridge or Oxford in patients aged less than 18 are included in the adult section, and any transplants carried out at Birmingham or King's College Hospital in patients 18 or over are included in the paediatric section. Unadjusted [patient survival rates](#) are calculated for these two groups at 90 days, one year and five years post-transplantation; these should be regarded as guidance only due to the limited number of transplants performed.

### 2.1 Transplant list


**Figure 2.1** shows the total number of patients on the intestine [active transplant list](#) at 31 March of each year between 2017 and 2026. The number of patients waiting for a transplant has decreased in recent years, with a peak of 18 patients in 2020 compared to 14 in 2026.

**Figure 2.1** Patients on the active intestine transplant list at 31 March




**Figure 2.2** shows the number of adult and paediatric patients on the [active transplant list](#) on 31 March 2026, by centre. In total, there were 10 adults and 4 paediatric patients on the active transplant list. Ten-year trends of the number of adult and paediatric patients on the active transplant list by transplant centre are shown later in the report, in **Figure 3.2** and **Figure 7.2**, respectively.

**Figure 2.2** Patients on the active intestine transplant list at31 March 2026, by centre




**Figure 2.3** shows the number of registrations made onto the active intestine transplant list each year in the last 10 years. The number of registrations has fluctuated over the decade, with 24 registrations in 2025/2026.

**Figure 2.3** Number of registrations onto the active intestine transplant list 1 April 2016 - 31 March 2026



**Figure 2.4** shows the number of registrations onto the active intestine transplant list in 2025/2026, by centre. There were 20 adult registrations and 4 paediatric registrations.


**Figure 2.4** Registrations onto the active intestine transplant list 1 April 2025 - 31 March 2026, by centre



2.1.1 Post-registration outcomes, 1 April 2022 – 31 March 2024

The registration outcomes of patients listed between 1 April 2022 and 31 March 2024 for an elective intestine transplant are summarised in **Figure 2.5**. This shows the proportion of patients transplanted or still waiting six months, one year and two years after joining the transplant list. At two years post-registration, 71% of patients had received a transplant and 12% were still waiting.

**Figure 2.5** Post-registration outcome for 58 elective registrations for an intestine transplant made in the UK, 1 April 2022 - 31 March 2024



\*Removals due to condition deteriorating

## 2.1.2 Median waiting time to transplant, 1 April 2022 – 31 March 2025

**Table 2.1** shows [median waiting time](#) to [elective](#) intestine transplant by registration type for adult and paediatric patients, combined, registered between 1 April 2022 and 31 March 2025. Patients who received a living donor transplant were excluded. A small proportion of patients requiring a liver containing graft are removed from the intestinal transplant waiting list to receive a liver transplant without the small bowel; the liver transplant is counted as receiving a transplant. Overall, patients waited a median total of 155 days (approximately six months) for a transplant.


| <b>Table 2.1      Median waiting time to elective intestine transplant in the UK,<br/>for patients registered 1 April 2022 - 31 March 2025</b> |                                              |                                              |                                    |                                   |                                    |
|------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------|----------------------------------------------|------------------------------------|-----------------------------------|------------------------------------|
| <b>Registration type</b>                                                                                                                       | <b>Number of<br/>patients<br/>registered</b> | <b>Total waiting time (days)<sup>1</sup></b> |                                    | <b>Active waiting time (days)</b> |                                    |
|                                                                                                                                                |                                              | <b>Median</b>                                | <b>95% Confidence<br/>interval</b> | <b>Median</b>                     | <b>95% Confidence<br/>interval</b> |
| Liver containing <sup>2</sup>                                                                                                                  | 30                                           | 250                                          | 41 – 459                           | 183                               | 25 – 341                           |
| Non liver containing                                                                                                                           | 46                                           | 144                                          | 86 – 202                           | 131                               | 72 – 190                           |
| <b>TOTAL</b>                                                                                                                                   | <b>76</b>                                    | <b>180</b>                                   | <b>126 – 194</b>                   | <b>144</b>                        | <b>105 - 183</b>                   |

<sup>1</sup> Periods of suspension from the list are included in the calculation of total median waiting times  
<sup>2</sup> Patients who received a liver only transplant are classed as transplanted

## 2.2 Transplants


**Figure 2.6** shows the number of intestine transplants performed each year in the last 10 years. Currently in the UK, intestine transplants are performed from donors after brain death ([DBD](#)), however, there have been two transplants, one in 2017/2018 and one in 2023/2024, performed using living donors. The total number of transplants over the decade was 201, with annual figures fluctuating between 12 and 27. In 2025/26, 17 transplants were performed, the same as the previous year.

**Figure 2.6** Total number of intestine transplants 1 April 2016 - 31 March 2026



**Figure 2.7** details the 201 intestine transplants performed in the UK in the 10 year period. Of these, 49 (24%) were in paediatric recipients and 152 (76%) were in adult recipients. Most of both paediatric and adult transplants were in first time recipients.

**Figure 2.7 UK intestine transplants, 1 April 2016 to 31 March 2026**



<sup>1</sup> Excludes one known abdominal wall only transplant (October 2016)

<sup>2</sup> Includes one living donor liver and bowel transplant (June 2017) and one living donor bowel-only transplant (September 2023)


\* Two of these transplants were super-urgent

## 2.3 Geographical variation in registration and transplant rates

**Figure 2.8** shows rates of registration to the intestinal transplant list per million population (pmp) between 1 April 2016 and 31 March 2026 compared with intestinal transplant rates pmp for the same time period, by recipient country/NHS region of residence. **Table 2.2** shows the breakdown of these numbers by recipient country/NHS region of residence. No adjustments have been made for potential demographic differences in populations. If a patient has had more than one registration/transplant in the period, each registration/transplant is considered. Note that this analysis only considered NHS Group 1 patients.

Since there will inevitably be some random variation in rates between areas, the systematic coefficient component of variation (SCV) was used to identify if the variation is more (or less) than a random effect for the different NHS regions in England only. Only first registrations and transplants in this period were considered. The larger the SCV the greater the evidence of a high level of systematic variation between areas. Registration and transplant rates yielded an SCV of 0 (p-value >0.999) and 0 (p-value >0.999), respectively. The p-value shows the probability that an SCV of this size (or higher) would be observed by chance if only random variation existed and therefore, no evidence of geographical variation beyond what would be expected at random. No adjustment has been made for area-specific demographic characteristics that may impact the rates of registration to the transplant list and transplantation such as age and sex. Therefore, these results should be interpreted with caution.

**Figure 2.8 Comparison of mean annual intestine registration rates (pmp) with mean annual transplant rates (pmp) by recipient country/NHS region of residence, 1 April 2016 – 31 March 2026**



**Table 2.2 Intestinal registration and transplant rates per million population (pmp) in the UK, 1 April 2016 - 31 March 2026, by Country/NHS region**

| <b>Country/<br/>NHS region</b> | <b>Registrations (pmp)</b> |               | <b>Transplants (pmp)</b> |               |
|--------------------------------|----------------------------|---------------|--------------------------|---------------|
| North East and Yorkshire       | 2.8                        | (0.33)        | 2.2                      | (0.26)        |
| North West                     | 2.5                        | (0.32)        | 2                        | (0.26)        |
| Midlands                       | 3.8                        | (0.34)        | 3.1                      | (0.28)        |
| East of England                | 3.8                        | (0.58)        | 2.7                      | (0.41)        |
| London                         | 2.4                        | (0.26)        | 2.1                      | (0.23)        |
| South East                     | 4.4                        | (0.46)        | 2.6                      | (0.27)        |
| South West                     | 1.7                        | (0.29)        | 0.9                      | (0.15)        |
| <b>England</b>                 | <b>21.4</b>                | <b>(0.36)</b> | <b>15.6</b>              | <b>(0.27)</b> |
| <b>Isle of Man</b>             | -                          |               | -                        |               |
| <b>Channel Islands</b>         | -                          |               | -                        |               |
| <b>Wales</b>                   | <b>0.7</b>                 | <b>(0.22)</b> | <b>0.3</b>               | <b>(0.09)</b> |
| <b>Scotland</b>                | <b>2.3</b>                 | <b>(0.41)</b> | <b>1.6</b>               | <b>(0.29)</b> |
| <b>Northern Ireland</b>        | <b>1.4</b>                 | <b>(0.71)</b> | <b>1</b>                 | <b>(0.51)</b> |
| <b>TOTAL</b>                   | <b>26<sup>1</sup></b>      | <b>(0.37)</b> | <b>18.8<sup>2</sup></b>  | <b>(0.27)</b> |

<sup>1</sup> Registrations include 2 recipients whose postcode was unknown and excludes 6 recipients who reside in the Republic of Ireland and 5 recipients who reside overseas

<sup>2</sup> Transplants include 3 recipients whose postcode was unknown and excludes 4 recipients who reside in the Republic of Ireland and 4 recipients who reside overseas

# **ADULT INTESTINE TRANSPLANTATION**




### 3. Transplant list

#### 3.1 Adult intestine transplant list as at 31 March, 2017 – 2026

**Figure 3.1** shows the number of adults active or suspended on the intestine transplant list on 31 March of each year between 2017 and 2026. The number of adults on the [active intestine transplant list](#) remained low over most of the decade with 10 adults active at 31 March 2026.


**Figure 3.1** Adult patients on the intestine transplant list at 31 March, 2017 - 2026



<sup>1</sup> Excludes one patient at Oxford registered for abdominal wall only

**Figure 3.2** shows the number of adults on the intestine transplant list at 31 March of each year between 2017 and 2026, at each transplant centre. Cambridge had generally more adults on the national [active transplant list](#) than Oxford.

**Figure 3.2** Adult patients on the intestine transplant list at 31 March, by centre, 2017 - 2026




The demographic characteristics of 203 adult intestine transplant recipient registrations in the 10-year period are shown by centre and overall in **Table 3.1**. Nationally, 51% of registrations were male and the [median](#) age was 45 years old. The most common reported indication for transplantation was short bowel syndrome (29%). The median recipient BMI was 22 kg/m<sup>2</sup>. For some characteristics, percentages may not add up to 100 due to rounding.

**Table 3.1 Demographic characteristics of adult intestine transplant recipient registrations, 1 April 2016 - 31 March 2026**

|                              |                                  | Cambridge<br>N (%) | Oxford<br>N (%) | TOTAL<br>N (%)    |
|------------------------------|----------------------------------|--------------------|-----------------|-------------------|
| Number of registrations      |                                  | 143                | 60              | <b>203 (100)</b>  |
| Number of patients           |                                  | 140 (100)          | 59 (100)        | <b>199 (100)</b>  |
| Registration type            | Elective                         | 139 (97)           | 60 (100)        | <b>199 (98)</b>   |
|                              | Super-urgent                     | 4 (3)              | 0               | <b>4 (2)</b>      |
| Transplant type required     | Non liver containing             | 69 (48)            | 60 (100)        | <b>129 (64)</b>   |
|                              | Liver containing                 | 74 (52)            | 0               | <b>74 (36)</b>    |
| Recipient sex                | Male                             | 71 (50)            | 33 (55)         | <b>104 (51)</b>   |
|                              | Female                           | 72 (50)            | 27 (45)         | <b>99 (49)</b>    |
| Recipient ethnicity group    | White                            | 126 (88)           | 55 (92)         | <b>181 (89)</b>   |
|                              | Other                            | 17 (12)            | 5 (8)           | <b>22 (11)</b>    |
| Indication group             | Short bowel syndrome             | 42 (30)            | 17 (28)         | <b>59 (29)</b>    |
|                              | Motility disorders               | 5 (4)              | 1 (2)           | <b>6 (3)</b>      |
|                              | Primary mucosal disorders        | 2 (1)              | 0               | <b>2 (1)</b>      |
|                              | Tumour                           | 19 (13)            | 30 (50)         | <b>49 (24)</b>    |
|                              | Liver disease                    | 11 (8)             | 0               | <b>11 (6)</b>     |
|                              | Retransplant                     | 11 (8)             | 4 (7)           | <b>15 (8)</b>     |
|                              | Mesenteric thrombosis (venous)   | 7 (5)              | 0               | <b>7 (4)</b>      |
|                              | Mesenteric thrombosis (arterial) | 3 (2)              | 3 (5)           | <b>6 (3)</b>      |
|                              | Other/not reported               | 41 (29)            | 5 (8)           | <b>46 (23)</b>    |
| Patient location             | Out-patient                      | 82 (57)            | 36 (60)         | <b>118 (58)</b>   |
|                              | Ward                             | 16 (11)            | 4 (7)           | <b>20 (10)</b>    |
|                              | ICU/HDU                          | 3 (2)              | 0               | <b>3 (2)</b>      |
|                              | Not reported                     | 42 (29)            | 20 (33)         | <b>62 (31)</b>    |
| Pre-transplant renal support | No                               | 134 (94)           | 58 (97)         | <b>192 (95)</b>   |
|                              | Yes                              | 8 (6)              | 2 (3)           | <b>10 (5)</b>     |
|                              | Not reported                     | 1 (1)              | 0               | <b>1 (1)</b>      |
| Previous abdominal surgery   | No                               | 13 (9)             | 0               | <b>13 (6)</b>     |
|                              | Yes                              | 125 (87)           | 60 (100)        | <b>185 (91)</b>   |
|                              | Not reported                     | 5 (3)              | 0               | <b>5 (3)</b>      |
| Recipient blood group        | O                                | 59 (41)            | 25 (42)         | <b>84 (41)</b>    |
|                              | A                                | 54 (38)            | 29 (48)         | <b>83 (41)</b>    |
|                              | B                                | 20 (14)            | 4 (7)           | <b>24 (12)</b>    |
|                              | AB                               | 10 (7)             | 2 (3)           | <b>12 (6)</b>     |
| Recipient age (years)        | Median (IQR)                     | 43 (33,53)         | 50 (40,56)      | <b>45 (34,54)</b> |
| Recipient BMI (kg/m2)        | Median (IQR)                     | 22 (20,26)         | 22 (21,25)      | <b>22 (20,26)</b> |


An indication of registration outcomes for adult elective patients registered on the intestine transplant list in the UK between 1 April 2022 and 31 March 2024 is summarised in **Figure 3.3**.

**Figure 3.3** Post-registration outcome for 39 elective adult registrations for an intestine transplant made in the UK, 1 April 2022 - 31 March 2024



**Figure 3.4** shows the proportion of patients transplanted, removed, died while waiting, or still waiting on the list at 6 months after joining the list, by transplant centre. 61% of registrations at Cambridge received a transplant within 6 months of listing compared to 73% at Oxford. Note there were 28 registrations at Cambridge, and 11 registrations at Oxford between 1 April 2022 and 31 March 2024.

**Figure 3.4** Six month post-registration outcome for 39 elective adult registrations for an intestine transplant made in the UK, 1 April 2022 - 31 March 2024



**Table 3.2** shows [median waiting time](#) to [elective](#) intestine transplant by registration type for adult patients, registered between 1 April 2022 and 31 March 2025. Registrations ending in a living donor transplant are excluded. The national UK median total waiting time to transplant for adult elective patients is 136 days.

| <b>Table 3.2      Median total waiting time to elective intestine transplant in the UK,<br/>for adults registered 1 April 2022 - 31 March 2025</b> |                                              |                                              |                                    |                                   |                                    |
|----------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------|----------------------------------------------|------------------------------------|-----------------------------------|------------------------------------|
| <b>Registration type</b>                                                                                                                           | <b>Number of<br/>patients<br/>registered</b> | <b>Total waiting time (days)<sup>1</sup></b> |                                    | <b>Active waiting time (days)</b> |                                    |
|                                                                                                                                                    |                                              | <b>Median</b>                                | <b>95% Confidence<br/>interval</b> | <b>Median</b>                     | <b>95% Confidence<br/>interval</b> |
| Liver containing                                                                                                                                   | 13                                           | 54                                           | 0 – 115                            | 36                                | 2 – 70                             |
| Non liver containing                                                                                                                               | 43                                           | 144                                          | 88 – 200                           | 131                               | 65 – 197                           |
| <b>Total</b>                                                                                                                                       | <b>56</b>                                    | <b>136</b>                                   | <b>69 – 203</b>                    | <b>112</b>                        | <b>44 – 180</b>                    |
| <sup>1</sup> Periods of suspension from the list are included in the calculation of total median waiting times                                     |                                              |                                              |                                    |                                   |                                    |

## 4. Response to offers

Potential [DBD](#) donors aged under 60 years and with a weight of less than 90 kg are considered for intestine donation. However, centres are highly selective when accepting donor organs which leads to high decline rates. Between 1 April 2025 and 31 March 2026, Cambridge received 190 intestine offers from 120 donors and Oxford received 37 intestine offers from 32 donors. Their [donor offer decline rates](#) were 94% and 89%, respectively.


| Table 4.1 Offer decline rates for each centre, 1 April 2025 - 31 March 2026 |            |           |            |           |
|-----------------------------------------------------------------------------|------------|-----------|------------|-----------|
| Centre                                                                      | Offers     | %Decline  | Donors     | %Decline  |
| Cambridge                                                                   | 190        | 94        | 120        | 91        |
| Oxford                                                                      | 37         | 89        | 32         | 88        |
| <b>Total</b>                                                                | <b>227</b> | <b>93</b> | <b>152</b> | <b>90</b> |

## 5. Transplants

### 5.1 Adult intestine transplants, 1 April 2016 – 31 March 2026

**Figure 5.1** shows the number of adult intestine transplants performed in the last 10 years, by [transplant type](#). The annual number of adult transplants has increased from 11 in 2024/2025 to 15 in 2025/2026.


**Figure 5.1 Adult intestine transplants in the UK, 1 April 2016 - 31 March 2026**



Transplants may also include any of stomach, spleen, abdominal wall, kidney, color


**Figure 5.2** shows the number of adult intestine transplants performed in 2025/2026, by centre and [transplant type](#). Oxford performed 4 transplants in the last financial year. Cambridge performed 11 transplants (3 liver containing and 8 non liver containing grafts).

**Figure 5.2 Adult intestine transplants in the UK by centre 1 April 2025 - 31 March 2026**



**Figure 5.3** shows the number of adult intestine transplants performed in the last 10 years, by centre and [type of transplant](#). Oxford performed their highest number of transplants over the decade in 2018/2019, while Cambridge performed their highest number in 2022/2023. Overall, Cambridge performed 111 transplants (73%) and Oxford performed 41 (27%).

**Figure 5.3 Adult intestine transplants by centre, 1 April 2016 - 31 March 2026**



The demographic characteristics of 152 adult intestine transplant recipients in the 10 year period are shown by centre and overall in **Table 5.1**. Nationally, 55% of recipients were male and the [median](#) age of recipients was 46 years old, while the median age of donors was 30 years old. The most common indication for transplantation was short bowel syndrome. Two of the transplants were in super-urgent patients, with the rest in [elective](#) recipients. For some characteristics, percentages may not add up to 100 due to rounding.

| <b>Table 5.1 Demographic characteristics of adult intestine transplant recipients, 1 April 2016 - 31 March 2026</b> |                                  |                    |                 |                  |
|---------------------------------------------------------------------------------------------------------------------|----------------------------------|--------------------|-----------------|------------------|
|                                                                                                                     |                                  | Cambridge<br>N (%) | Oxford<br>N (%) | TOTAL<br>N (%)   |
| <b>Number of transplants</b>                                                                                        |                                  | <b>111</b>         | <b>41</b>       | <b>152 (100)</b> |
| Transplant type                                                                                                     | Non liver containing             | 60 (54)            | 41 (100)        | <b>101 (66)</b>  |
|                                                                                                                     | Liver containing                 | 51 (46)            | 0               | <b>51 (34)</b>   |
| Urgency                                                                                                             | Elective                         | 109 (98)           | 41 (100)        | <b>150 (99)</b>  |
|                                                                                                                     | Super-urgent                     | 2 (2)              | 0               | <b>2 (1)</b>     |
| Recipient sex                                                                                                       | Male                             | 57 (51)            | 27 (66)         | <b>84 (55)</b>   |
|                                                                                                                     | Female                           | 54 (49)            | 14 (34)         | <b>68 (45)</b>   |
| Recipient ethnicity group                                                                                           | White                            | 99 (89)            | 37 (90)         | <b>136 (90)</b>  |
|                                                                                                                     | Other                            | 12 (11)            | 4 (10)          | <b>16 (11)</b>   |
| Indication group                                                                                                    | Short bowel syndrome             | 41 (37)            | 17 (41)         | <b>58 (38)</b>   |
|                                                                                                                     | Motility disorders               | 5 (5)              | 1 (2)           | <b>6 (4)</b>     |
|                                                                                                                     | Tumour                           | 21 (19)            | 15 (37)         | <b>36 (24)</b>   |
|                                                                                                                     | Liver disease                    | 9 (8)              | 0               | <b>9 (6)</b>     |
|                                                                                                                     | Retransplant                     | 10 (9)             | 5 (12)          | <b>15 (10)</b>   |
|                                                                                                                     | Mesenteric thrombosis (venous)   | 6 (5)              | 0               | <b>6 (4)</b>     |
|                                                                                                                     | Mesenteric thrombosis (arterial) | 4 (4)              | 2 (5)           | <b>6 (4)</b>     |
|                                                                                                                     | Other/not reported               | 15 (14)            | 1 (2)           | <b>16 (11)</b>   |
| Patient location                                                                                                    | Out-patient                      | 76 (68)            | 39 (95)         | <b>115 (76)</b>  |
|                                                                                                                     | Ward                             | 26 (23)            | 2 (5)           | <b>28 (18)</b>   |
|                                                                                                                     | ICU/HDU                          | 6 (5)              | 0               | <b>6 (4)</b>     |
|                                                                                                                     | Not reported                     | 3 (3)              | 0               | <b>3 (2)</b>     |
| Pre-transplant renal support                                                                                        | No                               | 97 (87)            | 39 (95)         | <b>136 (90)</b>  |
|                                                                                                                     | Yes                              | 10 (9)             | 2 (5)           | <b>12 (8)</b>    |
|                                                                                                                     | Not reported                     | 4 (4)              | 0               | <b>4 (3)</b>     |
| Previous abdominal surgery                                                                                          | No                               | 9 (8)              | 1 (2)           | <b>10 (7)</b>    |
|                                                                                                                     | Yes                              | 94 (85)            | 40 (98)         | <b>134 (88)</b>  |
|                                                                                                                     | Not reported                     | 8 (7)              | 0               | <b>8 (5)</b>     |
| Life style activity                                                                                                 | Normal                           | 11 (10)            | 0               | <b>11 (7)</b>    |
|                                                                                                                     | Restricted                       | 26 (23)            | 10 (24)         | <b>36 (24)</b>   |
|                                                                                                                     | Self-care                        | 38 (34)            | 28 (68)         | <b>66 (43)</b>   |
|                                                                                                                     | Confined                         | 8 (7)              | 1 (2)           | <b>9 (6)</b>     |
|                                                                                                                     | Reliant                          | 9 (8)              | 0               | <b>9 (6)</b>     |
|                                                                                                                     | Not reported                     | 19 (17)            | 2 (5)           | <b>21 (14)</b>   |


**Table 5.1 Demographic characteristics of adult intestine transplant recipients, 1 April 2016 - 31 March 2026**

|                                          |              | Cambridge<br>N (%) | Oxford<br>N (%) | TOTAL<br>N (%) |
|------------------------------------------|--------------|--------------------|-----------------|----------------|
| Restricted venous access at registration | No           | 64 (58)            | 27 (66)         | 91 (60)        |
|                                          | Yes          | 39 (35)            | 11 (27)         | 50 (33)        |
|                                          | Not reported | 8 (7)              | 3 (7)           | 11 (7)         |
| Recipient age (years)                    | Median (IQR) | 43 (34,54)         | 52 (41,57)      | 46 (34,55)     |
|                                          | Not reported | 0                  | 0               | 0              |
| Recipient BMI (kg/m2)                    | Median (IQR) | 23 (20,26)         | 22 (21,25)      | 23 (20,26)     |
|                                          | Not reported | 3                  | 3               | 6              |
| Serum bilirubin (umol/l)                 | Median (IQR) | 15 (8,32)          | 11 (6,17)       | 14 (7,29)      |
|                                          | Not reported | 4                  | 0               | 4              |
| Serum bilirubin (umol/l) non-liver       | Median (IQR) | 10 (6,17)          | 11 (6,17)       | 11 (6,17)      |
|                                          | Not reported | 3                  | 0               | 3              |
| Serum bilirubin (umol/l) including liver | Median (IQR) | 28 (14,175)        | 0               | 28 (14,175)    |
|                                          | Not reported | 1                  | 0               | 1              |
| Total time on list (days)                | Median (IQR) | 88 (30,198)        | 59 (28,116)     | 75 (29,183)    |
|                                          | Not reported | 1                  | 0               | 1              |
| Donor sex                                | Male         | 41 (37)            | 18 (44)         | 59 (39)        |
|                                          | Female       | 70 (63)            | 23 (56)         | 93 (61)        |
| Donor ethnicity group                    | White        | 98 (88)            | 39 (95)         | 137 (90)       |
|                                          | Other        | 11 (10)            | 1 (2)           | 12 (8)         |
|                                          | Not reported | 2 (2)              | 1 (2)           | 3 (2)          |
| Donor cause of death group               | Stroke       | 95 (86)            | 30 (73)         | 125 (82)       |
|                                          | Trauma       | 7 (6)              | 4 (10)          | 11 (7)         |
|                                          | Other        | 9 (8)              | 7 (17)          | 16 (11)        |
| Donor history of diabetes                | No           | 111 (100)          | 41 (100)        | 152 (100)      |
| Donor age (years)                        | Median (IQR) | 26 (17,44)         | 34 (20,42)      | 30 (19,44)     |
|                                          | Not reported | 0                  | 0               | 0              |
| Donor BMI (kg/m2)                        | Median (IQR) | 21 (19,23)         | 23 (21,23)      | 22 (20,23)     |
|                                          | Not reported | 0                  | 0               | 0              |
| ABO match                                | Identical    | 76 (68)            | 35 (85)         | 111 (73)       |
|                                          | Compatible   | 35 (32)            | 6 (15)          | 41 (27)        |
| Total preservation time (hours)          | Median (IQR) | 5.1 (4.4,5.6)      | 6.3 (5.5,6.7)   | 5.4 (4.7,6.1)  |
|                                          | Not reported | 5                  | 3               | 5              |

## 5.2 Total preservation time, 1 April 2016 – 31 March 2026

**Figure 5.4** shows [boxplots](#) of the [total preservation time](#) of deceased donor organs used in adult intestine transplants over the last 10 years. This is the elapsed time from removal of the organs from the donor to its transplantation into the recipient. The line inside the box indicates the [median](#) value. The median total preservation time has generally remained stable over the decade and was 5.4 hours for 2025/2026.


**Figure 5.4** Median total preservation time in adult intestine transplant:  
1 April 2016 - 31 March 2026



**Figure 5.5** shows the median total preservation times in adult intestine transplants over the last 10 years for each transplant centre. Generally, total preservation times were longer for transplants performed by Oxford.

The total preservation time used is cold ischaemia time as reported on the intestinal transplant record form. These boxplots represent a small number of observations and as shown in **Table 5.1**, several preservation times have not been reported.

**Figure 5.5** Median total preservation time in all adult intestine transplants  
1 April 2016 - 31 March 2026



## 6. Post-transplant survival

One hundred and forty seven of the 152 adult intestinal transplants performed between 1 April 2016 to 31 March 2026 were first graft. This section presents [patient survival](#) after first adult intestine transplantation by transplanting centre and transplant type. Survival information was known for 146 of the 147 transplants performed in the time period. Due to small numbers, [unadjusted survival rates](#) are presented and do not account for differences in the [case mix](#) at each centre.

**Table 6.1** shows the 90-day [patient survival](#) rates by transplant type. There was little evidence of a statistically significant difference in survival between those receiving a liver containing graft and those who did not at 90 days (log-rank  $p=0.07$ ).

**Table 6.2** shows the 90-day [patient survival](#) rates by indication (malignancy versus non-malignant) for recipients who did not receive a liver containing transplant.

| <b>Table 6.1 Unadjusted 90-day patient survival (%) for adult first intestine transplants between 1 April 2016 and 31 March 2026, by transplant type</b> |                              |                                   |                    |
|----------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|-----------------------------------|--------------------|
| <b>Transplant type</b>                                                                                                                                   | <b>Number of transplants</b> | <b>90-day survival (95% CI)</b>   |                    |
|                                                                                                                                                          |                              | <b><a href="#">Unadjusted</a></b> |                    |
| <b>Cambridge</b>                                                                                                                                         |                              |                                   |                    |
| Liver containing <sup>1</sup>                                                                                                                            | 48                           | 91.7                              | (79.3-96.8)        |
| Non liver containing <sup>2</sup>                                                                                                                        | 59                           | 100.0                             | -                  |
| <b>Oxford</b>                                                                                                                                            |                              |                                   |                    |
| Non liver containing <sup>2</sup>                                                                                                                        | 39                           | 94.9                              | (81.0-98.7)        |
| <b>Total</b>                                                                                                                                             |                              |                                   |                    |
| Liver containing <sup>1</sup>                                                                                                                            | 48                           | 91.7                              | (79.3-96.8)        |
| Non liver containing <sup>2</sup>                                                                                                                        | 98                           | 98.0                              | (92.1-99.5)        |
| <b>Total</b>                                                                                                                                             | <b>146</b>                   | <b>95.9</b>                       | <b>(91.0-98.1)</b> |

<sup>1</sup> Includes liver, bowel & pancreas, multivisceral and liver & bowel transplants  
<sup>2</sup> Includes intestine only, bowel & pancreas, modified multivisceral transplants

| <b>Table 6.2 Unadjusted 90-day patient survival (%) for adult first NON-LIVER CONTAINING intestine transplants between 1 April 2016 and 31 March 2026, by indication</b> |                              |                                   |                    |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|-----------------------------------|--------------------|
| <b>Indication</b>                                                                                                                                                        | <b>Number of transplants</b> | <b>90-day survival (95% CI)</b>   |                    |
|                                                                                                                                                                          |                              | <b><a href="#">Unadjusted</a></b> |                    |
| <b>Cambridge</b>                                                                                                                                                         |                              |                                   |                    |
| Malignant indication <sup>1</sup>                                                                                                                                        | 1                            | -                                 | -                  |
| Non-malignant indication                                                                                                                                                 | 58                           | 100.0                             | -                  |
| <b>Oxford</b>                                                                                                                                                            |                              |                                   |                    |
| Malignant indication                                                                                                                                                     | 15                           | 93.3                              | (61.3-99.0)        |
| Non-malignant indication                                                                                                                                                 | 24                           | 95.8                              | (73.9-99.4)        |
| <b>Total</b>                                                                                                                                                             |                              |                                   |                    |
| Malignant indication                                                                                                                                                     | 16                           | 93.8                              | (63.2-99.1)        |
| Non-malignant indication                                                                                                                                                 | 82                           | 98.8                              | (91.6-99.8)        |
| <b>Total</b>                                                                                                                                                             | <b>98</b>                    | <b>98.0</b>                       | <b>(92.1-99.5)</b> |

<sup>1</sup>Survival for transplant types with less than 10 transplants is not presented due to small numbers.

One-year [patient survival](#) rates are shown in **Table 6.3**. The estimated one-year survival rate was 88.1% for patients receiving a non-liver containing graft, and 78.3% for patients receiving a liver containing graft. There was no evidence of a statistically significant difference at a 5% level in survival between those receiving the liver and those who did not at one year post-transplant (log-rank  $p=0.11$ ). **Table 6.4** shows the one-year patient survival rates by indication (malignancy versus non-malignant) for recipients who did not receive a liver containing transplant.

| <b>Table 6.3      Unadjusted one-year patient survival (%) for adult first intestine transplants between 1 April 2016 and 31 March 2026, by transplant type</b> |                              |                                                                      |                    |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|----------------------------------------------------------------------|--------------------|
| <b>Transplant type</b>                                                                                                                                          | <b>Number of transplants</b> | <b>1-year survival (95% CI)</b><br><b><a href="#">Unadjusted</a></b> |                    |
| <b>Cambridge</b>                                                                                                                                                |                              |                                                                      |                    |
| Liver containing <sup>1</sup>                                                                                                                                   | 48                           | 78.3                                                                 | (63.3-87.7)        |
| Non liver containing <sup>2</sup>                                                                                                                               | 59                           | 90.9                                                                 | (79.5-96.1)        |
| <b>Oxford</b>                                                                                                                                                   |                              |                                                                      |                    |
| Non liver containing <sup>2</sup>                                                                                                                               | 39                           | 83.7                                                                 | (67.2-92.3)        |
| <b>Total</b>                                                                                                                                                    |                              |                                                                      |                    |
| <b>Liver containing<sup>1</sup></b>                                                                                                                             | <b>48</b>                    | <b>78.3</b>                                                          | <b>(63.3-87.7)</b> |
| <b>Non liver containing<sup>2</sup></b>                                                                                                                         | <b>98</b>                    | <b>88.1</b>                                                          | <b>(79.5-93.2)</b> |
| <b>Total</b>                                                                                                                                                    | <b>146</b>                   | <b>84.6</b>                                                          | <b>(77.4-89.7)</b> |

<sup>1</sup> Includes liver, bowel & pancreas, multivisceral and liver & bowel transplants  
<sup>2</sup> Includes intestine only, bowel & pancreas, modified multivisceral transplants

| <b>Table 6.4      Unadjusted one-year patient survival (%) for adult first NON-LIVER CONTAINING intestine transplants between 1 April 2016 and 31 March 2026, by indication</b> |                              |                                                                      |                    |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|----------------------------------------------------------------------|--------------------|
| <b>Indication</b>                                                                                                                                                               | <b>Number of transplants</b> | <b>1-year survival (95% CI)</b><br><b><a href="#">Unadjusted</a></b> |                    |
| <b>Cambridge</b>                                                                                                                                                                |                              |                                                                      |                    |
| Malignant indication <sup>1</sup>                                                                                                                                               | 1                            | -                                                                    | -                  |
| Non-malignant indication                                                                                                                                                        | 58                           | 90.7                                                                 | (79.2-96.0)        |
| <b>Oxford</b>                                                                                                                                                                   |                              |                                                                      |                    |
| Malignant indication                                                                                                                                                            | 15                           | 73.3                                                                 | (43.6-89.1)        |
| Non-malignant indication                                                                                                                                                        | 24                           | 91.0                                                                 | (68.6-97.7)        |
| <b>Total</b>                                                                                                                                                                    |                              |                                                                      |                    |
| <b>Malignant indication</b>                                                                                                                                                     | <b>16</b>                    | <b>75.0</b>                                                          | <b>(46.3-89.8)</b> |
| <b>Non-malignant indication</b>                                                                                                                                                 | <b>82</b>                    | <b>90.8</b>                                                          | <b>(81.6-95.5)</b> |
| <b>Total</b>                                                                                                                                                                    | <b>98</b>                    | <b>88.1</b>                                                          | <b>(79.5-93.2)</b> |

<sup>1</sup>Survival for transplant types with less than 10 transplants is not presented due to small numbers.

Five-year [patient survival rates](#) are shown in **Table 6.5**. The survival rate was 71.2% for patients not receiving the liver, and was 55.1% for patients receiving the liver. There was a evidence of a statistically significant difference at a 5% level in survival between those receiving the liver and those who did not at 5 years (log-rank p=0.03). **Table 6.6** shows the breakdown of survival rates by malignancy, for recipients not receiving the liver.

| <b>Table 6.5      Unadjusted five-year patient survival (%) for adult first intestine transplants between 1 April 2016 and 31 March 2026, by transplant type</b> |                              |                                                       |                    |
|------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|-------------------------------------------------------|--------------------|
| <b>Transplant type</b>                                                                                                                                           | <b>Number of transplants</b> | <b>5-year survival (95% CI)<br/><u>Unadjusted</u></b> |                    |
| <b>Cambridge</b>                                                                                                                                                 |                              |                                                       |                    |
| Liver containing <sup>1</sup>                                                                                                                                    | 48                           | 55.9                                                  | (38.4-70.2)        |
| Non liver containing <sup>2</sup>                                                                                                                                | 59                           | 85.4                                                  | (71.3-92.9)        |
| <b>Oxford</b>                                                                                                                                                    |                              |                                                       |                    |
| Non liver containing <sup>2</sup>                                                                                                                                | 39                           | 54.6                                                  | (33.6-71.4)        |
| <b>Total</b>                                                                                                                                                     |                              |                                                       |                    |
| <b>Liver containing<sup>1</sup></b>                                                                                                                              | <b>48</b>                    | <b>55.9</b>                                           | <b>(38.4-70.2)</b> |
| <b>Non liver containing<sup>2</sup></b>                                                                                                                          | <b>98</b>                    | <b>71.6</b>                                           | <b>(58.6-81.2)</b> |
| <b>Total</b>                                                                                                                                                     | <b>146</b>                   | <b>66.3</b>                                           | <b>(56.1-74.7)</b> |

<sup>1</sup> Includes liver, bowel & pancreas, multivisceral and liver & bowel transplants  
<sup>2</sup> Includes intestine only, bowel & pancreas, modified multivisceral transplants

| <b>Table 6.6      Unadjusted five-year patient survival (%) for adult first NON-LIVER CONTAINING intestine transplants between 1 April 2016 and 31 March 2026, by indication</b> |                              |                                                       |                    |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|-------------------------------------------------------|--------------------|
| <b>Indication</b>                                                                                                                                                                | <b>Number of transplants</b> | <b>5-year survival (95% CI)<br/><u>Unadjusted</u></b> |                    |
| <b>Cambridge</b>                                                                                                                                                                 |                              |                                                       |                    |
| Malignant indication <sup>1</sup>                                                                                                                                                | 1                            | -                                                     | -                  |
| Non-malignant indication                                                                                                                                                         | 58                           | 85.1                                                  | (70.7-92.7)        |
| <b>Oxford</b>                                                                                                                                                                    |                              |                                                       |                    |
| Malignant indication                                                                                                                                                             | 15                           | 37.7                                                  | (12.5-63.3)        |
| Non-malignant indication                                                                                                                                                         | 24                           | 68.8                                                  | (39.0-86.2)        |
| <b>Total</b>                                                                                                                                                                     |                              |                                                       |                    |
| Malignant indication                                                                                                                                                             | <b>16</b>                    | <b>42.6</b>                                           | <b>(16.6-66.7)</b> |
| Non-malignant indication                                                                                                                                                         | <b>82</b>                    | <b>79.8</b>                                           | <b>(66.3-88.4)</b> |
| <b>Total</b>                                                                                                                                                                     | <b>98</b>                    | <b>71.6</b>                                           | <b>(58.6-81.2)</b> |

<sup>1</sup>Survival rates for transplant types with less than 10 transplants are not presented due to small numbers.

# **PAEDIATRIC INTESTINE TRANSPLANTATION**




## 7. Transplant list

### 7.1 Paediatric intestine transplant list as at 31 March, 2017 – 2026


**Figure 7.1** shows the number of paediatric patients (aged <18 years) active and suspended on the intestine transplant list at 31 March of each year between 2017 and 2026. The number of paediatric patients on the [active transplant list](#) ranged between 2 and 9 each year and was 4 on 31 March 2026.

**Figure 7.1** Paediatric patients on the intestine transplant list at 31 March, 2017 - 2026



**Figure 7.2** shows the number of paediatric patients on the intestine transplant list at 31 March of each year between 2017 and 2026, at each transplant centre. The number of paediatric patients on the active transplant list at 31 March 2026 was 4 with 2 patients at Birmingham and 2 at King's college.

**Figure 7.2** Paediatric patients on the intestine transplant list at 31 March, by centre, 2017 - 2026




**Table 7.1** shows the demographic characteristics of 74 paediatric intestine transplant recipient registrations in the 10-year period by centre and overall. Nationally, 54% of registrations were male and the [median](#) age was 3 years old. The most commonly reported indication for transplantation was short bowel syndrome (46%). The median recipient BMI was 16 kg/m<sup>2</sup>. For some characteristics, percentages may not add up to 100 due to rounding.

| <b>Table 7.1 Demographic characteristics of paediatric intestine transplant recipient registrations, 1 April 2016 - 31 March 2026</b> |                           |            |                         |                   |
|---------------------------------------------------------------------------------------------------------------------------------------|---------------------------|------------|-------------------------|-------------------|
|                                                                                                                                       |                           | Birmingham | King's College Hospital | TOTAL             |
|                                                                                                                                       |                           | N (%)      | N (%)                   | N (%)             |
| Number of registrations                                                                                                               |                           | 33         | 41                      | <b>74 (100)</b>   |
| Number of patients                                                                                                                    |                           | 31 (100)   | 35 (100)                | <b>66 (100)</b>   |
| Registration type                                                                                                                     | Elective                  | 33 (100)   | 41 (100)                | <b>74 (100)</b>   |
| Transplant type required                                                                                                              | Non liver containing      | 10 (30)    | 10 (24)                 | <b>20 (27)</b>    |
|                                                                                                                                       | Liver containing          | 23 (70)    | 31 (76)                 | <b>54 (73)</b>    |
| Recipient sex                                                                                                                         | Male                      | 19 (58)    | 21 (51)                 | <b>40 (54)</b>    |
|                                                                                                                                       | Female                    | 14 (42)    | 20 (49)                 | <b>34 (46)</b>    |
| Recipient ethnicity group                                                                                                             | White                     | 28 (85)    | 28 (68)                 | <b>56 (76)</b>    |
|                                                                                                                                       | Other                     | 3 (9)      | 13 (32)                 | <b>16 (22)</b>    |
|                                                                                                                                       | Not reported              | 2 (6)      | 0                       | <b>2 (3)</b>      |
| Indication group                                                                                                                      | Short bowel syndrome      | 18 (55)    | 16 (39)                 | <b>34 (46)</b>    |
|                                                                                                                                       | Motility disorders        | 7 (21)     | 7 (17)                  | <b>14 (19)</b>    |
|                                                                                                                                       | Primary mucosal disorders | 1 (3)      | 3 (7)                   | <b>4 (5)</b>      |
|                                                                                                                                       | Tumour                    | 3 (9)      | 1 (2)                   | <b>4 (5)</b>      |
|                                                                                                                                       | Other/not reported        | 4 (12)     | 5 (12)                  | <b>9 (12)</b>     |
|                                                                                                                                       | Retransplant              | 0          | 9 (22)                  | <b>9 (12)</b>     |
| Patient location                                                                                                                      | Out-patient               | 21 (64)    | 18 (44)                 | <b>39 (53)</b>    |
|                                                                                                                                       | Ward                      | 3 (9)      | 9 (22)                  | <b>12 (16)</b>    |
|                                                                                                                                       | ICU/HDU                   | 1 (3)      | 1 (2)                   | <b>2 (3)</b>      |
|                                                                                                                                       | Not reported              | 8 (24)     | 13 (32)                 | <b>21 (28)</b>    |
| Pre-transplant renal support                                                                                                          | No                        | 33 (100)   | 40 (98)                 | <b>73 (99)</b>    |
|                                                                                                                                       | Yes                       | 0          | 1 (2)                   | <b>1 (1)</b>      |
| Previous abdominal surgery                                                                                                            | No                        | 4 (12)     | 4 (10)                  | <b>8 (11)</b>     |
|                                                                                                                                       | Yes                       | 29 (88)    | 37 (90)                 | <b>66 (89)</b>    |
| Recipient blood group                                                                                                                 | O                         | 16 (48)    | 15 (37)                 | <b>31 (42)</b>    |
|                                                                                                                                       | A                         | 12 (36)    | 19 (46)                 | <b>31 (42)</b>    |
|                                                                                                                                       | B                         | 1 (3)      | 5 (12)                  | <b>6 (8)</b>      |
|                                                                                                                                       | AB                        | 4 (12)     | 2 (5)                   | <b>6 (8)</b>      |
| Recipient age (years)                                                                                                                 | Median (IQR)              | 5 (2,10)   | 3 (1,5)                 | <b>3 (2,7)</b>    |
| Recipient BMI (kg/m2)                                                                                                                 | Median (IQR)              | 16 (15,19) | 16 (16,18)              | <b>16 (15,18)</b> |

An indication of registration outcomes for paediatric elective patients registered on the intestine transplant list in the UK between 1 April 2022 and 31 March 2024 is summarised in **Figure 7.3**. The proportion of paediatric patients transplanted within six months of listing was 53%.

**Figure 7.3** Post-registration outcome for 19 elective paediatric registrations for an intestine transplant made in the UK, 1 April 2022 - 31 March 2024



**Table 7.2** shows [median waiting time](#) to [elective](#) intestine transplant by registration type for paediatric patients, registered between 1 April 2022 and 31 March 2025. A small proportion of patients requiring a liver are removed from the intestinal transplant waiting list to receive a liver transplant without the small bowel; the liver transplant is counted as receiving a transplant. Registrations ending in a living donor transplant are excluded. The national UK median total waiting time to transplant for paediatric elective patients is 316 days.

| Table 7.2 Median waiting time to elective intestine transplant in the UK, for paediatric patients registered 1 April 2022 - 31 March 2025 |                               |                                        |                         |                            |                         |
|-------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------|----------------------------------------|-------------------------|----------------------------|-------------------------|
| Registration type                                                                                                                         | Number of patients registered | Total waiting time (days) <sup>1</sup> |                         | Active waiting time (days) |                         |
|                                                                                                                                           |                               | Median                                 | 95% Confidence interval | Median                     | 95% Confidence interval |
| Liver containing                                                                                                                          | 18                            | 316                                    | 203 – 429               | 261                        | 142 – 380               |
| Non liver containing                                                                                                                      | 3                             | 82                                     | 24 - 140                | 82                         | 24 – 140                |
| <b>Total</b>                                                                                                                              | <b>21</b>                     | <b>316</b>                             | <b>191 - 441</b>        | <b>261</b>                 | <b>168 - 354</b>        |

<sup>1</sup> Periods of suspension from the list are included in the calculation of total median waiting times  
<sup>2</sup> Patients who received a liver only transplant are classed as transplanted

## 8. Response to offers

Between 1 April 2025 and 31 March 2026, Birmingham and King's College Hospital received offers from 104 and 5 donors, respectively, for intestine transplant patients at their centres. Their [donor offer decline rates](#) were 98% and 100% respectively. These rates are high to some extent because centres are very selective, particularly about the size of the donor.


| Table 8.1 Offer decline rates for each centre, 1 April 2025 - 31 March 2026 |            |           |            |           |
|-----------------------------------------------------------------------------|------------|-----------|------------|-----------|
| Centre                                                                      | Offers     | %Decline  | Donors     | %Decline  |
| Birmingham                                                                  | 106        | 98        | 104        | 98        |
| King's College                                                              | 5          | 100       | 5          | 100       |
| <b>Total</b>                                                                | <b>111</b> | <b>98</b> | <b>109</b> | <b>98</b> |

## 9. Transplants

### 9.1 Paediatric intestine transplants, 1 April 2016 – 31 March 2026

**Figure 9.1** shows the number of paediatric intestine transplants performed in the last 10 years, by [transplant type](#). The number of paediatric transplants has fluctuated over the period, with 2 performed in 2025/2026. In 2017/2018, a living liver and bowel transplant was performed by King's College Hospital and in 2023/2024, a living bowel-only transplant was performed by King's College Hospital. These transplants are included in the numbers presented in this section.


**Figure 9.1 Paediatric intestine transplants in the UK 1 April 2016 - 31 March 2026**



Transplants may also include any of stomach, spleen, abdominal wall, kidney, color

**Figure 9.2** shows the number of paediatric intestine transplants performed in 2025/2026, by centre and [transplant type](#). Birmingham performed two intestinal transplants while none were reported as performed at King's College Hospital. One of the transplants were non liver containing and one included the liver.


**Figure 9.2 Paediatric intestine transplants in the UK by centre 1 April 2025 - 31 March 2026**



Transplants may also include any of stomach, spleen, abdominal wall, kidney, color

**Figure 9.3** shows the number of paediatric intestine transplants performed in the last 10 years, by centre and [type of transplant](#). The number of transplants performed by both centres fluctuated over the last 10 years. King's College performed 25 (51%) of transplants, whilst Birmingham performed 24 (49%) of transplants.

**Figure 9.3 Paediatric intestine transplants by centre, 1 April 2016 - 31 March 2026**



Transplants may also include any of stomach, spleen, abdominal wall, kidney, color

The demographic characteristics of 49 paediatric intestine transplant recipients in the 10-year period are shown by centre and overall in **Table 9.1**. Nationally, 57% of recipients were male and the [median](#) recipient age was 6 years old, while the median age of donors was 7 years old. The most common reported indication for transplantation was short bowel syndrome. All transplants were performed in [elective](#) recipients. For some characteristics, percentages may not add up to 100 due to rounding.

| <b>Table 9.1 Demographic characteristics of paediatric elective intestine transplant recipients, 1 April 2016 - 31 March 2026</b> |                           |                     |                         |                 |
|-----------------------------------------------------------------------------------------------------------------------------------|---------------------------|---------------------|-------------------------|-----------------|
|                                                                                                                                   |                           | Birmingham<br>N (%) | King's College<br>N (%) | TOTAL<br>N (%)  |
| Number of transplants                                                                                                             |                           | 24                  | 25                      | <b>49 (100)</b> |
| Transplant type                                                                                                                   | Non liver containing      | 8 (33)              | 10 (40)                 | <b>18 (37)</b>  |
|                                                                                                                                   | Liver containing          | 16 (67)             | 15 (60)                 | <b>31 (63)</b>  |
| Urgency                                                                                                                           | Elective                  | 24 (100)            | 25 (100)                | <b>49 (100)</b> |
| Recipient sex                                                                                                                     | Male                      | 14 (58)             | 14 (56)                 | <b>28 (57)</b>  |
|                                                                                                                                   | Female                    | 10 (42)             | 11 (44)                 | <b>21 (43)</b>  |
| Recipient ethnicity group                                                                                                         | White                     | 18 (75)             | 18 (72)                 | <b>36 (74)</b>  |
|                                                                                                                                   | Other                     | 4 (17)              | 7 (28)                  | <b>11 (22)</b>  |
|                                                                                                                                   | Not reported              | 2 (8)               | 0                       | <b>2 (4)</b>    |
| Indication group                                                                                                                  | Short bowel syndrome      | 12 (50)             | 6 (24)                  | <b>18 (37)</b>  |
|                                                                                                                                   | Motility disorders        | 4 (17)              | 6 (24)                  | <b>10 (20)</b>  |
|                                                                                                                                   | Primary mucosal disorders | 1 (4)               | 2 (8)                   | <b>3 (6)</b>    |
|                                                                                                                                   | Tumour                    | 3 (13)              | 1 (4)                   | <b>4 (8)</b>    |
|                                                                                                                                   | Other/not reported        | 3 (13)              | 4 (16)                  | <b>7 (14)</b>   |
|                                                                                                                                   | Retransplant              | 1 (4)               | 6 (24)                  | <b>7 (14)</b>   |
| Patient location                                                                                                                  | Out-patient               | 19 (79)             | 14 (56)                 | <b>33 (67)</b>  |
|                                                                                                                                   | Ward                      | 4 (17)              | 4 (16)                  | <b>8 (16)</b>   |
|                                                                                                                                   | ICU/HDU                   | 1 (4)               | 3 (12)                  | <b>4 (8)</b>    |
|                                                                                                                                   | Not reported              | 0                   | 4 (16)                  | <b>4 (8)</b>    |
| Pre-transplant renal support                                                                                                      | No                        | 23 (96)             | 21 (84)                 | <b>44 (90)</b>  |
|                                                                                                                                   | Yes                       | 1 (4)               | 0                       | <b>1 (2)</b>    |
|                                                                                                                                   | Not reported              | 0                   | 4 (16)                  | <b>4 (8)</b>    |
| Previous abdominal surgery                                                                                                        | No                        | 7 (29)              | 2 (8)                   | <b>9 (18)</b>   |
|                                                                                                                                   | Yes                       | 17 (71)             | 19 (76)                 | <b>36 (74)</b>  |
|                                                                                                                                   | Not reported              | 0                   | 4 (16)                  | <b>4 (8)</b>    |
| Life style activity                                                                                                               | Normal                    | 2 (8)               | 1 (4)                   | <b>3 (6)</b>    |
|                                                                                                                                   | Restricted                | 9 (38)              | 7 (28)                  | <b>16 (33)</b>  |
|                                                                                                                                   | Self-care                 | 3 (13)              | 0                       | <b>3 (6)</b>    |
|                                                                                                                                   | Reliant                   | 0                   | 2 (8)                   | <b>2 (4)</b>    |
|                                                                                                                                   | Aged five years or less   | 9 (38)              | 11 (44)                 | <b>20 (41)</b>  |
|                                                                                                                                   | Not reported              | 1 (4)               | 4 (16)                  | <b>5 (10)</b>   |
| Restricted venous access at registration                                                                                          | No                        | 15 (63)             | 15 (60)                 | <b>30 (61)</b>  |
|                                                                                                                                   | Yes                       | 9 (38)              | 9 (36)                  | <b>18 (37)</b>  |
|                                                                                                                                   | Not reported              | 0                   | 1 (4)                   | <b>1 (2)</b>    |


**Table 9.1 Demographic characteristics of paediatric elective intestine transplant recipients, 1 April 2016 - 31 March 2026**

|                                             |                                     | Birmingham<br>N (%)             | King's College<br>N (%)         | TOTAL<br>N (%)                                 |
|---------------------------------------------|-------------------------------------|---------------------------------|---------------------------------|------------------------------------------------|
| Recipient age (years)                       | Median (IQR)                        | 7 (3,10)                        | 5 (3,9)                         | <b>6 (3,10)</b>                                |
| Recipient BMI (kg/m2)                       | Median (IQR)<br>Not reported        | 17 (15,18)<br>0                 | 16 (16,19)<br>1                 | <b>17 (16,19)<br/>1</b>                        |
| Serum bilirubin (umol/l)                    | Median (IQR)<br>Not reported        | 15 (8,30)<br>2                  | 10 (9,25)<br>4                  | <b>13 (8,30)<br/>6</b>                         |
| Serum bilirubin (umol/l)<br>non-liver       | Median (IQR)<br>Not reported        | 15 (9,17)<br>2                  | 9 (3,20)<br>3                   | <b>9 (7,17)<br/>5</b>                          |
| Serum bilirubin (umol/l)<br>including liver | Median (IQR)<br>Not reported        | 14 (7,155)<br>0                 | 12 (9, 34)<br>1                 | <b>14 (9, 41)<br/>1</b>                        |
| Total time on list (days)                   | Median (IQR)<br>Not reported        | 150 (44,264)<br>0               | 276 (129,788)<br>2              | <b>190 (71,545)<br/>2</b>                      |
| Donor sex                                   | Male<br>Female                      | 15 (63)<br>9 (38)               | 11 (44)<br>14 (56)              | <b>26 (53)<br/>23 (47)</b>                     |
| Donor ethnicity group                       | White<br>Other<br>Not reported      | 19 (79)<br>1 (4)<br>4 (17)      | 16 (64)<br>3 (12)<br>6 (24)     | <b>35 (71)<br/>4 (8)<br/>10 (20)</b>           |
| Donor cause of death<br>group               | Stroke<br>Trauma<br>Other<br>Living | 20 (83)<br>1 (4)<br>3 (13)<br>0 | 16 (64)<br>0<br>7 (28)<br>2 (8) | <b>36 (74)<br/>1 (2)<br/>10 (20)<br/>2 (4)</b> |
| Donor history of diabetes                   | No<br>Not reported                  | 23 (96)<br>1 (4)                | 19 (76)<br>6 (24)               | <b>42 (86)<br/>7 (14)</b>                      |
| Donor age (years)                           | Median (IQR)                        | 11 (6,24)                       | 5 (2,36)                        | <b>7 (2,29)</b>                                |
| Donor BMI (kg/m2)                           | Median (IQR)<br>Not reported        | 19 (16,20)<br>0                 | 16 (15,20)<br>2                 | <b>18 (15,20)<br/>2</b>                        |
| ABO match                                   | Identical<br>Compatible             | 19 (79)<br>5 (21)               | 20 (80)<br>5 (20)               | <b>39 (80)<br/>10 (20)</b>                     |
| Total preservation time<br>(hours)          | Median (IQR)<br>Not reported        | 5.3 (4.2,6.1)<br>0              | 8.4 (7.2,9.8)<br>7              | <b>6.3 (5.2,8.2)<br/>7</b>                     |

## 9.2 Total preservation time, 1 April 2016 – 31 March 2026


**Figure 9.4** shows [boxplots](#) of the [total preservation times](#) of deceased donor organs used in paediatric intestine transplants over the last 10 years. This is the elapsed time from removal of the organs from the donor to removal from ice prior to implant. The line inside the box indicates the [median](#) value. The median total preservation time in paediatric transplants was 7.1 hours in 2025/2026.

**Figure 9.4** Median total preservation time in paediatric intestine transplants  
1 April 2016 - 31 March 2026



**Figure 9.5** shows the median total preservation times in paediatric intestine transplants by centre over the last 10 years for deceased donors. These boxplots represent a small number of observations and, as shown in **Table 9.1**, several preservation times have not been reported.

**Figure 9.5** Median total preservation time in all paediatric intestine transplants  
1 April 2016 - 31 March 2026



## 10. Post-transplant survival

This section presents [patient survival](#) post paediatric intestine transplantation for first intestine transplants between 1 April 2016 and 31 March 2026, by transplanting centre and transplant type. Of the 40 transplants of this kind in the time period, survival information was known for all cases. Due to small numbers, [unadjusted survival rates](#) are presented, which do not account for differences in the [case mix](#) at each centre and transplant type.

### 10.1 Survival by transplant type

**Table 10.1** shows the 90-day [unadjusted patient survival rates](#) by transplant type. The 90-day survival estimate was 96.0% for patients who received a liver and 100% for patients not receiving a liver. There was little evidence of a difference in survival between those who received a liver and those who did not at 90 days (log-rank  $p=0.44$ ).

| <b>Table 10.1 90-day patient survival (%) for paediatric first intestine transplants between 1 April 2016 and 31 March 2026, by transplant type</b> |                              |                                                                |                    |
|-----------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|----------------------------------------------------------------|--------------------|
| <b>Transplant type</b>                                                                                                                              | <b>Number of transplants</b> | <b>90-day survival (95% CI)<br/><a href="#">Unadjusted</a></b> |                    |
| <b>Birmingham</b>                                                                                                                                   |                              |                                                                |                    |
| Liver containing <sup>1</sup>                                                                                                                       | 14                           | 92.9                                                           | (59.1-99.0)        |
| Non liver containing <sup>2</sup>                                                                                                                   | 8 <sup>3</sup>               | -                                                              | -                  |
| <b>King's College Hospital</b>                                                                                                                      |                              |                                                                |                    |
| Liver containing <sup>1</sup>                                                                                                                       | 11                           | 100.0                                                          | -                  |
| Non liver containing <sup>2</sup>                                                                                                                   | 7 <sup>3</sup>               | -                                                              | -                  |
| <b>Total</b>                                                                                                                                        |                              |                                                                |                    |
| Liver containing <sup>1</sup>                                                                                                                       | <b>25</b>                    | <b>96.0</b>                                                    | <b>(74.8-99.4)</b> |
| Non liver containing <sup>2</sup>                                                                                                                   | <b>15</b>                    | <b>100.0</b>                                                   | <b>-</b>           |
| <b>Total</b>                                                                                                                                        | <b>40</b>                    | <b>97.5</b>                                                    | <b>(83.5-99.6)</b> |

<sup>1</sup> Includes liver, bowel & pancreas, multivisceral and liver & bowel transplants  
<sup>2</sup> Includes intestine only, bowel & pancreas, modified multivisceral transplants  
<sup>3</sup> Survival rates for transplant types with less than 10 transplants are not presented due to small numbers.

One-year [patient survival rates](#) are shown in **Table 10.2**. One year survival was 100.0% for patients not receiving a liver and 83.5% for patients receiving a liver. There was little evidence of a difference in survival between those who received a liver and those who did not at 1 year (log-rank  $p=0.11$ ).

| <b>Table 10.2 1-year patient survival (%) for paediatric first intestine transplants between 1 April 2016 and 31 March 2026, by transplant type</b> |                              |                                                       |                    |
|-----------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|-------------------------------------------------------|--------------------|
| <b>Transplant type</b>                                                                                                                              | <b>Number of transplants</b> | <b>1-year survival (95% CI)<br/><u>Unadjusted</u></b> |                    |
| <b>Birmingham</b>                                                                                                                                   |                              |                                                       |                    |
| Liver containing <sup>1</sup>                                                                                                                       | 14                           | 85.1                                                  | (52.3-96.1)        |
| Non liver containing <sup>2</sup>                                                                                                                   | 8 <sup>3</sup>               | -                                                     | -                  |
| <b>King's College Hospital</b>                                                                                                                      |                              |                                                       |                    |
| Liver containing <sup>1</sup>                                                                                                                       | 11                           | 81.8                                                  | (44.7-95.1)        |
| Non liver containing <sup>2</sup>                                                                                                                   | 7 <sup>3</sup>               | -                                                     | -                  |
| <b>Total</b>                                                                                                                                        |                              |                                                       |                    |
| <b>Liver containing<sup>1</sup></b>                                                                                                                 | <b>25</b>                    | <b>83.5</b>                                           | <b>(61.7-93.5)</b> |
| <b>Non liver containing<sup>2</sup></b>                                                                                                             | <b>15</b>                    | <b>100.0</b>                                          | <b>-</b>           |
| <b>Total</b>                                                                                                                                        | <b>40</b>                    | <b>89.6</b>                                           | <b>(74.6-96.0)</b> |
| <sup>1</sup> Includes liver, bowel & pancreas, multivisceral and liver & bowel transplants                                                          |                              |                                                       |                    |
| <sup>2</sup> Includes intestine only, bowel & pancreas, modified multivisceral transplants                                                          |                              |                                                       |                    |
| <sup>3</sup> Survival for transplant types with less than 10 transplants are not presented due to small numbers.                                    |                              |                                                       |                    |

Five-year [patient survival](#) rates are shown in **Table 10.3**. At five years post-transplant, the survival was 79.5% for patients not receiving a liver and was 67.8% for patients receiving a liver. There was little evidence of a difference in survival between those who received a liver and those who did not at 5 years (log-rank p=0.38).

| <b>Table 10.3 5-year patient survival (%) for paediatric first intestine transplants between 1 April 2016 and 31 March 2026, by transplant type</b> |                              |                                                       |                    |
|-----------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------|-------------------------------------------------------|--------------------|
| <b>Transplant type</b>                                                                                                                              | <b>Number of transplants</b> | <b>5-year survival (95% CI)<br/><u>Unadjusted</u></b> |                    |
| <b>Birmingham</b>                                                                                                                                   |                              |                                                       |                    |
| Liver containing <sup>1</sup>                                                                                                                       | 14                           | 77.4                                                  | (44.9-92.1)        |
| Non liver containing <sup>2</sup>                                                                                                                   | 8 <sup>3</sup>               | -                                                     | -                  |
| <b>King's College Hospital</b>                                                                                                                      |                              |                                                       |                    |
| Liver containing <sup>1</sup>                                                                                                                       | 11                           | 57.3                                                  | (20.5-82.2)        |
| Non liver containing <sup>2</sup>                                                                                                                   | 7 <sup>3</sup>               | -                                                     | -                  |
| <b>Total</b>                                                                                                                                        |                              |                                                       |                    |
| <b>Liver containing<sup>1</sup></b>                                                                                                                 | <b>25</b>                    | <b>67.8</b>                                           | <b>(43.3-83.5)</b> |
| <b>Non liver containing<sup>2</sup></b>                                                                                                             | <b>15</b>                    | <b>79.5</b>                                           | <b>(39.3-94.5)</b> |
| <b>Total</b>                                                                                                                                        | <b>40</b>                    | <b>71.7</b>                                           | <b>(51.6-84.6)</b> |
| <sup>1</sup> Includes liver, bowel & pancreas, multivisceral and liver & bowel transplants                                                          |                              |                                                       |                    |
| <sup>2</sup> Includes intestine only, bowel & pancreas, modified multivisceral transplants                                                          |                              |                                                       |                    |
| <sup>3</sup> Survival for transplant types with less than 10 transplants are not presented due to small numbers.                                    |                              |                                                       |                    |

# FORM RETURN RATES



## 11.1 Form return rates, 1 January 2025 – 31 December 2025

Form return rates are reported by follow-up centre in **Table 11.1** for adult transplants and **Table 11.2** for paediatric transplants. The forms included are the intestine transplant record form and the three month and annual intestine transplant follow-up forms that are reported to the UK Transplant Registry database. The tables show the number of forms issued between 1 January 2025 and 31 December 2025, for recipients being followed up by each centre, and the percentage of forms that had been returned at time of analysis (8 July 2025). Annual follow-up forms are broken down into those issued at one year post-transplant and “lifetime follow-up”, which is two years or longer. Note that some paediatric transplant patients have transitioned to be followed up at Cambridge and Oxford, both of which are adult transplant centres.

| Table 11.1 Form return rates by follow-up centre, 1 January 2025 to 31 December 2025 |                   |            |                   |            |                  |            |                    |            |
|--------------------------------------------------------------------------------------|-------------------|------------|-------------------|------------|------------------|------------|--------------------|------------|
| Follow-up centre                                                                     | Transplant record |            | 3 month follow-up |            | 1 year follow-up |            | Lifetime follow-up |            |
|                                                                                      | No. forms issued  | % returned | No. forms issued  | % returned | No. forms issued | % returned | No. forms issued   | % returned |
| Cambridge                                                                            | 7                 | 100        | 11                | 91         | 13               | 100        | 71                 | 97         |
| Oxford                                                                               | 4                 | 100        | 3                 | 100        | 1                | 100        | 27                 | 100        |
| <b>Total</b>                                                                         | <b>11</b>         | <b>100</b> | <b>14</b>         | <b>93</b>  | <b>14</b>        | <b>100</b> | <b>98</b>          | <b>98</b>  |

APPENDIX

| Table 11.2 Form return rates by follow-up centre, 1 January 2025 to 31 December 2025 |                   |            |                   |            |                  |            |                    |            |
|--------------------------------------------------------------------------------------|-------------------|------------|-------------------|------------|------------------|------------|--------------------|------------|
| Follow-up centre                                                                     | Transplant record |            | 3 month follow-up |            | 1 year follow-up |            | Lifetime follow-up |            |
|                                                                                      | No. forms issued  | % returned | No. forms issued  | % returned | No. forms issued | % returned | No. forms issued   | % returned |
| Birmingham                                                                           | 2                 | 100        | 2                 | 100        | 3                | 100        | 21                 | 100        |
| Cambridge                                                                            | 0                 | -          | 0                 | -          | 0                | -          | 15                 | 100        |
| King's College Hospital                                                              | 1                 | 100        | 1                 | 100        | 2                | 100        | 13                 | 100        |
| Oxford                                                                               | 0                 | -          | 0                 | -          | 0                | -          | 16                 | 100        |
| <b>Total</b>                                                                         | <b>3</b>          | <b>100</b> | <b>3</b>          | <b>100</b> | <b>5</b>         | <b>100</b> | <b>65</b>          | <b>100</b> |



## A1: Number of patients analysed

Data were obtained from the UK Transplant Registry for the 10-year time period, 1 April 2016 to 31 March 2026. NHS Group 2 transplants have been included in all analyses (except Section 2.3 Geographical variation in registration and transplant rates), while liver-only transplants because of intestine failure have been excluded. One transplant at Oxford, in 2016, involving the abdominal wall only where the patient was registered using the Intestinal Failure Transplant Recipient Registration form has been excluded. Those who are registered for an intestine transplant but receive a liver only transplant are included in the transplant list activity but are excluded in the transplant activity and post-transplant survival sections.

**Table A1** shows the number of adult transplants including the intestine in the 10-year period by centre and urgency status. The left hand columns show the total number of transplants (including re-transplants) and the right hand columns show first-time transplants only.

| <b>Table A1    Number of adult intestine transplants, by transplant centre and urgency status, 1 April 2016 to 31 March 2026</b> |                        |                     |                               |                     |
|----------------------------------------------------------------------------------------------------------------------------------|------------------------|---------------------|-------------------------------|---------------------|
| <b>Centre</b>                                                                                                                    | <b>All transplants</b> |                     | <b>First-time transplants</b> |                     |
|                                                                                                                                  | <b>Elective</b>        | <b>Super-urgent</b> | <b>Elective</b>               | <b>Super-urgent</b> |
| Cambridge                                                                                                                        | 109                    | 2                   | 106                           | 2                   |
| Oxford                                                                                                                           | 41                     | 0                   | 39                            | 0                   |
| <b>TOTAL</b>                                                                                                                     | <b>150</b>             | <b>2</b>            | <b>145</b>                    | <b>2</b>            |

**Table A2** shows the number of paediatric transplants including the intestine in the 10-year period by centre and urgency status. The left hand columns show the total number of transplants (including re-transplants) and the right hand columns shows first-time transplants only.

| <b>Table A2    Number of paediatric intestine transplants, by transplant centre and urgency status, 1 April 2016 to 31 March 2026</b> |                        |                     |                               |                     |
|---------------------------------------------------------------------------------------------------------------------------------------|------------------------|---------------------|-------------------------------|---------------------|
| <b>Centre</b>                                                                                                                         | <b>All transplants</b> |                     | <b>First-time transplants</b> |                     |
|                                                                                                                                       | <b>Elective</b>        | <b>Super-urgent</b> | <b>Elective</b>               | <b>Super-urgent</b> |
| Birmingham                                                                                                                            | 24                     | 0                   | 22                            | 0                   |
| King's College Hospital                                                                                                               | 25                     | 0                   | 18                            | 0                   |
| <b>TOTAL</b>                                                                                                                          | <b>49</b>              | <b>0</b>            | <b>40</b>                     | <b>0</b>            |

## **Geographical variation analysis**

### **Registration rates**

All NHS group 1 patients who were registered onto the intestinal transplant lists with an active status between 1 April 2016 and 31 March 2026 were extracted from the UK Transplant Registry on 10 July 2026 (numerator). Patients were assigned to NHS regions in England using their postcode of residence, as reported at registration. The number of registrations per million population (pmp) by NHS region was obtained using mid-2025 population estimates based on the Office for National Statistics (ONS) 2021 Census figures (denominator). No NHS region age- or sex-specific standardisation of rates was performed.

The registration rates pmp were categorised into four groups – low, low-medium, medium-high and high – based on the quartiles of their distribution and visualised in a map using contrasting colours.

### **Transplant rates**

Transplant rates pmp were obtained as the number of intestinal transplants on NHS group 1 recipients from deceased donors between 1 April 2016 and 31 March 2026 (numerator), divided by the mid-2025 population estimates from the ONS (denominator). Transplant rates pmp were categorised and visualised in a map as done for the registration rates.

### **Systematic component of variation**

Only registrations or transplants in England between 1 April 2016 and 31 March 2026 were included. If a patient was re-registered during the time period, only the first registration was considered. If a patient underwent more than one intestinal transplant in the time period, only the first transplant was considered.

## A2: Methods

### Waiting time to transplant

Waiting time was calculated from date of registration to date of transplant, for [elective](#) patients registered between 1 April 2021 and 31 March 2024 for an intestine transplant. Any periods of suspension were included in the calculation. Registrations for a re-transplant were included too. [Kaplan-Meier](#) methods were used to calculate median waiting times, where patients who were removed or died on the transplant list were censored at the date of event. If the patient was removed and registered on the liver list on the date of removal, their waiting time is recorded as the time from intestinal registration to liver transplantation, as opposed to censored at the date of removal. Often patients receive a liver transplant on the date of removal from intestinal list. Patients who were still active on the transplant list at time of analysis, 08 July 2026, were censored at that time. Registrations ending in a living donor transplant are excluded.

### Unadjusted survival rates

[Unadjusted patient survival](#) rates were estimated using [Kaplan-Meier](#) methods. Patient survival rates are based on the number of patients transplanted and the number and timing of those that die within the post-transplant period of interest. Patients can be included in this method of analysis irrespective of the length of follow-up recorded. If a patient is alive at the end of the follow-up, then information about the survival of the patient is censored at the time of analysis. Death, irrespective of whether the graft is still functioning or not, is classed as an event.

### Donor offer decline rates

Donor intestine offer decline rates were calculated for each intestine transplant centre. The denominator was equal to the number of intestines offered to them from UK [DBD](#) donors who met the criteria for intestine donation and whose family granted consent for intestine donation. The numerator was equal to the number of intestines each centre declined. Therefore, if a centre received two offers from the same donor for two of their patients and declined it for both, this counts as one offer and one decline; if they accepted it for one of these patients it counts as one offer and one acceptance. The general criteria for intestine donation is donor age less than 60 years and weight less than 90 kg at time of death. However, centres are highly selective when accepting a donor organ which leads to high decline rates. The time period analysed was 1 April 2025 to 31 March 2026.

### Geographical variation analysis

For a given individual who is a resident in a given NHS region, registration to the transplant list is modelled as a Bernoulli trial. At the whole area level, this becomes a Binomial process which can be approximated by a Poisson distribution when rare events are modelled. Transplant counts follow similar assumptions.

To allow for the possibility that, even after allowing for area-specific Poisson rates, area differences remain, introduce an additional multiplicative rate factor which varies from area to area. Postulate a non-parametric distribution for the multiplicative factor, with variance  $\sigma^2$ . If the factor is one for all areas, then area differences are fully explained by the area-specific Poisson rate. If the factor varies with a nonzero variance,  $\sigma^2$ , then we conclude that there are unexplained area differences.

The systematic component of variation (SCV; McPherson et al., N Engl J Med 1982, 307: 1310-4) is the moment estimator of  $\sigma^2$ . Under the null hypothesis of homogeneity across areas, the SCV would be zero. The SCV, therefore, allows us to detect variability across areas beyond that expected by chance; the larger the SCV, the greater the evidence of systematic variation across areas.

A one-sided p-value for the hypothesis that the SCV is greater than zero versus the null hypothesis that the SCV is equal to zero was derived using a parametric bootstrap where data were simulated from the Poisson distribution that would be consistent with the null hypothesis (multiplicative rate factor is equal to one in all areas and  $\sigma^2$  equal to zero). The observed SCV was then compared against this simulated data to calculate the probability that an SCV of at least this size would be observed due to chance if the null hypothesis were true.

10,000 bootstrap samples of size 7 (number of areas) were simulated, where the registration/transplant count in each area was drawn from a Poisson distribution with its expected value being the area-specific expected count (the rate of transplants/registrations in the total population multiplied by the population of the area). The SCV was then calculated in each of the 10,000 samples and a bootstrap p-value for the SCV in the observed data was estimated as:

$$P_{boot} = \frac{1 + \#\{SCV_{sim} \geq SCV_{obs}\}}{10000 + 1}$$

where  $\#\{SCV_{sim} \geq SCV_{obs}\}$  is the number of SCV values in the simulated datasets which are greater than or equal to the SCV in the observed data. This follows the simulation method given in Ibanez et al., BMC Health Services Research, 2009, 9:60. No adjustment was made for area-specific demographic characteristics that may impact the rates of registration to the transplant list and transplantation such as age and sex.

### **A3: Glossary of terms**

#### **Active transplant list**

When a patient is registered for a transplant, they may be registered on what is called the 'active' transplant list. This means that when a donor organ becomes available, the patient is included among those who are matched against the donor to determine whether or not the organ is suitable for them. It may sometimes be necessary to take a patient off the active transplant list, either temporarily or permanently. This may be done, for example, if someone becomes too ill to receive a transplant. The patient is told about the decision to suspend them from the list and is informed whether the suspension is temporary or permanent. If a patient is suspended from the list, they are not included in the matching of any donor organs that become available.

#### **Boxplots**

The length of the box in this plot represents the [inter-quartile range](#). The line inside the box indicates the [median](#) value. The vertical lines issuing from the box are called the whiskers and indicate the range of values that are outside of the inter-quartile range but are close enough not to be considered outliers. The circles that are outside the box indicate the outliers.

#### **Case mix**

The types of patients treated at a unit for a common condition. This can vary across units depending on the facilities available at the unit as well as the types of people in the catchment area of the unit. The definition of what type of patient a person is depends on the patient characteristics that influence the outcome of the treatment.

#### **Confidence interval (CI)**

When an estimate of a quantity such as a survival rate is obtained from data, the value of the estimate depends on the set of patients whose data were used. If, by chance, data from a different set of patients had been used, the value of the estimate may have been different. There is therefore some uncertainty linked with any estimate. A confidence interval is a range of values whose width gives an indication of the uncertainty or precision of an estimate. The number of transplants or patients analysed influences the width of a confidence interval. Smaller data sets tend to lead to wider confidence intervals compared to larger data sets. Estimates from larger data sets are therefore more precise than those from smaller data sets. Confidence intervals are calculated with a stated probability, usually 95%.

#### **Confidence limit**

The upper and lower bounds of a [confidence interval](#).

**Donor after brain death (DBD)**

Donation after brain death (DBD) is a type of donation that takes place following the diagnosis of death using neurological criteria.

**Elective registration**

A patient who is registered to the intestine transplant list as a 'routine' rather than a 'super-urgent' patient. The two groups have a different range of indications for transplantation with markedly different short-term prognoses. Similarly, the process of offering a donor intestine is different for super-urgent and elective registrations, reflecting the difference in risk of death without transplantation for these two patient groups.

**Inter-quartile range (IQR)**

The values between which the middle 50% of the data are distributed. The lower boundary of the IQR is the lower quartile, the upper boundary is the upper quartile. Quartiles divide a rank-ordered data set into four equal parts. The values that divide each part are called the first (or lower), second, and third (or upper) quartiles.

**Kaplan-Meier method**

A method that allows patients with incomplete follow-up information to be included in estimating survival rates. For example, in a cohort for estimating one-year patient survival rates, a patient was followed up for only nine months before they relocated. If we calculated a crude survival estimate using the number of patients who survived for at least a year, this patient would have to be excluded as it is not known if the patient was still alive at one year after transplant. The Kaplan-Meier method allows information about such patients to be used for the length of time that they are followed-up, when this information would otherwise be discarded. Such instances of incomplete follow-up are not uncommon and the Kaplan-Meier method allows the computation of estimates that are more meaningful in these cases.

**Median**

It is a measure of central tendency of a series of observations. The median is the midpoint in a rank-ordered dataset, so that half the data values are larger than the median, and half are smaller.

**Patient survival rate**

The percentage of patients who are still alive (whether the graft is still functioning or not). This is usually specified for a given time point after first transplant. For example, a five-year patient survival rate is the percentage of patients who are still alive five years after their first transplant.

***p* value**

In the context of comparing survival rates across centres, the *p* value is the probability that the differences observed in the rates between centres occurred by chance. As this is a probability, it takes values between 0 and 1. If the *p* value is small, say less than 0.05, this implies that the differences are unlikely to be due to chance and there may be some identifiable cause for these differences. If the *p* value is large, say greater than 0.1, then it is quite likely that any differences seen are due to chance.

**Total preservation time**

The length of time that elapses between the chilling of the intestine after its blood supply has been cut off in the donor and its grafting into the recipient is called total preservation time. Generally, the shorter this time, the better the long-term survival of the recipient.

**Transplant type**

To achieve homogeneous groups in terms of clinical complexity and for simplicity in analysing intestine transplants, the range of transplants in this report are grouped into the following two groups:

Liver containing. This refers to a transplant where the small bowel and liver are transplanted together and may include one or more of: pancreas, kidney, spleen, stomach, abdominal wall, colon.

Non liver containing. This refers to a transplant where the small bowel is transplanted either on its own or with one or more of: kidney, spleen, stomach, abdominal wall, colon. This also refers to a transplant where the small bowel and pancreas are transplanted together and may include one or more of: kidney, spleen, stomach, abdominal wall, colon.

**Unadjusted survival rate**

Unadjusted survival rates are based only on the number of transplants at a given centre and the number and timing of those patients who die within the post-transplant period of interest. In this case, all patients are assumed to be equally likely to die at any given time. However, some centres may have lower unadjusted survival rates than others simply because they tend to undertake transplants that have increased risks of failure. Comparison of unadjusted survival rates across centres and to the national rate is therefore inappropriate.

Prepared by:

Statistics and Clinical Research, NHS Blood and Transplant

Phoebe Salmon  
Rhiannon Taylor

NHSBT Statistics Team are registered with the Office for Statistics Regulation TQV Voluntary Application scheme which demonstrates our commitment to applying the principles of Trustworthiness, Quality and Value from the Code of Practice for Statistics in all our analyses and reports.

