

**International Blood Group  
Reference Laboratory**500 North Bristol Park  
Northway  
Filton  
Bristol  
BS34 7QH**Protein Development  
and Production Unit**

Tel: +44 (0)117 921 7500

Fax: +44 (0)117 912 5796

Website: <http://ibgri.blood.co.uk>Email: [enquiries.IBGRL@nhsbt.nhs.uk](mailto:enquiries.IBGRL@nhsbt.nhs.uk)

<b>Antigen</b>	Glut 1
<b>Clone</b>	BRAC 67
<b>Product Code</b>	9484
<b>Immunoglobulin Class</b>	Rat IgG2c, kappa

**Antigen Description and Distribution**

Glucose transporter 1 (or GLUT1), also known as solute carrier family 2, facilitated glucose transporter member 1 (SLC2A1) is a protein that in humans is encoded by the SLC2A1 gene<sup>1</sup>. GLUT1 facilitates the transport of glucose across the plasma membranes of mammalian cells<sup>2</sup>. The glucose transporter of erythrocytes (called GLUT1 to distinguish it from related glucose transporters in other tissues) is a type III integral protein with 12 hydrophobic segments, each of which is believed to form a membrane-spanning helix. The detailed structure of GLUT1 is not known yet, but one plausible model suggests that the side-by-side assembly of several helices produces a transmembrane channel lined with hydrophilic residues that can hydrogen-bond with glucose as it moves through the channel<sup>3</sup>. GLUT1 is responsible for the low-level of basal glucose uptake required to sustain respiration in all cells. Expression levels of GLUT1 in cell membranes are increased by reduced glucose levels and decreased by increased glucose levels. GLUT1 is also a major receptor for uptake of Vitamin C as well as glucose, especially in non vitamin C producing mammals as part of an adaptation to compensate by participating in a Vitamin C recycling process. In mammals that do produce Vitamin C, GLUT4 is often expressed instead of GLUT1<sup>4</sup>.

**Clone**

BRAC 67 was made in response to a partial purified erythrocyte membrane preparation and has specificity for Glut 1. It has been used to elucidate protein distribution during human erythroblast enucleation<sup>5</sup>. BRAC 67 has been used by immunoblotting of red cell membrane proteins to reveal the membrane alterations in the absence of RHCE<sup>6</sup>.

**References**

1. Mueckler M, Caruso C, Baldwin SA, Panico M, Blench I, Morris HR, Allard WJ, Lienhard GE, Lodish HF (September 1985). "Sequence and structure of a human glucose transporter". *Science* **229** (4717): 941–5.
2. Olson AL, Pessin JE (1996). "Structure, function, and regulation of the mammalian facilitative glucose transporter gene family". *Annu. Rev. Nutr.* **16**: 235–56.
3. Nelson DL, Cox MM (2008). *Lehninger, Principles of Biochemistry*. W. H. Freeman and Company. ISBN 978-0-7167-7108-1.
4. Montel-hagen A, Kinet S, Manel N *et al.* (2008). "Erythrocyte Glut1 Triggers Dehydroascorbic Acid Uptake in Mammals Unable to Synthesize Vitamin C". *Cell* **132** (6): 1039–1048.
5. Bell AJ, Satchwell TJ, Heesom KJ, Hawley BR, Kupzig S, Hazell M, Mushens R, Herman A and Toye AM (2013). Protein Distribution during Human Erythroblast Enucleation *In Vitro*. PLoS ONE Volume **8** (Issue 4) e60300.
6. Joanna F. Flatt, Rozi H. Musa, Yasmin Ayob, Afifah Hassan, Norhanim Asidin, Nurul M. Yahya, Rosalind Mathlouthi, Nicole Thornton, David J. Anstee, Lesley J. Bruce (2012) Study of the D-- phenotype reveals erythrocyte membrane alterations in the absence of RHCE. *BJH* **158**, 262-273.